

Andreas Natsch

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8249356/publications.pdf>

Version: 2024-02-01

89
papers

5,083
citations

101384

36
h-index

88477

70
g-index

95
all docs

95
docs citations

95
times ranked

2785
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Nature-derived epoxy resins: Synthesis, allergenicity, and thermosetting properties of pinoresinol diglycidyl ether. <i>Toxicology and Industrial Health</i> , 2022, 38, 259-269. | 0.6 | 7 |
| 2 | A species specific metabolism leading to male rat reprotoxicity of Cyclamen aldehyde: in vivo and in vitro evaluation. <i>Food and Chemical Toxicology</i> , 2021, 153, 112243. | 1.8 | 1 |
| 3 | Scientific discrepancies in European regulatory proposals on endocrine disruptorsâ€”REACH regulation quo vadis?. <i>Archives of Toxicology</i> , 2021, 95, 3601-3609. | 1.9 | 4 |
| 4 | Assessing Experimental Uncertainty in Defined Approaches: Borderline Ranges for <i>In Chemico</i> and <i>In Vitro</i> Skin Sensitization Methods Determined from Ring Trial Data. <i>Applied in Vitro Toxicology</i> , 2021, 7, 102-111. | 0.6 | 4 |
| 5 | Benzoyl-CoA conjugate accumulation as an initiating event for male reprotoxic effects in the rat? Structureâ€”activity analysis, species specificity, and in vivo relevance. <i>Archives of Toxicology</i> , 2020, 94, 4115-4129. | 1.9 | 5 |
| 6 | Examining Uncertainty in In Vitroâ€”In Vivo Extrapolation Applied in Fish Bioconcentration Models. <i>Environmental Science & Technology</i> , 2020, 54, 9483-9494. | 4.6 | 27 |
| 7 | A New Family of Rigid Dienone Musks Challenges the Perceptive Range of the Human Olfactory Receptor OR5AN1. <i>Synlett</i> , 2020, 31, 972-976. | 1.0 | 8 |
| 8 | Replacing the refinement for skin sensitization testing: Considerations to the implementation of adverse outcome pathway (AOP)-based defined approaches (DA) in OECD guidelines. <i>Regulatory Toxicology and Pharmacology</i> , 2020, 115, 104713. | 1.3 | 32 |
| 9 | PeBiToSens, Φ : A Platform for PBT Screening of Fragrance Ingredients Without Animal Testing. <i>Chimia</i> , 2020, 74, 168. | 0.3 | 3 |
| 10 | The specific biochemistry of human axilla odour formation viewed in an evolutionary context. <i>Philosophical Transactions of the Royal Society B: Biological Sciences</i> , 2020, 375, 20190269. | 1.8 | 26 |
| 11 | A review of substances found positive in 1 of 3 in vitro tests for skin sensitization. <i>Regulatory Toxicology and Pharmacology</i> , 2019, 106, 352-368. | 1.3 | 23 |
| 12 | Exposure source for skin sensitizing hydroperoxides of limonene and linalool remains elusive: An analytical market surveillance. <i>Food and Chemical Toxicology</i> , 2019, 127, 156-162. | 1.8 | 24 |
| 13 | Repeatability and Reproducibility of the RTgill-W1 Cell Line Assay for Predicting Fish Acute Toxicity. <i>Toxicological Sciences</i> , 2019, 169, 353-364. | 1.4 | 52 |
| 14 | Letter to the editor regarding the article by Roberts, 2018. <i>Regulatory Toxicology and Pharmacology</i> , 2019, 102, 115-116. | 1.3 | 1 |
| 15 | Interlaboratory evaluation of methods to quantify skinâ€”sensitizing hydroperoxides of limonene and linalool (Φ): Analysis in cosmetic bases. <i>Flavour and Fragrance Journal</i> , 2018, 33, 322-330. | 1.2 | 4 |
| 16 | Accurate prediction of acute fish toxicity of fragrance chemicals with the RTgillâ€”W1 cell assay. <i>Environmental Toxicology and Chemistry</i> , 2018, 37, 931-941. | 2.2 | 32 |
| 17 | Effect of Fluorination on Skin Sensitization Potential and Fragrant Properties of Cinnamyl Compounds. <i>Chemistry and Biodiversity</i> , 2018, 15, e1800013. | 1.0 | 5 |
| 18 | Deriving a No Expected Sensitization Induction Level for Fragrance Ingredients Without Animal Testing: An Integrated Approach Applied to Specific Case Studies. <i>Toxicological Sciences</i> , 2018, 165, 170-185. | 1.4 | 28 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Interspecies assessment factors and skin sensitization risk assessment. <i>Regulatory Toxicology and Pharmacology</i> , 2018, 97, 186-188. | 1.3 | 11 |
| 20 | Skin Sensitization of Odorant Materials. , 2017, , 89-90. | | 0 |
| 21 | Interlaboratory evaluation of methods to quantify skin sensitizing hydroperoxides potentially formed from linalool and limonene in perfumes. <i>Flavour and Fragrance Journal</i> , 2017, 32, 277-285. | 1.2 | 7 |
| 22 | Reaction Chemistry to Characterize the Molecular Initiating Event in Skin Sensitization: A Journey to Be Continued. <i>Chemical Research in Toxicology</i> , 2017, 30, 315-331. | 1.7 | 18 |
| 23 | p-Alkyl-Benzoyl-CoA Conjugates as Relevant Metabolites of Aromatic Aldehydes With Rat Testicular Toxicityâ€”Studies Leading to the Design of a Safer New Fragrance Chemical. <i>Toxicological Sciences</i> , 2017, 160, 244-255. | 1.4 | 21 |
| 24 | Biochemistry and Genetics of Human Axilla Odor. , 2017, , 123-124. | | 3 |
| 25 | Nrf2 Activation as a Key Event Triggered by Skin Sensitisers: The Development of the Stable KeratinoSens Reporter Gene Assay. <i>ATLA Alternatives To Laboratory Animals</i> , 2016, 44, 443-451. | 0.7 | 23 |
| 26 | Response to the Letter to the Editor Regarding Our Article (Natsch et al., 2015). <i>Chemical Research in Toxicology</i> , 2015, 28, 2082-2084. | 1.7 | 1 |
| 27 | What Makes Us Smell: The Biochemistry of Body Odour and the Design of New Deodorant Ingredients. <i>Chimia</i> , 2015, 69, 414. | 0.3 | 13 |
| 28 | Bayesian integrated testing strategy (ITS) for skin sensitization potency assessment: a decision support system for quantitative weight of evidence and adaptive testing strategy. <i>Archives of Toxicology</i> , 2015, 89, 2355-2383. | 1.9 | 116 |
| 29 | Predicting Skin Sensitizer Potency Based on In Vitro Data from KeratinoSens and Kinetic Peptide Binding: Global Versus Domain-Based Assessment. <i>Toxicological Sciences</i> , 2015, 143, 319-332. | 1.4 | 82 |
| 30 | Oxidative Tryptophan Modification by Terpene- and Squalene-Hydroperoxides and a Possible Link to Cross-Reactions in Diagnostic Tests. <i>Chemical Research in Toxicology</i> , 2015, 28, 1205-1208. | 1.7 | 7 |
| 31 | Assessing skin sensitization hazard in mice and men using non-animal test methods. <i>Regulatory Toxicology and Pharmacology</i> , 2015, 71, 337-351. | 1.3 | 300 |
| 32 | A fast Resazurin-based live viability assay is equivalent to the MTT-test in the KeratinoSens assay. <i>Toxicology in Vitro</i> , 2015, 29, 688-693. | 1.1 | 29 |
| 33 | Dual regulation of skin sensitizer-induced HMOX1 expression by Bach1 and Nrf2: Comparison to regulation of the AKR1C2-ARE element in the KeratinoSens cell line. <i>Toxicology and Applied Pharmacology</i> , 2015, 288, 281-288. | 1.3 | 12 |
| 34 | Reporter cell lines for skin sensitization testing. <i>Archives of Toxicology</i> , 2015, 89, 1645-1668. | 1.9 | 25 |
| 35 | Systematic evaluation of non-animal test methods for skin sensitisation safety assessment. <i>Toxicology in Vitro</i> , 2015, 29, 259-270. | 1.1 | 112 |
| 36 | Epoxyalcohols: Bioactivation and Conjugation Required for Skin Sensitization. <i>Chemical Research in Toxicology</i> , 2014, 27, 1860-1870. | 1.7 | 10 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Detection of potentially skin sensitizing hydroperoxides of linalool in fragranced products. <i>Analytical and Bioanalytical Chemistry</i> , 2014, 406, 6165-6178. | 1.9 | 32 |
| 38 | A human chemosensory modality to detect peptides in the nose?. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2014, 281, 20131678. | 1.2 | 11 |
| 39 | Predicting the Bioconcentration of Fragrance Ingredients by Rainbow Trout Using Measured Rates of in Vitro Intrinsic Clearance. <i>Environmental Science & Technology</i> , 2014, 48, 9486-9495. | 4.6 | 54 |
| 40 | Stability of limonene and monitoring of a hydroperoxide in fragranced products. <i>Flavour and Fragrance Journal</i> , 2014, 29, 277-286. | 1.2 | 19 |
| 41 | Evaluating the performance of integrated approaches for hazard identification of skin sensitizing chemicals. <i>Regulatory Toxicology and Pharmacology</i> , 2014, 69, 371-379. | 1.3 | 78 |
| 42 | A dataset on 145 chemicals tested in alternative assays for skin sensitization undergoing prevalidation. <i>Journal of Applied Toxicology</i> , 2013, 33, 1337-1352. | 1.4 | 251 |
| 43 | Validation of a malodour-forming enzyme as a target for deodorant actives: in vivo testing of a glutamine conjugate targeting a corynebacterial N ^ε -acyl-glutamine-aminoacylase. <i>Flavour and Fragrance Journal</i> , 2013, 28, 262-268. | 1.2 | 1 |
| 44 | Gene expression changes induced by skin sensitizers in the KeratinoSens [®] cell line: Discriminating Nrf2-dependent and Nrf2-independent events. <i>Toxicology in Vitro</i> , 2013, 27, 2225-2232. | 1.1 | 38 |
| 45 | The sensitivity of the KeratinoSens [®] assay to evaluate plant extracts: A pilot study. <i>Toxicology in Vitro</i> , 2013, 27, 1220-1225. | 1.1 | 28 |
| 46 | Bayesian integrated testing strategy to assess skin sensitization potency: from theory to practice. <i>Journal of Applied Toxicology</i> , 2013, 33, 1353-1364. | 1.4 | 92 |
| 47 | Utility of Rat Liver S9 Fractions to Study Skin-Sensitizing Prohaptens in a Modified KeratinoSens Assay. <i>Toxicological Sciences</i> , 2013, 135, 356-368. | 1.4 | 56 |
| 48 | Bacteria and human (mal)-odours. <i>Flavour and Fragrance Journal</i> , 2013, 28, 199-199. | 1.2 | 3 |
| 49 | Persistence of a biocontrol <i>Pseudomonas</i> inoculant as high populations of culturable and non-culturable cells in 200-cm-deep soil profiles. <i>Soil Biology and Biochemistry</i> , 2012, 44, 122-129. | 4.2 | 31 |
| 50 | Analogues of the Epoxy Resin Monomer Diglycidyl Ether of Bisphenol F: Effects on Contact Allergic Potency and Cytotoxicity. <i>Chemical Research in Toxicology</i> , 2012, 25, 2469-2478. | 1.7 | 25 |
| 51 | Chemical Reactivity and Skin Sensitization Potential for Benzaldehydes: Can Schiff Base Formation Explain Everything?. <i>Chemical Research in Toxicology</i> , 2012, 25, 2203-2215. | 1.7 | 37 |
| 52 | Der menschlichen Duftchemie auf der Spur. <i>Chemie in Unserer Zeit</i> , 2012, 46, 110-116. | 0.1 | 3 |
| 53 | Structure-Activity Relationship between the in Vivo Skin Sensitizing Potency of Analogues of Phenyl Glycidyl Ether and the Induction of Nrf2-Dependent Luciferase Activity in the KeratinoSens in Vitro Assay. <i>Chemical Research in Toxicology</i> , 2011, 24, 1312-1318. | 1.7 | 26 |
| 54 | Relating Skin Sensitizing Potency to Chemical Reactivity: Reactive Michael Acceptors Inhibit NF- κ B Signaling and Are Less Sensitizing than S _N Ar- and S _N 2- Reactive Chemicals. <i>Chemical Research in Toxicology</i> , 2011, 24, 2018-2027. | 1.7 | 38 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | The intra- and inter-laboratory reproducibility and predictivity of the KeratinoSens assay to predict skin sensitizers in vitro: Results of a ring-study in five laboratories. <i>Toxicology in Vitro</i> , 2011, 25, 733-744. | 1.1 | 96 |
| 56 | Respiratory sensitization: Advances in assessing the risk of respiratory inflammation and irritation. <i>Toxicology in Vitro</i> , 2011, 25, 1251-1258. | 1.1 | 13 |
| 57 | Evaluating the sensitization potential of surfactants: Integrating data from the local lymph node assay, guinea pig maximization test, and in vitro methods in a weight-of-evidence approach. <i>Regulatory Toxicology and Pharmacology</i> , 2011, 60, 389-400. | 1.3 | 67 |
| 58 | Investigation of odors in the fragrance industry. <i>Chemoecology</i> , 2010, 20, 135-147. | 0.6 | 66 |
| 59 | Lack of Evidence for HLA-Linked Patterns of Odorous Carboxylic Acids Released from Glutamine Conjugates Secreted in the Human Axilla. <i>Journal of Chemical Ecology</i> , 2010, 36, 837-846. | 0.9 | 21 |
| 60 | Performance of a novel keratinocyte-based reporter cell line to screen skin sensitizers in vitro. <i>Toxicology and Applied Pharmacology</i> , 2010, 245, 281-290. | 1.3 | 267 |
| 61 | A Functional ABCC11 Allele Is Essential in the Biochemical Formation of Human Axillary Odor. <i>Journal of Investigative Dermatology</i> , 2010, 130, 529-540. | 0.3 | 137 |
| 62 | The Nrf2-Keap1-ARE Toxicity Pathway as a Cellular Sensor for Skin Sensitizers—Functional Relevance and a Hypothesis on Innate Reactions to Skin Sensitizers. <i>Toxicological Sciences</i> , 2010, 113, 284-292. | 1.4 | 149 |
| 63 | Chemical Basis for the Extreme Skin Sensitization Potency of (E)-4-(Ethoxymethylene)-2-phenyloxazol-5(4H)-one. <i>Chemical Research in Toxicology</i> , 2010, 23, 1913-1920. | 1.7 | 17 |
| 64 | Use of in vitro testing to identify an unexpected skin sensitizing impurity in a commercial product: A case study. <i>Toxicology in Vitro</i> , 2010, 24, 411-416. | 1.1 | 14 |
| 65 | High Throughput Kinetic Profiling Approach for Covalent Binding to Peptides: Application to Skin Sensitization Potency of Michael Acceptor Electrophiles. <i>Chemical Research in Toxicology</i> , 2009, 22, 592-603. | 1.7 | 120 |
| 66 | Filling the Concept with Data: Integrating Data from Different In Vitro and In Silico Assays on Skin Sensitizers to Explore the Battery Approach for Animal-Free Skin Sensitization Testing. <i>Toxicological Sciences</i> , 2009, 107, 106-121. | 1.4 | 106 |
| 67 | Development of a high-throughput keratinocyte-based standard assay to detect skin sensitizers based on ARE-dependent gene activity. <i>Toxicology Letters</i> , 2009, 189, S69. | 0.4 | 1 |
| 68 | Body odour of monozygotic human twins: a common pattern of odorant carboxylic acids released by a bacterial aminoacylase from axilla secretions contributing to an inherited body odour type. <i>Journal of the Royal Society Interface</i> , 2009, 6, 377-392. | 1.5 | 70 |
| 69 | Filling the concept with data: Integrating data from different in vitro and in silico assays on skin sensitizers to explore the battery approach for animal-free skin sensitization testing. <i>Toxicology Letters</i> , 2008, 180, S101. | 0.4 | 0 |
| 70 | The Sequential Action of a Dipeptidase and a β -Lyase Is Required for the Release of the Human Body Odorant 3-Methyl-3-sulfanylhexasan-1-ol from a Secreted Cys-Gly-(S) Conjugate by Corynebacteria. <i>Journal of Biological Chemistry</i> , 2008, 283, 20645-20652. | 1.6 | 57 |
| 71 | Skin Sensitizers Induce Antioxidant Response Element Dependent Genes: Application to the In Vitro Testing of the Sensitization Potential of Chemicals. <i>Toxicological Sciences</i> , 2008, 102, 110-119. | 1.4 | 182 |
| 72 | LC-MS-Based Characterization of the Peptide Reactivity of Chemicals to Improve the In Vitro Prediction of the Skin Sensitization Potential. <i>Toxicological Sciences</i> , 2008, 106, 464-478. | 1.4 | 173 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Chemical Reactivity Measurement and the Predictive Identification of Skin Sensitisers. ATLA Alternatives To Laboratory Animals, 2008, 36, 215-242. | 0.7 | 129 |
| 74 | Utility and limitations of a peptide reactivity assay to predict fragrance allergens in vitro. Toxicology in Vitro, 2007, 21, 1220-1226. | 1.1 | 133 |
| 75 | Biochemistry of Human Axilla Malodor and Chemistry of Deodorant Ingredients. Chimia, 2007, 61, 27-32. | 0.3 | 24 |
| 76 | Fragrance raw materials and essential oils can reduce prostaglandin E2 formation in keratinocytes and reconstituted human epidermis. International Journal of Cosmetic Science, 2007, 29, 369-376. | 1.2 | 3 |
| 77 | Autecology of the biocontrol strain <i>Pseudomonas fluorescens</i> CHA0 in the rhizosphere and inside roots at later stages of plant development. FEMS Microbiology Ecology, 2006, 23, 119-130. | 1.3 | 30 |
| 78 | A Broad Diversity of Volatile Carboxylic Acids, Released by a Bacterial Aminoacylase from Axilla Secretions, as Candidate Molecules for the Determination of Human-Body Odor Type. Chemistry and Biodiversity, 2006, 3, 1-20. | 1.0 | 128 |
| 79 | Isolation of a bacterial enzyme releasing axillary malodor and its use as a screening target for novel deodorant formulations I. International Journal of Cosmetic Science, 2005, 27, 115-122. | 1.2 | 36 |
| 80 | Identification of Odoriferous Sulfanylalkanols in Human Axilla Secretions and Their Formation through Cleavage of Cysteine Precursors by a Cys ₂ S Lyase Isolated from Axilla bacteria. Chemistry and Biodiversity, 2004, 1, 1058-1072. | 1.0 | 99 |
| 81 | A Specific Bacterial Aminoacylase Cleaves Odorant Precursors Secreted in the Human Axilla. Journal of Biological Chemistry, 2003, 278, 5718-5727. | 1.6 | 135 |
| 82 | Cosmopolitan distribution of pHID-containing dicotyledonous crop-associated biocontrol pseudomonads of worldwide origin. FEMS Microbiology Ecology, 2001, 37, 105-116. | 1.3 | 102 |
| 83 | Title is missing!. European Journal of Plant Pathology, 1998, 104, 631-643. | 0.8 | 122 |
| 84 | Impact of <i>Pseudomonas fluorescens</i> strain CHA0 and a derivative with improved biocontrol activity on the culturable resident bacterial community on cucumber roots. FEMS Microbiology Ecology, 1998, 27, 365-380. | 1.3 | 44 |
| 85 | Transport of a biocontrol <i>Pseudomonas fluorescens</i> through 2.5-M deep outdoor lysimeters and survival in the effluent water. Soil Biology and Biochemistry, 1998, 30, 621-631. | 4.2 | 23 |
| 86 | Influence of biocontrol strain <i>Pseudomonas fluorescens</i> CHA0 and its antibiotic overproducing derivative on the diversity of resident root colonizing pseudomonads. FEMS Microbiology Ecology, 1997, 23, 341-352. | 1.3 | 45 |
| 87 | Importance of Preferential Flow and Soil Management in Vertical Transport of a Biocontrol Strain of <i>Pseudomonas fluorescens</i> in Structured Field Soil. Applied and Environmental Microbiology, 1996, 62, 33-40. | 1.4 | 89 |
| 88 | Conservation of the 2,4-diacetylphloroglucinol biosynthesis locus among fluorescent <i>Pseudomonas</i> strains from diverse geographic locations. Applied and Environmental Microbiology, 1996, 62, 552-563. | 1.4 | 270 |
| 89 | Contribution of the Global Regulator Gene <i>gacA</i> to Persistence and Dissemination of <i>Pseudomonas fluorescens</i> Biocontrol Strain CHA0 Introduced into Soil Microcosms. Applied and Environmental Microbiology, 1994, 60, 2553-2560. | 1.4 | 91 |