

Kazuya Fujihara

List of Publications by Citations

Source: <https://exaly.com/author-pdf/825557/kazuya-fujihara-publications-by-citations.pdf>
Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

37 papers	337 citations	10 h-index	16 g-index
47 ext. papers	487 ext. citations	4.8 avg, IF	2.97 L-index

#	Paper	IF	Citations
37	In Search of the Ideal Resistance Training Program to Improve Glycemic Control and its Indication for Patients with Type 2 Diabetes Mellitus: A Systematic Review and Meta-Analysis. <i>Sports Medicine</i> , 2016 , 46, 67-77	10.6	45
36	Carotid artery plaque and LDL-to-HDL cholesterol ratio predict atherosclerotic status in coronary arteries in asymptomatic patients with type 2 diabetes mellitus. <i>Journal of Atherosclerosis and Thrombosis</i> , 2013 , 20, 452-64	4	29
35	High risk of failing eradication of <i>Helicobacter pylori</i> in patients with diabetes: a meta-analysis. <i>Diabetes Research and Clinical Practice</i> , 2014 , 106, 81-7	7.4	28
34	Physical Fitness Tests and Type 2 Diabetes Among Japanese: A Longitudinal Study From the Niigata Wellness Study. <i>Journal of Epidemiology</i> , 2019 , 29, 139-146	3.4	27
33	The Relationship between Diabetic Neuropathy and Sleep Apnea Syndrome: A Meta-Analysis. <i>Sleep Disorders</i> , 2013 , 2013, 150371	1.7	24
32	Comparison of the Framingham risk score, UK Prospective Diabetes Study (UKPDS) Risk Engine, Japanese Atherosclerosis Longitudinal Study-Existing Cohorts Combine (JALS-ECC) and maximum carotid intima-media thickness for predicting coronary artery stenosis in patients with asymptomatic type 2 diabetes. <i>Journal of Atherosclerosis and Thrombosis</i> , 2014 , 21, 799-815	4	23
31	Comparison of baseline characteristics and clinical course in Japanese patients with type 2 diabetes among whom different types of oral hypoglycemic agents were chosen by diabetes specialists as initial monotherapy (JDDM 42). <i>Medicine (United States)</i> , 2017 , 96, e6122	1.8	15
30	Association of <i>Helicobacter pylori</i> infection with glycemic control in patients with diabetes: a meta-analysis. <i>Journal of Diabetes Research</i> , 2014 , 2014, 250620	3.9	15
29	Utility of the triglyceride level for predicting incident diabetes mellitus according to the fasting status and body mass index category: the Ibaraki Prefectural Health Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2014 , 21, 1152-69	4	13
28	Relationship Between Number of Multiple Risk Factors and Coronary Artery Disease Risk With and Without Diabetes Mellitus. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019 , 104, 5084-5090	5.6	12
27	Comparison of clinical characteristics in patients with type 2 diabetes among whom different antihyperglycemic agents were prescribed as monotherapy or combination therapy by diabetes specialists. <i>Journal of Diabetes Investigation</i> , 2016 , 7, 260-9	3.9	10
26	Pulse Pressure is a Stronger Predictor Than Systolic Blood Pressure for Severe Eye Diseases in Diabetes Mellitus. <i>Journal of the American Heart Association</i> , 2019 , 8, e010627	6	9
25	Malondialdehyde-modified LDL-related variables are associated with diabetic kidney disease in type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2018 , 141, 237-243	7.4	8
24	Risk of coronary artery disease according to glucose abnormality status and prior coronary artery disease in Japanese men. <i>Metabolism: Clinical and Experimental</i> , 2019 , 101, 153991	12.7	8
23	Association between all-cause mortality and severity of depressive symptoms in patients with type 2 diabetes: Analysis from the Japan Diabetes Complications Study (JDCS). <i>Journal of Psychosomatic Research</i> , 2017 , 99, 34-39	4.1	7
22	Relationships among cardiorespiratory fitness, muscular fitness, and cardiometabolic risk factors in Japanese adolescents: Niigata screening for and preventing the development of non-communicable disease study-Agano (NICE EVIDENCE Study-Agano) 2. <i>Pediatric Diabetes</i> , 2018 , 19, 593-602	3.6	7
21	Ability of Current Machine Learning Algorithms to Predict and Detect Hypoglycemia in Patients With Diabetes Mellitus: Meta-analysis. <i>JMIR Diabetes</i> , 2021 , 6, e22458	2.7	7

20	Association of increased hepatic insulin clearance and change in serum triglycerides or hydroxybutyrate concentration via the sodium/glucose-cotransporter 2 inhibitor tofogliflozin. <i>Diabetes, Obesity and Metabolism</i> , 2020 , 22, 947-956	6.7	6
19	Accuracy of Japanese claims data in identifying diabetes-related complications. <i>Pharmacoepidemiology and Drug Safety</i> , 2021 , 30, 594-601	2.6	6
18	Higher pulse pressure predicts initiation of dialysis in Japanese patients with diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2019 , 35, e3120	7.5	6
17	Circulating malondialdehyde-modified LDL-related variables and coronary artery stenosis in asymptomatic patients with type 2 diabetes. <i>Journal of Diabetes Research</i> , 2015 , 2015, 507245	3.9	5
16	Development and evaluation of the Japanese version of the Audit of Diabetes-Dependent Quality of Life for patients with diabetes. <i>Diabetology International</i> , 2016 , 7, 384-390	2.3	4
15	Associations of Systolic Blood Pressure and Diastolic Blood Pressure With the Incidence of Coronary Artery Disease or Cerebrovascular Disease According to Glucose Status. <i>Diabetes Care</i> , 2021 , 44, 2124-2131	14.6	4
14	Skipping breakfast, late-night eating and current smoking are associated with medication adherence in Japanese patients with diabetes. <i>Primary Care Diabetes</i> , 2020 , 14, 753-759	2.4	3
13	Physical Fitness and Dyslipidemia Among Japanese: A Cohort Study From the Niigata Wellness Study. <i>Journal of Epidemiology</i> , 2021 , 31, 287-296	3.4	3
12	Association of estimated plasma volume and weight loss after long-term administration and subsequent discontinuation of the sodium-glucose cotransporter-2 inhibitor tofogliflozin. <i>Diabetes, Obesity and Metabolism</i> , 2021 , 23, 1660-1665	6.7	3
11	Brain adaptations of insulin signaling kinases, GLUT 3, p-BADser155 and nitrotyrosine expression in various hypoglycemic models of mice. <i>Neurochemistry International</i> , 2020 , 137, 104745	4.4	2
10	Combination of diabetes mellitus and lack of habitual physical activity is a risk factor for functional disability in Japanese. <i>BMJ Open Diabetes Research and Care</i> , 2020 , 8,	4.5	2
9	Machine Learning Approach to Decision Making for Insulin Initiation in Japanese Patients With Type 2 Diabetes (JDDM 58): Model Development and Validation Study. <i>JMIR Medical Informatics</i> , 2021 , 9, e22148	3.6	2
8	Combined Effects of Energy Intake and Physical Activity on Obesity in Japanese Patients with Type 2 Diabetes (JDDM 50): A Cross-Sectional Study. <i>Diabetes Therapy</i> , 2019 , 10, 1133-1138	3.6	1
7	Lactic Acidosis with Metformin Use in a Patient with Type 1 Diabetes Mellitus. <i>General Medicine</i> , 2013 , 14, 72-75		1
6	Rates and risk factors for amputation in people with diabetes in Japan: a historical cohort study using a nationwide claims database. <i>Journal of Foot and Ankle Research</i> , 2021 , 14, 29	3.2	1
5	Differences in occupational stress by smoking intensity and gender in cross-sectional study of 59 355 Japanese employees using the Brief Job Stress Questionnaire (BJSQ): the Niigata Wellness Study.. <i>BMJ Open</i> , 2022 , 12, e055577	3	1
4	Severity of hypertension as a predictor of initiation of dialysis among study participants with and without diabetes mellitus. <i>Journal of Investigative Medicine</i> , 2021 , 69, 724-729	2.9	0
3	Impact of prior cerebrovascular disease and glucose status on incident cerebrovascular disease in Japanese. <i>Cardiovascular Diabetology</i> , 2021 , 20, 174	8.7	0

2 Clinical Problem Solving: What Causes Tachycardia in a Hip Injury?. *General Medicine*, **2012**, 13, 5-10

1 Weight and cardiometabolic risk among adolescents in Agano city, Japan: NICE EVIDENCE
Study-Agano 1. *Asia Pacific Journal of Clinical Nutrition*, **2020**, 29, 856-866

1