
Kari Furu

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8255967/publications.pdf Version: 2024-02-01



Κλρι Ειιρι

#	Article	IF	CITATIONS
1	The Nordic Countries as a Cohort for Pharmacoepidemiological Research. Basic and Clinical Pharmacology and Toxicology, 2010, 106, 86-94.	2.5	460
2	Trends in attention-deficit hyperactivity disorder medication use: a retrospective observational study using population-based databases. Lancet Psychiatry,the, 2018, 5, 824-835.	7.4	187
3	International trends in antipsychotic use: A study in 16 countries, 2005–2014. European Neuropsychopharmacology, 2017, 27, 1064-1076.	0.7	185
4	Selective serotonin reuptake inhibitors and venlafaxine in early pregnancy and risk of birth defects: population based cohort study and sibling design. BMJ, The, 2015, 350, h1798-h1798.	6.0	152
5	Attention-Deficit/Hyperactivity Disorder in Offspring of Mothers With Inflammatory and Immune System Diseases. Biological Psychiatry, 2017, 81, 452-459.	1.3	141
6	Association Between Methylphenidate and Amphetamine Use in Pregnancy and Risk of Congenital Malformations. JAMA Psychiatry, 2018, 75, 167.	11.0	93
7	Association of Prenatal Exposure to Antiseizure Medication With Risk of Autism and Intellectual Disability. JAMA Neurology, 2022, 79, 672.	9.0	87
8	Use of SSRI and SNRI Antidepressants during Pregnancy: A Population-Based Study from Denmark, Iceland, Norway and Sweden. PLoS ONE, 2015, 10, e0144474.	2.5	84
9	Trends in use of warfarin and direct oral anticoagulants in atrial fibrillation in Norway, 2010 to 2015. European Journal of Clinical Pharmacology, 2017, 73, 1417-1425.	1.9	69
10	β-Blocker Use in Pregnancy and the Risk for Congenital Malformations. Annals of Internal Medicine, 2018, 169, 665.	3.9	65
11	Cancer Risk in Women Treated with Fertility Drugs According to Parity Status—A Registry-based Cohort Study. Cancer Epidemiology Biomarkers and Prevention, 2017, 26, 953-962.	2.5	58
12	Use of anti-asthmatic medications as a proxy for prevalence of asthma in children and adolescents in Norway:a nationwide prescription database analysis. European Journal of Clinical Pharmacology, 2007, 63, 693-698.	1.9	57
13	Cancer risk among insulin users: comparing analogues with human insulin in the CARING five-country cohort study. Diabetologia, 2017, 60, 1691-1703.	6.3	57
14	Trends in prescription drug use during pregnancy and postpartum in Norway, 2005 to 2015. Pharmacoepidemiology and Drug Safety, 2018, 27, 995-1004.	1.9	53
15	High validity of mother-reported use of antiasthmatics among children: a comparison with a population-based prescription database. Journal of Clinical Epidemiology, 2011, 64, 878-884.	5.0	42
16	NORRISK 2: A Norwegian risk model for acute cerebral stroke and myocardial infarction. European Journal of Preventive Cardiology, 2017, 24, 773-782.	1.8	42
17	Drug exposure: inclusion of dispensed drugs before pregnancy may lead to underestimation of risk associations. Journal of Clinical Epidemiology, 2013, 66, 964-972.	5.0	39
18	Use of drugs for ADHD among adults—a multinational study among 15.8 million adults in the Nordic countries. European Journal of Clinical Pharmacology, 2016, 72, 1507-1514.	1.9	37

Kari Furu

#	Article	IF	CITATIONS
19	Antidiabetic medication use during pregnancy: an international utilization study. BMJ Open Diabetes Research and Care, 2019, 7, e000759.	2.8	35
20	ADHD treatment and diagnosis in relation to children's birth month: Nationwide cohort study from Norway. Scandinavian Journal of Public Health, 2017, 45, 343-349.	2.3	34
21	Antipsychotic drug use in pregnancy: A multinational study from ten countries. Schizophrenia Research, 2020, 220, 106-115.	2.0	30
22	Prevalence trends and individual patterns of antiepileptic drug use in pregnancy 2006â€⊋016: A study in the five Nordic countries, United States, and Australia. Pharmacoepidemiology and Drug Safety, 2020, 29, 913-922.	1.9	23
23	Utilization of Stimulants and Atomoxetine for Attentionâ€Deficit/Hyperactivity Disorder among 5.4 Million Children Using Populationâ€Based Longitudinal Data. Basic and Clinical Pharmacology and Toxicology, 2017, 120, 373-379.	2.5	20
24	Neuraminidase inhibitors during pregnancy and risk of adverse neonatal outcomes and congenital malformations: population based European register study. BMJ: British Medical Journal, 2017, 356, j629.	2.3	19
25	Individualâ€based versus aggregate metaâ€analysis in multiâ€database studies of pregnancy outcomes: the Nordic example of selective serotonin reuptake inhibitors and venlafaxine in pregnancy. Pharmacoepidemiology and Drug Safety, 2016, 25, 1160-1169.	1.9	17
26	Oral Fluconazole in Pregnancy and Risk of Stillbirth and Neonatal Death. JAMA - Journal of the American Medical Association, 2018, 319, 2333.	7.4	16
27	Antipsychotic use in pregnancy and risk of attention/deficit-hyperactivity disorder and autism spectrum disorder: a Nordic cohort study. Evidence-Based Mental Health, 2022, 25, 54-62.	4.5	13
28	<p>Use of Antidepressants and Risk of Cutaneous Melanoma: A Prospective Registry-Based Case-Control Study</p> . Clinical Epidemiology, 2020, Volume 12, 193-202.	3.0	12
29	Preterm births and use of medication in early adulthood: a populationâ€based registry study. Pharmacoepidemiology and Drug Safety, 2017, 26, 742-751.	1.9	9
30	Incidence of Malformations After Early Pregnancy Exposure to Modafinil in Sweden and Norway. JAMA - Journal of the American Medical Association, 2020, 324, 895.	7.4	9
31	Prescription patterns of inhaled corticosteroids for preschool children – A Norwegian register study. Pediatric Allergy and Immunology, 2015, 26, 655-661.	2.6	7
32	Interrupted time series analysis to assess changes in prescription filling around conception and implications for exposure misclassification. Pharmacoepidemiology and Drug Safety, 2020, 29, 745-749.	1.9	7
33	Child and Parental Characteristics of Medication Use for Attention-Deficit/Hyperactivity Disorder. Journal of Child and Adolescent Psychopharmacology, 2020, 30, 456-464.	1.3	6
34	Risk factors for stroke and choice of oral anticoagulant in atrial fibrillation. European Journal of Clinical Pharmacology, 2018, 74, 1653-1662.	1.9	5
35	Adjusting for unmeasured confounding using validation data: Simplified twoâ€stage calibration for survival and dichotomous outcomes. Statistics in Medicine, 2019, 38, 2719-2734.	1.6	4
36	Cardiovascular, antidepressant and immunosuppressive drug use in relation to risk of cutaneous melanoma: a protocol for a prospective case–control study. BMJ Open, 2019, 9, e025246.	1.9	4

Kari Furu

#	Article	IF	CITATIONS
37	Dispensing of prescribed analgesics in Norway among young people with foreign-or Norwegian-born parents. Scandinavian Journal of Pain, 2011, 2, 36-44.	1.3	3
38	Effect of the market withdrawal of dextropropoxyphene on use of other prescribed analgesics. Scandinavian Journal of Pain, 2018, 18, 667-674.	1.3	3
39	Prescribed drugs in 27 000 individuals after diagnosis of colorectal cancer: A populationâ€based cohort study. Pharmacoepidemiology and Drug Safety, 2021, 30, 1037-1048.	1.9	2
40	Use of sildenafil and other phosphodiesterase type 5 inhibitors among pregnant women in Scandinavia. Acta Obstetricia Et Gynecologica Scandinavica, 2021, 100, 2111-2118.	2.8	2
41	Validation of Assisted Reproductive Technology in the Medical Birth Registry of Norway Versus the Norwegian Prescription Database. Epidemiology, 2020, 31, 681-686.	2.7	1
42	Use of nonâ€insulin antidiabetic drugs in children and young adults – A Scandinavian drug utilization study from 2010–2019. British Journal of Clinical Pharmacology, 2021, 87, 4470-4475.	2.4	1