

# Akihiko Sekizawa

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/8257947/akihiko-sekizawa-publications-by-year.pdf>

**Version:** 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

111  
papers

2,944  
citations

31  
h-index

50  
g-index

117  
ext. papers

3,327  
ext. citations

3.4  
avg, IF

4.41  
L-index

#	Paper	IF	Citations
111	Antimicrobial Resistance for Genital Infection during Pregnancy in Japan.. <i>Infection and Chemotherapy</i> , <b>2022</b> , 54, 173-175	3.9	
110	Medical Professional Enhancement Using Explainable Artificial Intelligence in Fetal Cardiac Ultrasound Screening.. <i>Biomedicines</i> , <b>2022</b> , 10,	4.8	7
109	Predictive ability of serum advanced glycation end products at 11 to 13 weeks of gestation for early-onset preeclampsia. <i>AJOG Global Reports</i> , <b>2022</b> , 2, 100052		
108	Prenatal Identification of Confined Placental Mosaicism in Pregnant Women with Fetal Growth Restriction. <i>Reproductive Sciences</i> , <b>2021</b> , 1	3	
107	Infective endocarditis due to Streptococcus agalactiae in the puerperal period. <i>Journal of Obstetrics and Gynaecology Research</i> , <b>2021</b> , 47, 2238-2241	1.9	0
106	Tadalafil treatment for preeclampsia (medication in preeclampsia; MIE): a multicenter phase II clinical trial. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , <b>2021</b> , 34, 3709-3715	2	4
105	Accuracy of the FMF Bayes theorem-based model for predicting preeclampsia at 11-13 weeks of gestation in a Japanese population. <i>Hypertension Research</i> , <b>2021</b> , 44, 685-691	4.7	4
104	Postpartum questionnaire survey of women who tested negative in a non-invasive prenatal testing: examining negative emotions towards the test. <i>Journal of Human Genetics</i> , <b>2021</b> , 66, 579-584	4.3	0
103	Retrospective details of false-positive and false-negative results in non-invasive prenatal testing for fetal trisomies 21, 18 and 13. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , <b>2021</b> , 256, 75-81	2.4	7
102	Shadow Estimation for Ultrasound Images Using Auto-Encoding Structures and Synthetic Shadows. <i>Applied Sciences (Switzerland)</i> , <b>2021</b> , 11, 1127	2.6	11
101	The routine use of prophylactic Bakri balloon tamponade contributes to blood loss control in major placenta previa. <i>International Journal of Gynecology and Obstetrics</i> , <b>2021</b> , 154, 508-514	4	3
100	Examination of clinical factors affecting intrauterine microbiota.. <i>Reproduction and Fertility</i> , <b>2021</b> , 2, 1-6	1.1	1
99	Evaluation of the clinical performance of noninvasive prenatal testing at a Japanese laboratory. <i>Journal of Obstetrics and Gynaecology Research</i> , <b>2021</b> , 47, 3437-3446	1.9	0
98	Detection of Cardiac Structural Abnormalities in Fetal Ultrasound Videos Using Deep Learning. <i>Applied Sciences (Switzerland)</i> , <b>2021</b> , 11, 371	2.6	21
97	Assessment of the value of measuring soluble fms-like tyrosine kinase-1 and placental growth factor levels following administration of tadalafil to treat fetal growth restriction.. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , <b>2021</b> , 1-5	2	
96	Next-Generation Sequencing Reveals Downregulation of the Wnt Signaling Pathway in Human Dysmature Cumulus Cells as a Hallmark for Evaluating Oocyte Quality. <i>Reproductive Medicine</i> , <b>2020</b> , 1, 205-215	0.5	1
95	Declined use of cervical ripening balloon did not reduce the incidence of umbilical cord prolapse in Japan. <i>Journal of Obstetrics and Gynaecology Research</i> , <b>2020</b> , 46, 1349-1354	1.9	1

94	Successful monozygotic triplet pregnancy after a single blastocyst transfer following in vitro maturation of oocytes from a woman with polycystic ovary syndrome: a case report. <i>BMC Pregnancy and Childbirth</i> , <b>2020</b> , 20, 57	3.2	3
93	Three-Dimensional Peripheral Bloodstream Model of the Uterus for Laparoscopic Radical Hysterectomy. <i>Journal of Minimally Invasive Gynecology</i> , <b>2020</b> , 27, 1196-1202	2.2	
92	Image Segmentation of the Ventricular Septum in Fetal Cardiac Ultrasound Videos Based on Deep Learning Using Time-Series Information. <i>Biomolecules</i> , <b>2020</b> , 10,	5.9	20
91	Successful Pregnancy in a Case of Behçet Disease after Treatment with Prednisolone. <i>Case Reports in Obstetrics and Gynecology</i> , <b>2020</b> , 2020, 8862651	0.8	
90	Model-Agnostic Method for Thoracic Wall Segmentation in Fetal Ultrasound Videos. <i>Biomolecules</i> , <b>2020</b> , 10,	5.9	13
89	Distribution of PAPP-A and total hCG between 11 and 13 weeks of gestation in Japanese pregnant women. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , <b>2020</b> , 33, 2017-2022	2	
88	Whole-Genome Analysis of Human Papillomavirus Type 16 Prevalent in Japanese Women with or without Cervical Lesions. <i>Viruses</i> , <b>2019</b> , 11,	6.2	3
87	Safety Evaluation of Tadalafil Treatment for Fetuses with Early-Onset Growth Restriction (TADAFER): Results from the Phase II Trial. <i>Journal of Clinical Medicine</i> , <b>2019</b> , 8,	5.1	8
86	Prospective evaluation of screening performance of first-trimester prediction models for preterm preeclampsia in an Asian population. <i>American Journal of Obstetrics and Gynecology</i> , <b>2019</b> , 221, 650.e1-650.e16	6.4	33
85	Classification of factors involved in nonreportable results of noninvasive prenatal testing (NIPT) and prediction of success rate of second NIPT. <i>Prenatal Diagnosis</i> , <b>2019</b> , 39, 100-106	3.2	18
84	Within-Host Variations of Human Papillomavirus Reveal APOBEC Signature Mutagenesis in the Viral Genome. <i>Journal of Virology</i> , <b>2018</b> , 92,	6.6	27
83	Fetal cell-free DNA fraction in maternal plasma for the prediction of hypertensive disorders of pregnancy. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , <b>2018</b> , 224, 165-169	2.4	11
82	Massively parallel sequencing of cell-free DNA in plasma for detecting gynaecological tumour-associated copy number alteration. <i>Scientific Reports</i> , <b>2018</b> , 8, 11205	4.9	13
81	TADAFER II: Tadalafil treatment for fetal growth restriction - a study protocol for a multicenter randomised controlled phase II trial. <i>BMJ Open</i> , <b>2018</b> , 8, e020948	3	5
80	Current status of non-invasive prenatal testing in Japan. <i>Journal of Obstetrics and Gynaecology Research</i> , <b>2017</b> , 43, 1245-1255	1.9	26
79	Evaluation of Second-generation HIFU Systems: Less-invasive Fetal Therapy for TRAP Sequence. <i>The Showa University Journal of Medical Sciences</i> , <b>2017</b> , 29, 241-251	0.1	2
78	Clinical risk factors for poor neonatal outcomes in umbilical cord prolapse. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , <b>2016</b> , 29, 1652-6	2	11
77	Nationwide survey for current clinical status of amniocentesis and maternal serum marker test in Japan. <i>Journal of Human Genetics</i> , <b>2016</b> , 61, 879-884	4.3	3

76	Factors affecting parental decisions to terminate pregnancy in the presence of chromosome abnormalities: a Japanese multicenter study. <i>Prenatal Diagnosis</i> , <b>2016</b> , 36, 1121-1126	3.2	9
75	Obstetric risk factors for umbilical cord prolapse: a nationwide population-based study in Japan. <i>Archives of Gynecology and Obstetrics</i> , <b>2016</b> , 294, 467-72	2.5	10
74	Fetal cell-free DNA fraction in maternal plasma is affected by fetal trisomy. <i>Journal of Human Genetics</i> , <b>2016</b> , 61, 647-52	4.3	40
73	Reference values of focused assessment with sonography for obstetrics (FASO) in low-risk population. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , <b>2016</b> , 29, 3449-53	2	
72	A survey on awareness of genetic counseling for non-invasive prenatal testing: the first year experience in Japan. <i>Journal of Human Genetics</i> , <b>2016</b> , 61, 995-1001	4.3	12
71	The use of balloons for uterine cervical ripening is associated with an increased risk of umbilical cord prolapse: population based questionnaire survey in Japan. <i>BMC Pregnancy and Childbirth</i> , <b>2015</b> , 15, 4	3.2	15
70	How do the trends in the prenatal diagnosis of aneuploidy change after a non-invasive prenatal test becomes available? A Japanese single center study. <i>Journal of Medical Ultrasonics (2001)</i> , <b>2015</b> , 42, 195-8	1.4	3
69	Nationwide demonstration project of next-generation sequencing of cell-free DNA in maternal plasma in Japan: 1-year experience. <i>Prenatal Diagnosis</i> , <b>2015</b> , 35, 331-6	3.2	51
68	Chromosome abnormalities diagnosed in utero: a Japanese study of 28 983 amniotic fluid specimens collected before 22 weeks gestations. <i>Journal of Human Genetics</i> , <b>2015</b> , 60, 133-7	4.3	11
67	Current Status of the Screening of Chlamydia trachomatis Infection Among Japanese Pregnant Women. <i>Journal of Clinical Medicine Research</i> , <b>2015</b> , 7, 582-4	2.9	6
66	Guidelines for obstetrical practice in Japan: Japan Society of Obstetrics and Gynecology (JSOG) and Japan Association of Obstetricians and Gynecologists (JAOG) 2014 edition. <i>Journal of Obstetrics and Gynaecology Research</i> , <b>2014</b> , 40, 1469-99	1.9	220
65	Maternal smoking and placental expression of a panel of genes related to angiogenesis and oxidative stress in early pregnancy. <i>Fetal Diagnosis and Therapy</i> , <b>2014</b> , 35, 289-95	2.4	9
64	Circulating mRNA for the PLAC1 gene as a second trimester marker (14-18 weeks gestation) in the screening for late preeclampsia. <i>Fetal Diagnosis and Therapy</i> , <b>2014</b> , 36, 196-201	2.4	11
63	Fibrin Adhesive Spray Occlusion using a Laparoscope for Intractable Chylous Ascites: Case Report. <i>Japanese Journal of Gynecologic and Obstetric Endoscopy</i> , <b>2014</b> , 30, 188-192	0	1
62	Physiological changes in the pattern of placental gene expression early in the first trimester. <i>Reproductive Sciences</i> , <b>2013</b> , 20, 710-4	3	11
61	Attitudes toward non-invasive prenatal diagnosis among pregnant women and health professionals in Japan. <i>Prenatal Diagnosis</i> , <b>2012</b> , 32, 674-9	3.2	35
60	Higher circulating mRNA levels of placental specific genes in a patient with placenta accreta. <i>Prenatal Diagnosis</i> , <b>2011</b> , 31, 827-9	3.2	8
59	Quantitative RT-PCR gene expression analysis of a laser microdissected placenta: an approach to study preeclampsia. <i>Methods in Molecular Biology</i> , <b>2011</b> , 755, 477-89	1.4	

58	Performance of messenger RNAs circulating in maternal blood in the prediction of preeclampsia at 10-14 weeks. <i>American Journal of Obstetrics and Gynecology</i> , <b>2010</b> , 203, 575.e1-7	6.4	40
57	Expression of angiogenesis-related genes in the cellular component of the blood of preeclamptic women. <i>Reproductive Sciences</i> , <b>2009</b> , 16, 857-64	3	28
56	PP13 mRNA expression in trophoblasts from preeclamptic placentas. <i>Reproductive Sciences</i> , <b>2009</b> , 16, 408-13	3	57
55	Prediction of preeclampsia by analysis of cell-free messenger RNA in maternal plasma. <i>American Journal of Obstetrics and Gynecology</i> , <b>2009</b> , 200, 386.e1-7	6.4	42
54	Cellular mRNA expressions of anti-oxidant factors in the blood of preeclamptic women. <i>Prenatal Diagnosis</i> , <b>2009</b> , 29, 691-6	3.2	36
53	Gene expression in chorionic villous samples at 11 weeks of gestation in women who develop preeclampsia later in pregnancy: implications for screening. <i>Prenatal Diagnosis</i> , <b>2009</b> , 29, 1038-44	3.2	20
52	PP13 mRNA expression in the cellular component of maternal blood as a marker for preeclampsia. <i>Prenatal Diagnosis</i> , <b>2009</b> , 29, 1231-6	3.2	25
51	Gene expression in chorionic villous samples at 11 weeks of gestation from women destined to develop preeclampsia. <i>Prenatal Diagnosis</i> , <b>2008</b> , 28, 956-61	3.2	74
50	Evaluation of physiological alterations of the placenta through analysis of cell-free messenger ribonucleic acid concentrations of angiogenic factors. <i>American Journal of Obstetrics and Gynecology</i> , <b>2008</b> , 198, 124.e1-7	6.4	29
49	Detection and quantification of fetal DNA in maternal plasma by using LightCycler technology. <i>Methods in Molecular Biology</i> , <b>2008</b> , 444, 231-8	1.4	7
48	Cell-free mRNA concentrations of CRH, PLAC1, and selectin-P are increased in the plasma of pregnant women with preeclampsia. <i>Prenatal Diagnosis</i> , <b>2007</b> , 27, 772-7	3.2	62
47	Development of noninvasive fetal DNA diagnosis from nucleated erythrocytes circulating in maternal blood. <i>Prenatal Diagnosis</i> , <b>2007</b> , 27, 846-8	3.2	15
46	Performance of a panel of maternal serum markers in predicting preeclampsia at 11-15 weeks of gestation. <i>Prenatal Diagnosis</i> , <b>2007</b> , 27, 1005-10	3.2	26
45	Recent advances in non-invasive prenatal DNA diagnosis through analysis of maternal blood. <i>Journal of Obstetrics and Gynaecology Research</i> , <b>2007</b> , 33, 747-64		24
44	Cell-free mRNA concentrations of plasminogen activator inhibitor-1 and tissue-type plasminogen activator are increased in the plasma of pregnant women with preeclampsia. <i>Clinical Chemistry</i> , <b>2007</b> , 53, 399-404	5.5	49
43	Placenta-derived, cellular messenger RNA expression in the maternal blood of preeclamptic women. <i>Obstetrics and Gynecology</i> , <b>2007</b> , 110, 1130-6	4.9	27
42	The role of p53 mutation in the carcinomas arising from endometriosis. <i>International Journal of Gynecological Pathology</i> , <b>2007</b> , 26, 345-51	3.2	36
41	Clinical potential for noninvasive prenatal diagnosis through detection of fetal cells in maternal blood. <i>Taiwanese Journal of Obstetrics and Gynecology</i> , <b>2006</b> , 45, 10-20	1.6	11

40	Enrichment of NRBC in maternal blood: a more feasible method for noninvasive prenatal diagnosis. <i>Prenatal Diagnosis</i> , <b>2006</b> , 26, 545-7	3.2	21
39	Quantitative distribution of a panel of circulating mRNA in preeclampsia versus controls. <i>Prenatal Diagnosis</i> , <b>2006</b> , 26, 1115-20	3.2	43
38	Cell-Free Fetal DNA in Plasma of Pregnant Women: Clinical Potential and Origin. <i>Taiwanese Journal of Obstetrics and Gynecology</i> , <b>2005</b> , 44, 116-122	1.6	6
37	Disappearance of steroid hormone dependency during malignant transformation of ovarian clear cell cancer. <i>International Journal of Gynecological Pathology</i> , <b>2005</b> , 24, 369-76	3.2	16
36	Fragmentation of cell-free fetal DNA in plasma and urine of pregnant women. <i>Prenatal Diagnosis</i> , <b>2005</b> , 25, 604-7	3.2	40
35	Lower maternal PLAC1 mRNA in pregnancies complicated with vaginal bleeding (threatened abortion). <i>Clinical Chemistry</i> , <b>2005</b> , 51, 224-7	5.5	23
34	Rapid clearance of mRNA for PLAC1 gene in maternal blood after delivery. <i>Fetal Diagnosis and Therapy</i> , <b>2005</b> , 20, 27-30	2.4	13
33	Fragmentation of Fetal DNA in Maternal Plasma and Urine. <i>The Showa University Journal of Medical Sciences</i> , <b>2005</b> , 17, 81-87	0.1	
32	Proteinuria and hypertension are independent factors affecting fetal DNA values: a retrospective analysis of affected and unaffected patients. <i>Clinical Chemistry</i> , <b>2004</b> , 50, 221-4	5.5	28
31	Malignant transformation of endometriosis: application of laser microdissection for analysis of genetic alterations according to pathological changes. <i>Medical Electron Microscopy: Official Journal of the Clinical Electron Microscopy Society of Japan</i> , <b>2004</b> , 37, 97-100		20
30	K-ras mutation may promote carcinogenesis of endometriosis leading to ovarian clear cell carcinoma. <i>Medical Electron Microscopy: Official Journal of the Clinical Electron Microscopy Society of Japan</i> , <b>2004</b> , 37, 188-92		38
29	Fetal DNA in maternal plasma as a screening variable for preeclampsia. A preliminary nonparametric analysis of detection rate in low-risk nonsymptomatic patients. <i>Prenatal Diagnosis</i> , <b>2004</b> , 24, 83-6	3.2	54
28	Cell-free fetal DNA (SRY locus) concentration in maternal plasma is directly correlated to the time elapsed from the onset of preeclampsia to the collection of blood. <i>Prenatal Diagnosis</i> , <b>2004</b> , 24, 293-7	3.2	25
27	beta-globin DNA in maternal plasma as a molecular marker of pre-eclampsia. <i>Prenatal Diagnosis</i> , <b>2004</b> , 24, 697-700	3.2	30
26	Total cell-free DNA (beta-globin gene) distribution in maternal plasma at the second trimester: a new prospective for preeclampsia screening. <i>Prenatal Diagnosis</i> , <b>2004</b> , 24, 722-6	3.2	52
25	Relationship between Malignant Transformation of Endometriosis and Genetic Alterations of K-ras and Microsatellite Instability. <i>The Showa University Journal of Medical Sciences</i> , <b>2004</b> , 16, 47-54	0.1	
24	Placenta increta: Postpartum monitoring of plasma cell-free fetal DNA. <i>Clinical Chemistry</i> , <b>2003</b> , 49, 1540-5	0.1	30
23	Evaluation of bidirectional transfer of plasma DNA through placenta. <i>Human Genetics</i> , <b>2003</b> , 113, 307-106.3	0.3	42

22	Cell-free fetal DNA in the plasma of pregnant women with severe fetal growth restriction. <i>American Journal of Obstetrics and Gynecology</i> , <b>2003</b> , 188, 480-4	6.4	104
21	p53 mutations and overexpression affect prognosis of ovarian endometrioid cancer but not clear cell cancer. <i>Gynecologic Oncology</i> , <b>2003</b> , 88, 318-25	4.9	86
20	Relationship between severity of hyperemesis gravidarum and fetal DNA concentration in maternal plasma. <i>Clinical Chemistry</i> , <b>2003</b> , 49, 1667-9	5.5	44
19	Testing normality of fetal DNA concentration in maternal plasma at 10-12 completed weeksR gestation: a preliminary approach to a new marker for genetic screening. <i>Prenatal Diagnosis</i> , <b>2002</b> , 22, 148-52	3.2	24
18	Fate of Fetal Nucleated Erythrocytes Circulating in Maternal Blood: Apoptosis Is Induced by Maternal Oxygen Concentration. <i>Clinical Chemistry</i> , <b>2002</b> , 48, 1618-1620	5.5	17
17	Increased Cell-free Fetal DNA in Plasma of Two Women with Invasive Placenta. <i>Clinical Chemistry</i> , <b>2002</b> , 48, 353-354	5.5	112
16	Increased cell-free fetal DNA in plasma of two women with invasive placenta. <i>Clinical Chemistry</i> , <b>2002</b> , 48, 353-4	5.5	21
15	Fate of fetal nucleated erythrocytes circulating in maternal blood: apoptosis is induced by maternal oxygen concentration. <i>Clinical Chemistry</i> , <b>2002</b> , 48, 1618-20	5.5	6
14	Accuracy of Fetal Gender Determination by Analysis of DNA in Maternal Plasma. <i>Clinical Chemistry</i> , <b>2001</b> , 47, 1856-1858	5.5	113
13	Cell-free Fetal DNA Is Increased in Plasma of Women with Hyperemesis Gravidarum. <i>Clinical Chemistry</i> , <b>2001</b> , 47, 2164-2165	5.5	77
12	Comparison of fetal cell recovery from maternal blood using a high density gradient for the initial separation step: 1.090 versus 1.119 g/ml. <i>Prenatal Diagnosis</i> , <b>2000</b> , 20, 281-6	3.2	32
11	Apoptosis in fetal nucleated erythrocytes circulating in maternal blood. <i>Prenatal Diagnosis</i> , <b>2000</b> , 20, 886-9	3.2	89
10	Prenatal DNA diagnosis of a single-gene disorder from maternal plasma. <i>Lancet, The</i> , <b>2000</b> , 356, 1170	4.0	207
9	Female fetal cells in maternal blood: use of DNA polymorphisms to prove origin. <i>Human Genetics</i> , <b>2000</b> , 107, 28-32	6.3	34
8	Female fetal cells in maternal blood: use of DNA polymorphisms to prove origin <b>2000</b> , 107, 28		10
7	Fetal cell recycling: diagnosis of gender and RhD genotype in the same fetal cell retrieved from maternal blood. <i>American Journal of Obstetrics and Gynecology</i> , <b>1999</b> , 181, 1237-42	6.4	29
6	Improvement of fetal cell recovery from maternal blood: suitable density gradient for FACS separation. <i>Fetal Diagnosis and Therapy</i> , <b>1999</b> , 14, 229-33	2.4	33
5	Prenatal diagnosis of ornithine transcarbamylase deficiency by using a single nucleated erythrocyte from maternal blood. <i>Human Genetics</i> , <b>1998</b> , 102, 611-5	6.3	31

4	3Beta-hydroxysteroid dehydrogenase activity in human osteoblast-like cells. <i>Endocrine Journal</i> , <b>1997</b> , 44, 847-53	2.9	14
3	A study of gamma-aminobutyric acid (GABA) in amniotic fluid. <i>Journal of Obstetrics and Gynaecology Research</i> , <b>1997</b> , 23, 471-7		2
2	Prenatal diagnosis of the fetal RhD blood type using a single fetal nucleated erythrocyte from maternal blood. <i>Obstetrics and Gynecology</i> , <b>1996</b> , 87, 501-5	4.9	57
1	A study of monoamine oxidase activity in fetal membranes. <i>Acta Obstetricia Et Gynecologica Scandinavica</i> , <b>1996</b> , 75, 423-7	3.8	4