Xabier Guinda

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8260811/publications.pdf

Version: 2024-02-01

933447 888059 19 465 10 17 citations h-index g-index papers 19 19 19 682 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Changes in the distribution of intertidal macroalgae along a longitudinal gradient in the northern coast of Spain. Marine Environmental Research, 2020, 157, 104930.	2.5	11
2	Distributional shifts of canopy-forming seaweeds from the Atlantic coast of Southern Europe. Biodiversity and Conservation, 2019, 28, 1151-1172.	2.6	73
3	AMBEMAR-DSS: A Decision Support System for the Environmental Impact Assessment of Marine Renewable Energies. , 2018, , .		O
4	Co-location opportunities for renewable energies and aquaculture facilities in the Canary Archipelago. Ocean and Coastal Management, 2018, 166, 62-71.	4.4	28
5	A hierarchical ecological classification system along the NE Atlantic coast: focusing on the local scale (Cantabria, N Spain). European Journal of Phycology, 2017, 52, 75-89.	2.0	4
6	A first approach to stock assessment of the sea urchin <i>Paracentrotus lividus</i> (Lamarck, 1816) in Cantabria (Bay of Biscay). Journal of the Marine Biological Association of the United Kingdom, 2017, 97, 561-570.	0.8	9
7	Essence of the patterns of cover and richness of intertidal hard bottom communities: a pan-European study. Journal of the Marine Biological Association of the United Kingdom, 2017, 97, 525-538.	0.8	10
8	Consistent patterns of spatial variability between NE Atlantic and Mediterranean rocky shores. Journal of the Marine Biological Association of the United Kingdom, 2017, 97, 539-547.	0.8	11
9	Geographic patterns of biodiversity in European coastal marine benthos. Journal of the Marine Biological Association of the United Kingdom, 2017, 97, 507-523.	0.8	14
10	The role of physical variables in biodiversity patterns of intertidal macroalgae along European coasts. Journal of the Marine Biological Association of the United Kingdom, 2017, 97, 549-560.	0.8	10
11	Distribution Patterns of the Gooseneck Barnacle (<i>Pollicipes pollicipes</i> Cantabria Region (N Spain): Exploring Different Population Assessment Methods. Journal of Shellfish Research, 2017, 36, 787-797.	0.9	O
12	A comparison of the degree of implementation of marine biodiversity indicators by European countries in relation to the Marine Strategy Framework Directive (MSFD). Journal of the Marine Biological Association of the United Kingdom, 2015, 95, 1519-1531.	0.8	35
13	Assessing the suitability of the minimum capture size and protection regimes in the gooseneck barnacle shellfishery. Ocean and Coastal Management, 2015, 104, 150-158.	4.4	3
14	The Quality of Rocky Bottoms index (CFR): A validated method for the assessment of macroalgae according to the European Water Framework Directive. Marine Environmental Research, 2014, 102, 3-10.	2.5	15
15	Application of landscape mosaics for the assessment of subtidal macroalgae communities using the CFR index. Deep-Sea Research Part II: Topical Studies in Oceanography, 2014, 106, 207-215.	1.4	8
16	Bloom forming and toxic phytoplankton in transitional and coastal waters of Cantabria region coast (Southeastern Bay of Biscay, Spain). Marine Pollution Bulletin, 2012, 64, 2860-2866.	5.0	14
17	Spatial distribution pattern analysis of subtidal macroalgae assemblages by a non-destructive rapid assessment method. Journal of Sea Research, 2012, 67, 34-43.	1.6	23
18	Macroalgae, a suitable indicator of the ecological status of coastal rocky communities in the NE Atlantic. Ecological Indicators, 2008, 8, 351-359.	6.3	140

#	Article	IF	CITATIONS
19	Comparison of two methods for quality assessment of macroalgae assemblages, under different pollution types. Ecological Indicators, 2008, 8, 743-753.	6.3	57