

Tine W Hansen

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8269915/tine-w-hansen-publications-by-year.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

183
papers

11,170
citations

45
h-index

104
g-index

205
ext. papers

13,053
ext. citations

5.7
avg, IF

6.31
L-index

#	Paper	IF	Citations
183	Acute and Long-Term Treatment With Dapagliflozin and Association With Serum Soluble Urokinase Plasminogen Activator Receptor.. <i>Frontiers in Pharmacology</i> , 2022 , 13, 799915	5.6	0
182	Effect of 26 Weeks of Liraglutide Treatment on Coronary Artery Inflammation in Type 2 Diabetes Quantified by [Cu]Cu-DOTATATE PET/CT: Results from the LIRAFLAME Trial.. <i>Frontiers in Endocrinology</i> , 2021 , 12, 790405	5.7	3
181	Linking Kidney and Cardiovascular Complications in Diabetes-Impact on Prognostication and Treatment: The 2019 Edwin Bierman Award Lecture. <i>Diabetes</i> , 2021 , 70, 39-50	0.9	5
180	Cardiovascular autonomic neuropathy and the impact on progression of diabetic kidney disease in type 1 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2021 , 9,	4.5	3
179	Isolated Diastolic Hypertension in the IDACO Study: An Age-Stratified Analysis Using 24-Hour Ambulatory Blood Pressure Measurements. <i>Hypertension</i> , 2021 , 78, 1222-1231	8.5	3
178	Starting Antihypertensive Drug Treatment With Combination Therapy: Controversies in Hypertension - Con Side of the Argument. <i>Hypertension</i> , 2021 , 77, 788-798	8.5	4
177	Relative and Absolute Risk to Guide the Management of Pulse Pressure, an Age-Related Cardiovascular Risk Factor. <i>American Journal of Hypertension</i> , 2021 , 34, 929-938	2.3	4
176	ASSOCIATION OF FATAL AND NONFATAL CARDIOVASCULAR OUTCOMES WITH 24 HOUR MEAN ARTERIAL PRESSURE. <i>Journal of Hypertension</i> , 2021 , 39, e88	1.9	0
175	PREDICTIVE POWER OF 24-HOUR AMBULATORY PULSE PRESSURE AND ITS COMPONENTS FOR MORTALITY AND CARDIOVASCULAR OUTCOMES IN 11,848 PARTICIPANTS RECRUITED FROM 13 POPULATIONS. <i>Journal of Hypertension</i> , 2021 , 39, e3-e4	1.9	
174	The effect of liraglutide and sitagliptin on oxidative stress in persons with type 2 diabetes. <i>Scientific Reports</i> , 2021 , 11, 10624	4.9	3
173	380-P: Endothelial Glycocalyx Dimensions and Cardiovascular Risk Factors in Type 1 Diabetes. <i>Diabetes</i> , 2021 , 70, 380-P	0.9	
172	430-P: Urinary Proteome and Diabetic Retinopathy in the Direct-Protect 1 and 2 Trials. <i>Diabetes</i> , 2021 , 70, 430-P	0.9	
171	Endothelial glycocalyx and cardio-renal risk factors in type 1 diabetes. <i>PLoS ONE</i> , 2021 , 16, e0254859	3.7	1
170	Urinary peptidome and diabetic retinopathy in the DIRECT-Protect 1 and 2 trials. <i>Diabetic Medicine</i> , 2021 , 38, e14634	3.5	1
169	Visit-to-visit variability of clinical risk markers in relation to long-term complications in type 1 diabetes. <i>Diabetic Medicine</i> , 2021 , 38, e14459	3.5	1
168	Association of Fatal and Nonfatal Cardiovascular Outcomes With 24-Hour Mean Arterial Pressure. <i>Hypertension</i> , 2021 , 77, 39-48	8.5	7
167	Comparison of Natriuretic Peptides as Risk Markers for All-Cause Mortality and Cardiovascular and Renal Complications in Individuals With Type 1 Diabetes. <i>Diabetes Care</i> , 2021 , 44, 595-603	14.6	1

166	Non-invasive assessment of temporal changes in myocardial microvascular function in persons with type 2 diabetes and healthy controls. <i>Diabetic Medicine</i> , 2021 , 38, e14517	3.5	2
165	Plasma trimethylamine N-oxide and its metabolic precursors and risk of mortality, cardiovascular and renal disease in individuals with type 2-diabetes and albuminuria. <i>PLoS ONE</i> , 2021 , 16, e0244402	3.7	8
164	Effect of Liraglutide on Arterial Inflammation Assessed as [F]FDG Uptake in Patients With Type 2 Diabetes: A Randomized, Double-Blind, Placebo-Controlled Trial. <i>Circulation: Cardiovascular Imaging</i> , 2021 , 14, e012174	3.9	8
163	Faecal biomarkers in type 1 diabetes with and without diabetic nephropathy. <i>Scientific Reports</i> , 2021 , 11, 15208	4.9	0
162	Urinary proteomics combined with home blood pressure telemonitoring for health care reform trial: rationale and protocol. <i>Blood Pressure</i> , 2021 , 30, 269-281	1.7	2
161	Effect of Liraglutide on Vascular Inflammation Evaluated by [Cu]DOTATATE. <i>Diagnostics</i> , 2021 , 11,	3.8	2
160	Liraglutide reduces cardiac adipose tissue in type 2 diabetes: A secondary analysis of the LIRAFLAME randomized placebo-controlled trial. <i>Diabetes, Obesity and Metabolism</i> , 2021 , 23, 2651-2659	6.7	1
159	Ceramides and phospholipids are downregulated with liraglutide treatment: results from the LiraFlame randomized controlled trial. <i>BMJ Open Diabetes Research and Care</i> , 2021 , 9,	4.5	3
158	Effect of liraglutide on expression of inflammatory genes in type 2 diabetes. <i>Scientific Reports</i> , 2021 , 11, 18522	4.9	1
157	The Association Between Cardiovascular Autonomic Function and Changes in Kidney and Myocardial Function in Type 2 Diabetes and Healthy Controls.. <i>Frontiers in Endocrinology</i> , 2021 , 12, 780679	5.7	0
156	Early detection of diabetic kidney disease by urinary proteomics and subsequent intervention with spironolactone to delay progression (PRIORITY): a prospective observational study and embedded randomised placebo-controlled trial. <i>Lancet Diabetes and Endocrinology</i> , 2020 , 8, 301-312	18.1	75
155	Relation of cardiac adipose tissue to coronary calcification and myocardial microvascular function in type 1 and type 2 diabetes. <i>Cardiovascular Diabetology</i> , 2020 , 19, 16	8.7	10
154	Lipoprotein(a) and renal function decline, cardiovascular disease and mortality in type 2 diabetes and microalbuminuria. <i>Journal of Diabetes and Its Complications</i> , 2020 , 34, 107593	3.2	2
153	Assessment of the sublingual microcirculation with the GlycoCheck system: Reproducibility and examination conditions. <i>PLoS ONE</i> , 2020 , 15, e0243737	3.7	7
152	Covid-19 Effects on ARTERIAL Stiffness and Vascular Ageing: CARTESIAN Study Rationale and Protocol.. <i>Artery Research</i> , 2020 , 27, 59	2.2	9
151	28-LB: Improved Time in Glucose Range over One Year Is Associated with Reduced Albuminuria in Sensor-Augmented Insulin Pump-Treated Type 1 Diabetes. <i>Diabetes</i> , 2020 , 69, 28-LB	0.9	1
150	1532-P: Investigating Biomarkers of the Immune Response and Tissue Remodeling in Patients with Type 2 Diabetes with Microalbuminuria. <i>Diabetes</i> , 2020 , 69, 1532-P	0.9	
149	Copeptin and renal function decline, cardiovascular events and mortality in type 1 diabetes. <i>Nephrology Dialysis Transplantation</i> , 2020 ,	4.3	2

148	Lancet Commission on Hypertension group position statement on the global improvement of accuracy standards for devices that measure blood pressure. <i>Journal of Hypertension</i> , 2020 , 38, 21-29	1.9	46
147	MR-proANP and incident cardiovascular disease in patients with type 2 diabetes with and without heart failure with preserved ejection fraction. <i>Cardiovascular Diabetology</i> , 2020 , 19, 180	8.7	2
146	Carotid-Femoral Pulse Wave Velocity as a Risk Marker for Development of Complications in Type 1 Diabetes Mellitus. <i>Journal of the American Heart Association</i> , 2020 , 9, e017165	6	6
145	Circulating Metabolites and Lipids Are Associated to Diabetic Retinopathy in Individuals With Type 1 Diabetes. <i>Diabetes</i> , 2020 , 69, 2217-2226	0.9	12
144	Gut microbiota profile and selected plasma metabolites in type 1 diabetes without and with stratification by albuminuria. <i>Diabetologia</i> , 2020 , 63, 2713-2724	10.3	8
143	Improved Time in Range Over 1 Year Is Associated With Reduced Albuminuria in Individuals With Sensor-Augmented Insulin Pump-Treated Type 1 Diabetes. <i>Diabetes Care</i> , 2020 , 43, 2882-2885	14.6	26
142	Pleiotropic effects of liraglutide in patients with type 2 diabetes and moderate renal impairment: Individual effects of treatment. <i>Diabetes, Obesity and Metabolism</i> , 2019 , 21, 1261-1265	6.7	7
141	Utility of Plasma Concentration of Trimethylamine N-Oxide in Predicting Cardiovascular and Renal Complications in Individuals With Type 1 Diabetes. <i>Diabetes Care</i> , 2019 , 42, 1512-1520	14.6	41
140	Soluble Urokinase Plasminogen Activator Receptor Predicts Cardiovascular Events, Kidney Function Decline, and Mortality in Patients With Type 1 Diabetes. <i>Diabetes Care</i> , 2019 , 42, 1112-1119	14.6	19
139	Uric Acid Is an Independent Risk Factor for Decline in Kidney Function, Cardiovascular Events, and Mortality in Patients With Type 1 Diabetes. <i>Diabetes Care</i> , 2019 , 42, 1088-1094	14.6	33
138	A marker of type VI collagen formation (PRO-C6) is associated with higher arterial stiffness in type 1 diabetes. <i>Acta Diabetologica</i> , 2019 , 56, 711-712	3.9	3
137	Cardiac Autonomic Function Is Associated With Myocardial Flow Reserve in Type 1 Diabetes. <i>Diabetes</i> , 2019 , 68, 1277-1286	0.9	7
136	Outcome-Driven Thresholds for Ambulatory Blood Pressure Based on the New American College of Cardiology/American Heart Association Classification of Hypertension. <i>Hypertension</i> , 2019 , 74, 776-783	8.5	13
135	Association of Office and Ambulatory Blood Pressure With Mortality and Cardiovascular Outcomes. <i>JAMA - Journal of the American Medical Association</i> , 2019 , 322, 409-420	27.4	136
134	Markers of Collagen Formation and Degradation Reflect Renal Function and Predict Adverse Outcomes in Patients With Type 1 Diabetes. <i>Diabetes Care</i> , 2019 , 42, 1760-1768	14.6	15
133	Opposing Age-Related Trends in Absolute and Relative Risk of Adverse Health Outcomes Associated With Out-of-Office Blood Pressure. <i>Hypertension</i> , 2019 , 74, 1333-1342	8.5	13
132	Response to Comment on Pilemann-Lyberg et al. Uric Acid Is an Independent Risk Factor for Decline in Kidney Function, Cardiovascular Events, and Mortality in Patients With Type 1 Diabetes. <i>Diabetes Care</i> 2019;42:1088-1094. <i>Diabetes Care</i> , 2019 , 42, e188	14.6	1
131	Prevalence of heart failure and the diagnostic value of MR-proANP in outpatients with type 2 diabetes. <i>Diabetes, Obesity and Metabolism</i> , 2019 , 21, 736-740	6.7	9

130	Myocardial flow reserve assessed by cardiac 82Rb positron emission tomography/computed tomography is associated with albumin excretion in patients with Type 1 diabetes. <i>European Heart Journal Cardiovascular Imaging</i> , 2019 , 20, 796-803	4.1	8
129	Uric acid is not associated with diabetic nephropathy and other complications in type 1 diabetes. <i>Nephrology Dialysis Transplantation</i> , 2019 , 34, 659-666	4.3	7
128	Higher Collagen VI Formation Is Associated With All-Cause Mortality in Patients With Type 2 Diabetes and Microalbuminuria. <i>Diabetes Care</i> , 2018 , 41, 1493-1500	14.6	30
127	Growth differentiation factor-15 and fibroblast growth factor-23 are associated with mortality in type 2 diabetes - An observational follow-up study. <i>PLoS ONE</i> , 2018 , 13, e0196634	3.7	23
126	Cardiovascular autonomic neuropathy and bone metabolism in Type 1 diabetes. <i>Diabetic Medicine</i> , 2018 , 35, 1596-1604	3.5	8
125	Urinary tubular biomarkers as predictors of kidney function decline, cardiovascular events and mortality in microalbuminuric type 2 diabetic patients. <i>Acta Diabetologica</i> , 2018 , 55, 1143-1150	3.9	15
124	A3993 Myocardial flow reserve assessed by Cardiac 82Rb PET/CT is associated with albumin excretion in patients with type 1 diabetes. <i>Journal of Hypertension</i> , 2018 , 36, e147	1.9	
123	A18355 Age- sex- and ethnicity-specific prediction of cardiovascular outcomes by in-office and out-of-the-office blood pressure. <i>Journal of Hypertension</i> , 2018 , 36, e310-e311	1.9	
122	A18029 Outcome-Driven Thresholds for Ambulatory Blood Pressure Based on the New ACC/AHA Classification of Hypertension. <i>Journal of Hypertension</i> , 2018 , 36, e345	1.9	
121	A17393 Systolic and diastolic nocturnal blood pressure dipping differentially predict adverse health outcomes in 8857 untreated participants from 12 populations. <i>Journal of Hypertension</i> , 2018 , 36, e341	1.9	
120	SP418COLLAGEN TYPE III DEGRADATION IS ASSOCIATED WITH DETERIORATION OF KIDNEY FUNCTION IN PATIENTS WITH TYPE 2 DIABETES WITH MICROALBUMINURIA.. <i>Nephrology Dialysis Transplantation</i> , 2018 , 33, i488-i488	4.3	
119	Evidence-based proposal for the number of ambulatory readings required for assessing blood pressure level in research settings: an analysis of the IDACO database. <i>Blood Pressure</i> , 2018 , 27, 341-350	1.7	17
118	Higher Parathyroid Hormone Level Is Associated With Increased Arterial Stiffness in Type 1 Diabetes. <i>Diabetes Care</i> , 2017 , 40, e32-e33	14.6	2
117	Effects of liraglutide on cardiovascular risk biomarkers in patients with type 2 diabetes and albuminuria: A sub-analysis of a randomized, placebo-controlled, double-blind, crossover trial. <i>Diabetes, Obesity and Metabolism</i> , 2017 , 19, 901-905	6.7	31
116	Prevalence, Treatment, and Control Rates of Conventional and Ambulatory Hypertension Across 10 Populations in 3 Continents. <i>Hypertension</i> , 2017 , 70, 50-58	8.5	49
115	Cardiovascular and metabolic effects of metformin in patients with type 1 diabetes (REMOVAL): a double-blind, randomised, placebo-controlled trial. <i>Lancet Diabetes and Endocrinology</i> , 2017 , 5, 597-609	18.1	177
114	Effect of large weight reductions on measured and estimated kidney function. <i>BMC Nephrology</i> , 2017 , 18, 52	2.7	23
113	Symmetric and asymmetric dimethylarginine as risk markers of cardiovascular disease, all-cause mortality and deterioration in kidney function in persons with type 2 diabetes and microalbuminuria. <i>Cardiovascular Diabetology</i> , 2017 , 16, 88	8.7	27

112	Effect of weight reductions on estimated kidney function: Post-hoc analysis of two randomized trials. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 1164-1168	3.2	4
111	Plasma high-sensitivity troponin T predicts end-stage renal disease and cardiovascular and all-cause mortality in patients with type 1 diabetes and diabetic nephropathy. <i>Kidney International</i> , 2017 , 92, 1242-1248	9.9	17
110	Toe-brachial index as a predictor of cardiovascular disease and all-cause mortality in people with type 2 diabetes and microalbuminuria. <i>Diabetologia</i> , 2017 , 60, 1883-1891	10.3	14
109	The effect of liraglutide on renal function: A randomized clinical trial. <i>Diabetes, Obesity and Metabolism</i> , 2017 , 19, 239-247	6.7	55
108	Pleiotropic effects of liraglutide treatment on renal risk factors in type 2 diabetes: Individual effects of treatment. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 162-168	3.2	10
107	Pulse pressure is not an independent predictor of outcome in type 2 diabetes patients with chronic kidney disease and anemia—the Trial to Reduce Cardiovascular Events with Aranesp Therapy (TREAT). <i>Journal of Human Hypertension</i> , 2016 , 30, 46-52	2.6	9
106	High osteoprotegerin is associated with development of foot ulcer in type 1 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2016 , 30, 1603-1608	3.2	5
105	The Cardiovascular Risk of White-Coat Hypertension. <i>Journal of the American College of Cardiology</i> , 2016 , 68, 2033-2043	15.1	96
104	Impaired coronary microcirculation in type 2 diabetic patients is associated with elevated circulating regulatory T cells and reduced number of IL-21R+ T cells. <i>Cardiovascular Diabetology</i> , 2016 , 15, 67	8.7	7
103	Markers of inflammation and endothelial dysfunction are associated with incident cardiovascular disease, all-cause mortality, and progression of coronary calcification in type 2 diabetic patients with microalbuminuria. <i>Journal of Diabetes and Its Complications</i> , 2016 , 30, 248-55	3.2	39
102	Cardiac (82)Rb PET/CT for fast and non-invasive assessment of microvascular function and structure in asymptomatic patients with type 2 diabetes. <i>Diabetologia</i> , 2016 , 59, 371-8	10.3	45
101	Blood Pressure Variability as Elusive Harbinger of Adverse Health Outcomes 2016 , 129-148		
100	Cardiac Autonomic Function Is Associated With the Coronary Microcirculatory Function in Patients With Type 2 Diabetes. <i>Diabetes</i> , 2016 , 65, 3129-38	0.9	16
99	Urinary biomarkers are associated with incident cardiovascular disease, all-cause mortality and deterioration of kidney function in type 2 diabetic patients with microalbuminuria. <i>Diabetologia</i> , 2016 , 59, 1549-1557	10.3	17
98	Relationship Between Two Common Lipoprotein Lipase Variants and the Metabolic Syndrome and Its Individual Components. <i>Metabolic Syndrome and Related Disorders</i> , 2016 , 14, 442-448	2.6	1
97	Global Changes in Food Supply and the Obesity Epidemic. <i>Current Obesity Reports</i> , 2016 , 5, 449-455	8.4	98
96	Improved prognosis of diabetic nephropathy in type 1 diabetes. <i>Kidney International</i> , 2015 , 87, 417-26	9.9	45
95	Prognosis in relation to blood pressure variability: con side of the argument. <i>Hypertension</i> , 2015 , 65, 1170-9; discussion 1179	8.5	28

94	Glucagon-like peptide 1 receptor agonist (GLP-1 RA): long-term effect on kidney function in patients with type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2015 , 29, 670-4	3.2	47
93	Increased plasma concentrations of midregional proatrial natriuretic Peptide is associated with risk of cardiorenal dysfunction in type 1 diabetes. <i>American Journal of Hypertension</i> , 2015 , 28, 772-9	2.3	8
92	Soluble urokinase plasminogen activator receptor levels are elevated and associated with complications in patients with type 1 diabetes. <i>Journal of Internal Medicine</i> , 2015 , 277, 362-371	10.8	50
91	Additive prognostic value of plasma N-terminal pro-brain natriuretic peptide and coronary artery calcification for cardiovascular events and mortality in asymptomatic patients with type 2 diabetes. <i>Cardiovascular Diabetology</i> , 2015 , 14, 59	8.7	28
90	Effect of 4 years subcutaneous insulin infusion treatment on albuminuria, kidney function and HbA1c compared with multiple daily injections: a longitudinal follow-up study. <i>Diabetic Medicine</i> , 2015 , 32, 1445-52	3.5	18
89	Cuff inflations do not affect night-time blood pressure: comparison of 24 h ambulatory blood pressure measured by a cuff and a tonometric device in type 2 diabetes. <i>Blood Pressure Monitoring</i> , 2015 , 20, 369-72	1.3	2
88	Effect of Sensor-Augmented Pump Treatment Versus Multiple Daily Injections on Albuminuria: A 1-Year Randomized Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015 , 100, 4181-8	5.6	15
87	Risk stratification by ambulatory blood pressure monitoring across JNC classes of conventional blood pressure. <i>American Journal of Hypertension</i> , 2014 , 27, 956-65	2.3	37
86	Cardiovascular risk stratification and blood pressure variability on ambulatory and home blood pressure measurement. <i>Current Hypertension Reports</i> , 2014 , 16, 470	4.7	18
85	Setting thresholds to varying blood pressure monitoring intervals differentially affects risk estimates associated with white-coat and masked hypertension in the population. <i>Hypertension</i> , 2014 , 64, 935-42	8.5	108
84	Age-specific differences between conventional and ambulatory daytime blood pressure values. <i>Hypertension</i> , 2014 , 64, 1073-9	8.5	60
83	Ambulatory hypertension subtypes and 24-hour systolic and diastolic blood pressure as distinct outcome predictors in 8341 untreated people recruited from 12 populations. <i>Circulation</i> , 2014 , 130, 466-74	16.7	58
82	Improved survival and renal prognosis of patients with type 2 diabetes and nephropathy with improved control of risk factors. <i>Diabetes Care</i> , 2014 , 37, 1660-7	14.6	50
81	Aortic pulse wave velocity improves cardiovascular event prediction: an individual participant meta-analysis of prospective observational data from 17,635 subjects. <i>Journal of the American College of Cardiology</i> , 2014 , 63, 636-646	15.1	1076
80	Risk Associated with Pulse Pressure on Out-of-Office Blood Pressure Measurement. <i>Pulse</i> , 2014 , 2, 42-51	1.6	4
79	Pulse wave reflection is associated with diabetes duration, albuminuria and cardiovascular disease in type 1 diabetes. <i>Acta Diabetologica</i> , 2014 , 51, 973-80	3.9	6
78	How many measurements are needed to estimate blood pressure variability without loss of prognostic information?. <i>American Journal of Hypertension</i> , 2014 , 27, 46-55	2.3	30
77	Treatment with continuous subcutaneous insulin infusion is associated with lower arterial stiffness. <i>Acta Diabetologica</i> , 2014 , 51, 955-62	3.9	8

76	Ambulatory blood pressure monitoring for risk stratification in obese and non-obese subjects from 10 populations. <i>Journal of Human Hypertension</i> , 2014 , 28, 535-42	2.6	1
75	Outcome-driven thresholds for ambulatory pulse pressure in 9938 participants recruited from 11 populations. <i>Hypertension</i> , 2014 , 63, 229-37	8.5	30
74	Association between albuminuria, atherosclerotic plaques, elevated pulse wave velocity, age, risk category and prognosis in apparently healthy individuals. <i>Journal of Hypertension</i> , 2014 , 32, 1034-41; discussion 1041	1.9	8
73	Central hemodynamics are associated with cardiovascular disease and albuminuria in type 1 diabetes. <i>American Journal of Hypertension</i> , 2014 , 27, 1152-9	2.3	17
72	Blood pressure load does not add to ambulatory blood pressure level for cardiovascular risk stratification. <i>Hypertension</i> , 2014 , 63, 925-33	8.5	30
71	Blood pressure variability in risk stratification: What does it add?. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2014 , 41, 1-8	3	13
70	Outcome-Driven Thresholds for Pulse Pressure on Office and Out-of-the-Office Blood Pressure Measurement 2014 , 447-457		
69	24-hour central aortic systolic pressure and 24-hour central pulse pressure are related to diabetic complications in type 1 diabetes - a cross-sectional study. <i>Cardiovascular Diabetology</i> , 2013 , 12, 122	8.7	22
68	CRP and suPAR are differently related to anthropometry and subclinical organ damage. <i>International Journal of Cardiology</i> , 2013 , 167, 781-5	3.2	82
67	Cardiovascular risk prediction in the general population with use of suPAR, CRP, and Framingham Risk Score. <i>International Journal of Cardiology</i> , 2013 , 167, 2904-11	3.2	98
66	Masked hypertension in diabetes mellitus: treatment implications for clinical practice. <i>Hypertension</i> , 2013 , 61, 964-71	8.5	114
65	Response to Masked hypertension in untreated and treated patients with diabetes mellitus: attractive but questionable interpretations and response to Is masked hypertension related to diabetes mellitus?. <i>Hypertension</i> , 2013 , 62, e23-5	8.5	7
64	Tonometric devices for central aortic systolic pressure measurements in patients with type 1 diabetes: comparison of the BPro and SphygmoCor devices. <i>Blood Pressure Monitoring</i> , 2013 , 18, 156-60	1.3	10
63	Risk stratification by 24-hour ambulatory blood pressure and estimated glomerular filtration rate in 5322 subjects from 11 populations. <i>Hypertension</i> , 2013 , 61, 18-26	8.5	17
62	Double product reflects the predictive power of systolic pressure in the general population: evidence from 9,937 participants. <i>American Journal of Hypertension</i> , 2013 , 26, 665-72	2.3	25
61	White-coat hypertension: new insights from recent studies. <i>Hypertension</i> , 2013 , 62, 982-7	8.5	135
60	The immune marker soluble urokinase plasminogen activator receptor is associated with new-onset diabetes in non-smoking women and men. <i>Diabetic Medicine</i> , 2012 , 29, 479-87	3.5	21
59	Fibrillin-1 genotype and risk of prevalent hypertension: a study in two independent populations. <i>Blood Pressure</i> , 2012 , 21, 273-80	1.7	1

58	Genetic variation in the natriuretic peptide system, circulating natriuretic peptide levels, and blood pressure: an ambulatory blood pressure study. <i>American Journal of Hypertension</i> , 2012 , 25, 1095-100	2.3	10
57	Significance of white-coat hypertension in older persons with isolated systolic hypertension: a meta-analysis using the International Database on Ambulatory Blood Pressure Monitoring in Relation to Cardiovascular Outcomes population. <i>Hypertension</i> , 2012 , 59, 564-71	8.5	146
56	Can ambulatory blood pressure measurements substitute assessment of subclinical cardiovascular damage?. <i>Journal of Hypertension</i> , 2012 , 30, 513-21	1.9	7
55	Thresholds for pulse wave velocity, urine albumin creatinine ratio and left ventricular mass index using SCORE, Framingham and ESH/ESC risk charts. <i>Journal of Hypertension</i> , 2012 , 30, 1928-36	1.9	45
54	Are blood pressure and diabetes additive or synergistic risk factors? Outcome in 8494 subjects randomly recruited from 10 populations. <i>Hypertension Research</i> , 2011 , 34, 714-21	4.7	21
53	1.3 PROGNOSTIC VALUE OF CAROTID-FEMORAL PULSE WAVE VELOCITY FOR CARDIOVASCULAR EVENTS: AN IPD META-ANALYSIS OF PROSPECTIVE OBSERVATIONAL DATA FROM 14 STUDIES INCLUDING 16,358 SUBJECTS. <i>Artery Research</i> , 2011 , 5, 138	2.2	2
52	Soluble urokinase plasminogen activator receptor is associated with subclinical organ damage and cardiovascular events. <i>Atherosclerosis</i> , 2011 , 216, 237-43	3.1	66
51	Short-term blood pressure variability in relation to outcome in the International Database of Ambulatory blood pressure in relation to Cardiovascular Outcome (IDACO). <i>Acta Cardiologica</i> , 2011 , 66, 701-6	0.9	22
50	Ambulatory blood pressure monitoring in 9357 subjects from 11 populations highlights missed opportunities for cardiovascular prevention in women. <i>Hypertension</i> , 2011 , 57, 397-405	8.5	80
49	Predictive role of the nighttime blood pressure. <i>Hypertension</i> , 2011 , 57, 3-10	8.5	398
48	Increased plasma soluble uPAR level is a risk marker of respiratory cancer in initially cancer-free individuals. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2011 , 20, 609-18	4	36
47	Circulating soluble urokinase plasminogen activator receptor predicts cancer, cardiovascular disease, diabetes and mortality in the general population. <i>Journal of Internal Medicine</i> , 2010 , 268, 296-308	10.8	270
46	Urine albumin/creatinine ratio, high sensitivity C-reactive protein and N-terminal pro brain natriuretic peptide--three new cardiovascular risk markers--do they improve risk prediction and influence treatment?. <i>Current Vascular Pharmacology</i> , 2010 , 8, 134-9	3.3	10
45	Prognostic value of reading-to-reading blood pressure variability over 24 hours in 8938 subjects from 11 populations. <i>Hypertension</i> , 2010 , 55, 1049-57	8.5	332
44	Measures of overweight and obesity and risk of cardiovascular disease: a population-based study. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2010 , 17, 486-90		23
43	Prognostic value of the morning blood pressure surge in 5645 subjects from 8 populations. <i>Hypertension</i> , 2010 , 55, 1040-8	8.5	205
42	More information on the reproducibility of the ambulatory arterial stiffness index. <i>American Journal of Hypertension</i> , 2010 , 23, 113-4	2.3	6
41	Tyrosine hydroxylase polymorphism (C-824T) and hypertension: a population-based study. <i>American Journal of Hypertension</i> , 2010 , 23, 1306-11	2.3	7

40	Risk prediction is improved by adding markers of subclinical organ damage to SCORE. <i>European Heart Journal</i> , 2010 , 31, 883-91	9.5	195
39	Determinants of pulse wave velocity in healthy people and in the presence of cardiovascular risk factors: Reestablishing normal and reference values. <i>European Heart Journal</i> , 2010 , 31, 2338-50	9.5	1257
38	Blood pressure variability in relation to outcome in the International Database of Ambulatory blood pressure in relation to Cardiovascular Outcome. <i>Hypertension Research</i> , 2010 , 33, 757-66	4.7	71
37	Relationship between common lipoprotein lipase gene sequence variants, hyperinsulinemia, and risk of ischemic heart disease: A population-based study. <i>Atherosclerosis</i> , 2010 , 211, 506-11	3.1	9
36	From pioneering to implementing automated blood pressure measurement in clinical practice: Thomas Pickering's legacy. <i>Blood Pressure Monitoring</i> , 2010 , 15, 72-81	1.3	4
35	Prognostic value of isolated nocturnal hypertension on ambulatory measurement in 8711 individuals from 10 populations. <i>Journal of Hypertension</i> , 2010 , 28, 2036-45	1.9	263
34	New risk markers may change the HeartScore risk classification significantly in one-fifth of the population. <i>Journal of Human Hypertension</i> , 2009 , 23, 105-12	2.6	8
33	Blood pressure variability remains an elusive predictor of cardiovascular outcome. <i>American Journal of Hypertension</i> , 2009 , 22, 3-4	2.3	23
32	Sensor for vascular compliance and blood pressure 2009 ,		1
31	Which markers of subclinical organ damage to measure in individuals with high normal blood pressure?. <i>Journal of Hypertension</i> , 2009 , 27, 1165-71	1.9	22
30	Risk stratification with the risk chart from the European Society of Hypertension compared with SCORE in the general population. <i>Journal of Hypertension</i> , 2009 , 27, 2351-7	1.9	29
29	Comments on the reproducibility of ambulatory arterial stiffness index and QRS Korotkoff delay index. <i>Journal of Hypertension</i> , 2009 , 27, 436-437	1.9	
28	Diagnostic thresholds for ambulatory blood pressure moving lower: a review based on a meta-analysis-clinical implications. <i>Journal of Clinical Hypertension</i> , 2008 , 10, 377-81	2.3	30
27	Day or night blood pressures for prognosis [AuthorsPreply]. <i>Lancet, The</i> , 2008 , 371, 114-115	4.0	
26	Prognostic value of ambulatory heart rate revisited in 6928 subjects from 6 populations. <i>Hypertension</i> , 2008 , 52, 229-35	8.5	73
25	Independent prognostic value of the ambulatory arterial stiffness index and aortic pulse wave velocity in a general population. <i>Journal of Human Hypertension</i> , 2008 , 22, 214-6	2.6	61
24	C-reactive protein, insulin resistance and risk of cardiovascular disease: a population-based study. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2008 , 15, 594-8		33
23	Spotlights on ambulatory measures of arterial stiffness. <i>American Journal of Hypertension</i> , 2008 , 21, 368-9; author reply 370	2.3	1

22	Determinants of the ambulatory arterial stiffness index in 7604 subjects from 6 populations. <i>Hypertension</i> , 2008 , 52, 1038-44	8.5	34
21	Reproducibility of the ambulatory arterial stiffness index in hypertensive patients. <i>Journal of Hypertension</i> , 2008 , 26, 1993-2000	1.9	16
20	Cardiovascular risk prediction by N-terminal pro brain natriuretic peptide and high sensitivity C-reactive protein is affected by age and sex. <i>Journal of Hypertension</i> , 2008 , 26, 26-34	1.9	32
19	Is blood pressure during the night more predictive of cardiovascular outcome than during the day?. <i>Blood Pressure Monitoring</i> , 2008 , 13, 145-7	1.3	13
18	Insulin resistance, the metabolic syndrome, and risk of incident cardiovascular disease: a population-based study. <i>Journal of the American College of Cardiology</i> , 2007 , 49, 2112-9	15.1	193
17	Diagnostic thresholds for ambulatory blood pressure monitoring based on 10-year cardiovascular risk. <i>Circulation</i> , 2007 , 115, 2145-52	16.7	229
16	N-terminal pro-brain natriuretic peptide, but not high sensitivity C-reactive protein, improves cardiovascular risk prediction in the general population. <i>European Heart Journal</i> , 2007 , 28, 1374-81	9.5	109
15	Prognostic superiority of daytime ambulatory over conventional blood pressure in four populations: a meta-analysis of 7,030 individuals. <i>Journal of Hypertension</i> , 2007 , 25, 1554-64	1.9	266
14	The International Database of Ambulatory Blood Pressure in relation to Cardiovascular Outcome (IDACO): protocol and research perspectives. <i>Blood Pressure Monitoring</i> , 2007 , 12, 255-62	1.3	111
13	Diagnostic thresholds for ambulatory blood pressure monitoring based on 10-year cardiovascular risk. <i>Blood Pressure Monitoring</i> , 2007 , 12, 393-5	1.3	19
12	Added VALUE of an ancillary study on ambulatory blood pressure monitoring. <i>Journal of Hypertension</i> , 2007 , 25, 513-5	1.9	1
11	Cardiovascular outcome in relation to progression to hypertension in the Copenhagen MONICA cohort. <i>American Journal of Hypertension</i> , 2007 , 20, 483-91	2.3	37
10	Prognostic accuracy of day versus night ambulatory blood pressure: a cohort study. <i>Lancet, The</i> , 2007 , 370, 1219-29	40	655
9	Prognostic value of aortic pulse wave velocity as index of arterial stiffness in the general population. <i>Circulation</i> , 2006 , 113, 664-70	16.7	1142
8	Metabolic syndrome, low-density lipoprotein cholesterol, and risk of cardiovascular disease: a population-based study. <i>Atherosclerosis</i> , 2006 , 189, 369-74	3.1	43
7	Ambulatory blood pressure monitoring and risk of cardiovascular disease: a population based study. <i>American Journal of Hypertension</i> , 2006 , 19, 243-50	2.3	178
6	Ambulatory arterial stiffness index predicts stroke in a general population. <i>Journal of Hypertension</i> , 2006 , 24, 2247-53	1.9	114
5	High-sensitivity C-reactive protein is only weakly related to cardiovascular damage after adjustment for traditional cardiovascular risk factors. <i>Journal of Hypertension</i> , 2006 , 24, 655-61	1.9	32

4	Ambulatory blood pressure and mortality: a population-based study. <i>Hypertension</i> , 2005 , 45, 499-504	8.5	245
3	Letter regarding article by Sega et al, "Prognostic value of ambulatory and home blood pressures compared with office blood pressure in the general population". <i>Circulation</i> , 2005 , 112, e244; author reply e245-6	16.7	2
2	N-terminal pro brain natriuretic peptide is inversely related to metabolic cardiovascular risk factors and the metabolic syndrome. <i>Hypertension</i> , 2005 , 46, 660-6	8.5	140
1	Relation between insulin and aortic stiffness: a population-based study. <i>Journal of Human Hypertension</i> , 2004 , 18, 1-7	2.6	47