

Baoqi Fan

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/827011/publications.pdf>

Version: 2024-02-01

15
papers

240
citations

932766

10
h-index

1058022

14
g-index

16
all docs

16
docs citations

16
times ranked

201
citing authors

#	ARTICLE	IF	CITATIONS
1	Shortened Leukocyte Telomere Length Is Associated With Glycemic Progression in Type 2 Diabetes: A Prospective and Mendelian Randomization Analysis. <i>Diabetes Care</i> , 2022, 45, 701-709.	4.3	37
2	Shortened Relative Leukocyte Telomere Length Is Associated With Prevalent and Incident Cardiovascular Complications in Type 2 Diabetes: Analysis From the Hong Kong Diabetes Register. <i>Diabetes Care</i> , 2020, 43, 2257-2265.	4.3	31
3	MicroRNA and Diabetic Complications: A Clinical Perspective. <i>Antioxidants and Redox Signaling</i> , 2018, 29, 1041-1063.	2.5	27
4	Young age at diabetes diagnosis amplifies the effect of diabetes duration on risk of chronic kidney disease: a prospective cohort study. <i>Diabetologia</i> , 2021, 64, 1990-2000.	2.9	22
5	Development of genome-wide polygenic risk scores for lipid traits and clinical applications for dyslipidemia, subclinical atherosclerosis, and diabetes cardiovascular complications among East Asians. <i>Genome Medicine</i> , 2021, 13, 29.	3.6	18
6	Risk associations of long-term HbA1c variability and obesity on cancer events and cancer-specific death in 15,286 patients with diabetes - A prospective cohort study. <i>The Lancet Regional Health - Western Pacific</i> , 2022, 18, 100315.	1.3	13
7	Risk Associations of Glycemic Burden and Obesity With Liver Cancerâ€”A 10â€”Year Analysis of 15,280 Patients With Type 2 Diabetes. <i>Hepatology Communications</i> , 2022, 6, 1350-1360.	2.0	13
8	Detection of increased serum miR-122-5p and miR-455-3p levels before the clinical diagnosis of liver cancer in people with type 2 diabetes. <i>Scientific Reports</i> , 2021, 11, 23756.	1.6	13
9	Nonalbuminuric Diabetic Kidney Disease and Risk of All-Cause Mortality and Cardiovascular and Kidney Outcomes in Type 2 Diabetes: Findings From the Hong Kong Diabetes Biobank. <i>American Journal of Kidney Diseases</i> , 2022, 80, 196-206.e1.	2.1	12
10	Associations of the HOMA2â€”B and HOMA2â€”IR with progression to diabetes and glycaemic deterioration in young and middleâ€”aged Chinese. <i>Diabetes/Metabolism Research and Reviews</i> , 2022, 38, e3525.	1.7	12
11	Relative leukocyte telomere length is associated with incident end-stage kidney disease and rapid decline of kidney function in type 2 diabetes: analysis from the Hong Kong Diabetes Register. <i>Diabetologia</i> , 2022, 65, 375-386.	2.9	11
12	Pancreatic Sirtuin 3 Deficiency Promotes Hepatic Steatosis by Enhancing 5-Hydroxytryptamine Synthesis in Mice With Diet-Induced Obesity. <i>Diabetes</i> , 2021, 70, 119-131.	0.3	10
13	Shortened relative leukocyte telomere length is associated with all-cause mortality in type 2 diabetes- analysis from the Hong Kong Diabetes Register. <i>Diabetes Research and Clinical Practice</i> , 2021, 173, 108649.	1.1	10
14	Time-varying risk associations of renin angiotensin system inhibitors with pneumonia and related deaths in a cohort of 252,616 patients with diabetes (2002â€”2019). <i>Diabetes Research and Clinical Practice</i> , 2022, 185, 109233.	1.1	6
15	<i>CYP2C19</i> Loss-of-function Polymorphisms are Associated with Reduced Risk of Sulfonylurea Treatment Failure in Chinese Patients with Type 2 Diabetes. <i>Clinical Pharmacology and Therapeutics</i> , 2022, 111, 461-469.	2.3	5