

Ankur Pandey

List of Publications by Citations

Source: <https://exaly.com/author-pdf/8273238/ankur-pandey-publications-by-citations.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

23
papers

175
citations

9
h-index

11
g-index

24
ext. papers

253
ext. citations

7.9
avg, IF

2.7
L-index

#	Paper	IF	Citations
23	Optimized energy of spectral CT for infarct imaging: Experimental validation with human validation. <i>Journal of Cardiovascular Computed Tomography</i> , 2017 , 11, 171-178	2.8	15
22	Prediction of post-TACE necrosis of hepatocellular carcinoma using volumetric enhancement on MRI and volumetric oil deposition on CT, with pathological correlation. <i>European Radiology</i> , 2018 , 28, 3032-3040	8	15
21	Accuracy of apparent diffusion coefficient in differentiating pancreatic neuroendocrine tumour from intrapancreatic accessory spleen. <i>European Radiology</i> , 2018 , 28, 1560-1567	8	15
20	Follow-up of Incidentally Detected Pancreatic Cystic Neoplasms: Do Baseline MRI and CT Features Predict Cyst Growth?. <i>Radiology</i> , 2019 , 292, 647-654	20.5	13
19	Comparison between ROI-based and volumetric measurements in quantifying heterogeneity of liver stiffness using MR elastography. <i>European Radiology</i> , 2020 , 30, 1609-1615	8	13
18	Unresectable Intrahepatic Cholangiocarcinoma: Multiparametric MR Imaging to Predict Patient Survival. <i>Radiology</i> , 2018 , 288, 109-117	20.5	12
17	Utility of volumetric contrast-enhanced and diffusion-weighted MRI in differentiating between common primary hypervascular liver tumors. <i>Journal of Magnetic Resonance Imaging</i> , 2018 , 48, 1080-1090	5.6	11
16	Updates in hepatic oncology imaging. <i>Surgical Oncology</i> , 2017 , 26, 195-206	2.5	10
15	Role of baseline volumetric functional MRI in predicting histopathologic grade and patients survival in hepatocellular carcinoma. <i>European Radiology</i> , 2020 , 30, 3748-3758	8	10
14	Prognostic value of baseline volumetric multiparametric MR imaging in neuroendocrine liver metastases treated with transarterial chemoembolization. <i>European Radiology</i> , 2019 , 29, 5160-5171	8	7
13	Added value of apparent diffusion coefficient in distinguishing between serous and mucin-producing pancreatic cystic neoplasms. <i>European Radiology</i> , 2019 , 29, 4660-4669	8	7
12	Baseline Volumetric Multiparametric MRI: Can It Be Used to Predict Survival in Patients with Unresectable Intrahepatic Cholangiocarcinoma Undergoing Transcatheter Arterial Chemoembolization?. <i>Radiology</i> , 2018 , 289, 843-853	20.5	7
11	Baseline 3D-ADC outperforms 2D-ADC in predicting response to treatment in patients with colorectal liver metastases. <i>European Radiology</i> , 2020 , 30, 291-300	8	7
10	Correlation between quantitative liver and spleen volumes and disease severity in primary sclerosing cholangitis as determined by Mayo risk score. <i>European Journal of Radiology</i> , 2018 , 108, 254-260	4.7	6
9	Are pancreatic IPMN volumes measured on MRI images more reproducible than diameters? An assessment in a large single-institution cohort. <i>European Radiology</i> , 2018 , 28, 2790-2800	8	5
8	Cross-sectional imaging in patients with primary sclerosing cholangitis: Single time-point liver or spleen volume is associated with survival. <i>European Journal of Radiology</i> , 2020 , 132, 109331	4.7	4
7	Role of tumor margin and ADC change in defining the need for additional treatments after the first TACE in patients with unresectable HCC. <i>European Journal of Radiology</i> , 2020 , 133, 109389	4.7	4

6	Semi-quantitative visual assessment of hepatic tumor burden can reliably predict survival in neuroendocrine liver metastases treated with transarterial chemoembolization. <i>European Radiology</i> , 2019 , 29, 5804-5812	8	3
5	Assessing the Non-tumorous Liver: Implications for Patient Management and Surgical Therapy. <i>Journal of Gastrointestinal Surgery</i> , 2018 , 22, 344-360	3-3	3
4	Post-TACE changes in ADC histogram predict overall and transplant-free survival in patients with well-defined HCC: a retrospective cohort with up to 10 years follow-up. <i>European Radiology</i> , 2021 , 31, 1378-1390	8	3
3	Does the change in volumetric functional MR metrics post-TACE predict histopathologic grading of hepatocellular carcinoma?. <i>European Radiology</i> , 2020 , 30, 6709-6720	8	2
2	Role of volumetric multiparametric MRI in distinguishing between intraductal papillary mucinous neoplasms and serous cystadenoma. <i>Abdominal Radiology</i> , 2021 , 46, 1629-1639	3	2
1	Integrating baseline MR imaging biomarkers into BCLC and CLIP improves overall survival prediction of patients with hepatocellular carcinoma (HCC). <i>European Radiology</i> , 2021 , 31, 1630-1641	8	1