

Haibiao Xie

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/827756/publications.pdf>

Version: 2024-02-01

17
papers

371
citations

1040056

9
h-index

888059

17
g-index

17
all docs

17
docs citations

17
times ranked

585
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Clinical characteristics and risk factors for survival in affected offspring of von Hippel-Lindau disease patients. <i>Journal of Medical Genetics</i> , 2022, 59, 951-956. | 3.2 | 5 |
| 2 | TBK1 Is a Synthetic Lethal Target in Cancer with <i>VHL</i> Loss. <i>Cancer Discovery</i> , 2020, 10, 460-475. | 9.4 | 63 |
| 3 | Genome-wide Screening Identifies SFMBT1 as an Oncogenic Driver in Cancer with VHL Loss. <i>Molecular Cell</i> , 2020, 77, 1294-1306.e5. | 9.7 | 41 |
| 4 | A systematic review and meta-analysis of long noncoding RNA linc-UBC1 expression and prognosis and clinicopathological phenotypes in human cancers. <i>Artificial Cells, Nanomedicine and Biotechnology</i> , 2020, 48, 875-884. | 2.8 | 1 |
| 5 | Novel genetic characterisation and phenotype correlation in von Hippel-Lindau (VHL) disease based on the Elongin C binding site: a large retrospective study. <i>Journal of Medical Genetics</i> , 2020, 57, 744-751. | 3.2 | 1 |
| 6 | A revolutionary tool: CRISPR technology plays an important role in construction of intelligentized gene circuits. <i>Cell Proliferation</i> , 2019, 52, e12552. | 5.3 | 7 |
| 7 | Synthetic artificial "long non-coding RNAs" targeting oncogenic microRNAs and transcriptional factors inhibit malignant phenotypes of bladder cancer cells. <i>Cancer Letters</i> , 2018, 422, 94-106. | 7.2 | 6 |
| 8 | Lentivirus-mediated shRNA targeting <i>MUTYH</i> ; inhibits malignant phenotypes of bladder cancer SW780 cells. <i>OncoTargets and Therapy</i> , 2018, Volume 11, 6101-6109. | 2.0 | 4 |
| 9 | Oestrogen promotes tumorigenesis of bladder cancer by inducing the enhancer RNA "eGREB1". <i>Journal of Cellular and Molecular Medicine</i> , 2018, 22, 5919-5927. | 3.6 | 15 |
| 10 | High expression of enhancer RNA MARC1 or its activation by DHT is associated with the malignant behavior in bladder cancer. <i>Experimental Cell Research</i> , 2018, 370, 303-311. | 2.6 | 7 |
| 11 | Synthesizing a Genetic Sensor Based on CRISPR-Cas9 for Specifically Killing p53-Deficient Cancer Cells. <i>ACS Synthetic Biology</i> , 2018, 7, 1798-1807. | 3.8 | 24 |
| 12 | Long non-coding RNA CRNDE in cancer prognosis: Review and meta-analysis. <i>Clinica Chimica Acta</i> , 2018, 485, 262-271. | 1.1 | 38 |
| 13 | Tetracycline-controllable artificial microRNA-HOTAIR + EZH2 suppressed the progression of bladder cancer cells. <i>Molecular BioSystems</i> , 2017, 13, 1597-1607. | 2.9 | 12 |
| 14 | Colon cancer associated transcripts in human cancers. <i>Biomedicine and Pharmacotherapy</i> , 2017, 94, 531-540. | 5.6 | 32 |
| 15 | SPRY4-IT1: A novel oncogenic long non-coding RNA in human cancers. <i>Tumor Biology</i> , 2017, 39, 101042831771140. | 1.8 | 34 |
| 16 | LncRNA MALAT1 Inhibits Apoptosis and Promotes Invasion by Antagonizing miR-125b in Bladder Cancer Cells. <i>Journal of Cancer</i> , 2017, 8, 3803-3811. | 2.5 | 79 |
| 17 | Role of nuclear paraspeckle assembly transcript 1 as a common molecular marker for prognosis in various cancers. <i>Minerva Medica</i> , 2017, 108, 477-479. | 0.9 | 2 |