

Emmanuelle Lipka

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8284238/publications.pdf>

Version: 2024-02-01

66
papers

864
citations

566801

15
h-index

552369

26
g-index

67
all docs

67
docs citations

67
times ranked

981
citing authors

#	ARTICLE	IF	CITATIONS
1	Comparison of enhanced fluidity liquid chromatography with liquid chromatography for enantioseparation of selected l^3 -lactam derivatives. <i>Journal of Chromatography Open</i> , 2022, 2, 100026.	0.8	1
2	Contribution of supercritical fluid chromatography to serially coupling columns for chiral and achiral separations. <i>TrAC - Trends in Analytical Chemistry</i> , 2022, 149, 116563.	5.8	2
3	Supercritical fluid chromatography for separation of chiral planar metallocenes. <i>Journal of Chromatography A</i> , 2022, 1674, 463115.	1.8	4
4	Separation of planar chiral ferrocenes by capillary electrokinetic chromatography and liquid chromatography. <i>Journal of Chromatography A</i> , 2022, 1677, 463306.	1.8	6
5	Optimization of Detection of Native Amino Acids with Evaporative Light Scattering Detector in Chiral Supercritical Fluid Chromatography. <i>Chromatographia</i> , 2021, 84, 179-185.	0.7	4
6	Electrophoretic separation of multiple chiral center analyte with a three cyclodextrins mixture. <i>Electrophoresis</i> , 2021, 42, 1810-1817.	1.3	3
7	Recent developments in separation methods for enantiomeric ratio determination of amino acids specifically involved in cataract and Alzheimer's disease. <i>TrAC - Trends in Analytical Chemistry</i> , 2021, 141, 116287.	5.8	11
8	Supercritical fluid chromatography and liquid chromatography for isomeric separation of a multiple chiral centers analyte. <i>Journal of Chromatography A</i> , 2021, 1651, 462270.	1.8	10
9	Carbon dioxide transformation as a green alternative to phosgene and chloroformates: N-carboxyalkylation of lactams and analogues. <i>Journal of CO2 Utilization</i> , 2021, 54, 101782.	3.3	1
10	Pyroglutamide-Based P2X7 Receptor Antagonists Targeting Inflammatory Bowel Disease. <i>Journal of Medicinal Chemistry</i> , 2020, 63, 2074-2094.	2.9	24
11	Productivity and solvent waste in supercritical fluid chromatography for preparative chiral separations: a guide for a convenient strategy. <i>Journal of Chromatography A</i> , 2020, 1610, 460549.	1.8	10
12	Recent developments in preparative-scale supercritical fluid- and liquid chromatography for chiral separations. <i>TrAC - Trends in Analytical Chemistry</i> , 2020, 133, 116090.	5.8	27
13	Analytical and preparative enantioseparations in supercritical fluid chromatography using different brands of immobilized cellulose tris (3,5-dichlorophenylcarbamate) columns: Some differences. <i>Journal of Chromatography A</i> , 2020, 1622, 461125.	1.8	8
14	Comparison of dimethylated and methylchlorinated amylose stationary phases, coated and covalently immobilized on silica, for the separation of some chiral compounds in supercritical fluid chromatography. <i>Journal of Chromatography A</i> , 2020, 1621, 461053.	1.8	11
15	Applications of Chiral Supercritical Fluid Chromatography. <i>Methods in Molecular Biology</i> , 2019, 1985, 303-319.	0.4	0
16	Separations of antifungal compounds in capillary electrophoresis with two anionic cyclodextrins. <i>Electrophoresis</i> , 2019, 40, 1986-1991.	1.3	2
17	Cesium salts as superior catalysts for solvent-free modifications of biosourced pterolactam. <i>Molecular Catalysis</i> , 2019, 470, 32-39.	1.0	5
18	Performance comparison of chlorinated chiral stationary phases in supercritical fluid chromatography for separation of selected pyrrolidone derivatives. <i>Journal of Pharmaceutical Analysis</i> , 2019, 9, 248-253.	2.4	4

#	ARTICLE	IF	CITATIONS
19	Separation of enantiomers of native amino acids with polysaccharide-based chiral columns in supercritical fluid chromatography. <i>Journal of Chromatography A</i> , 2019, 1585, 207-212.	1.8	53
20	Application of tandem coupling of columns in supercritical fluid chromatography for stereoisomeric separation: Optimization and simulation. <i>Journal of Chromatography A</i> , 2019, 1588, 115-126.	1.8	17
21	Supercritical fluid chromatography versus high performance liquid chromatography for enantiomeric and diastereoisomeric separations on coated polysaccharides-based stationary phases: Application to dihydropyridone derivatives. <i>Journal of Chromatography A</i> , 2018, 1549, 39-50.	1.8	17
22	Single and dual cyclodextrins systems for the enantiomeric and diastereoisomeric separations of structurally related dihydropyridone analogues. <i>Electrophoresis</i> , 2017, 38, 1922-1931.	1.3	10
23	Small scale separation of isoxazole structurally related analogues by chiral supercritical fluid chromatography. <i>Journal of Chromatography A</i> , 2017, 1505, 106-113.	1.8	13
24	Development and validation of a reversed-phase HPLC method for the quantification of paclitaxel in different PLGA nanocarriers. <i>Electrophoresis</i> , 2017, 38, 2536-2541.	1.3	8
25	On the discovery of new potent human farnesyltransferase inhibitors: emerging pyroglutamic derivatives. <i>Organic and Biomolecular Chemistry</i> , 2017, 15, 8110-8118.	1.5	13
26	Discovery of highly functionalized scaffolds: Pyrroloimidazolediones as P2X7 receptor antagonists. <i>Tetrahedron</i> , 2017, 73, 5327-5336.	1.0	8
27	Supercritical fluid chromatography approach for a sustainable manufacture of new stereoisomeric anticancer agent. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2017, 145, 845-853.	1.4	6
28	1,3,5-Oxadiazine Framework by Oxygen vs. Nitrogen Trapping of an Acyliminium Ion Derived from N,O-bis(trimethylsilyl) Pyroglutamic Acid. <i>ChemistrySelect</i> , 2017, 2, 10654-10660.	0.7	5
29	Exploring chiral separation of 3-carboxamido-5-aryl isoxazole derivatives by supercritical fluid chromatography on amylose and cellulose tris dimethyl- and chloromethyl phenylcarbamate polysaccharide based stationary phases. <i>Journal of Chromatography A</i> , 2016, 1467, 473-481.	1.8	7
30	Preparative supercritical fluid chromatography: A powerful tool for chiral separations. <i>Journal of Chromatography A</i> , 2016, 1467, 33-55.	1.8	153
31	Impact of Functional Groups on the Copper-Initiated N-Arylation of 5-Functionalized Pyrrolidin-2-ones and Their Vinylogues. <i>Synthesis</i> , 2016, 48, 2226-2244.	1.2	9
32	ZrCl ₄ as a new catalyst for ester amidation: an efficient synthesis of h-P2X7R antagonists. <i>Tetrahedron Letters</i> , 2016, 57, 1165-1170.	0.7	16
33	Design of experiments for enantiomeric separation in supercritical fluid chromatography. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2016, 120, 297-305.	1.4	20
34	Evolution of availability of curcumin inside poly-lactic-co-glycolic acid nanoparticles: impact on antioxidant and antinitrosant properties. <i>International Journal of Nanomedicine</i> , 2015, 10, 5355.	3.3	15
35	Evaluation and comparison of three different separation techniques for analysis of retroamide enantiomers and their biological evaluation against h-P2X7 receptor. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2015, 986-987, 35-43.	1.2	4
36	Quantitative analysis of drugs in biological matrices by HPLC hyphenated to fluorescence detection. <i>Bioanalysis</i> , 2015, 7, 743-762.	0.6	9

#	ARTICLE	IF	CITATIONS
37	Enantioseparation and thermodynamic study of naphthalene derivatives, new melatonergic agonists, on coated amylose [<i>tris</i> (<i>S</i>)-phenylethylcarbamate] stationary phase. Transposition to preparative scale. <i>Biomedical Chromatography</i> , 2015, 29, 689-697.	0.8	3
38	Involvement of the P2X7 Purinergic Receptor in Inflammation: An Update of Antagonists Series Since 2009 and their Promising Therapeutic Potential. <i>Current Medicinal Chemistry</i> , 2015, 22, 713-729.	1.2	43
39	Development of HPLC/fluorescence detection method for chiral resolution of dansylated benzimidazoles derivatives. <i>Biomedical Chromatography</i> , 2014, 28, 4-9.	0.8	3
40	NMR studies of interactions of new CB2 cannabinoid receptor ligands with cyclodextrins hosts. Correlation with micellar electrokinetic chromatography and reversed phase high performance liquid chromatography. <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 2014, 78, 265-274.	0.9	0
41	Antagonists of the P2X7 receptor: Mechanism of enantioselective recognition using highly sulfated and sulfobutylether cyclodextrins by capillary electrokinetic chromatography. <i>Electrophoresis</i> , 2014, 35, 2892-2899.	1.3	9
42	Enantioseparation of pyroglutamide derivatives on polysaccharide based chiral stationary phases by high-performance liquid chromatography and supercritical fluid chromatography: A comparative study. <i>Journal of Chromatography A</i> , 2014, 1363, 257-269.	1.8	19
43	Enhanced detection for determination of enantiomeric purity of novel agomelatine analogs by EKC using single and dual cyclodextrin systems. <i>Electrophoresis</i> , 2014, 35, 2785-2792.	1.3	12
44	4-Oxo-1,4-dihydropyridines as Selective CB ₂ Cannabinoid Receptor Ligands Part 2: Discovery of New Agonists Endowed with Protective Effect Against Experimental Colitis. <i>Journal of Medicinal Chemistry</i> , 2012, 55, 8948-8952.	2.9	21
45	Analytical and Preparative Chiral Separation of $\hat{1}^2$ -Carboline Derivatives, LDL Oxidation Inhibitors, Using HPLC and CE Methodologies: Determination of Enantiomeric Purity. <i>Chromatographia</i> , 2012, 75, 337-345.	0.7	12
46	Synthesis of 2,3 and 4,5-Dihydro-hydroxy-isoxazoles and Isoxazoles Under Different pH Conditions. <i>Letters in Organic Chemistry</i> , 2010, 7, 32-38.	0.2	10
47	Determination of pKa values of benzimidazole derivatives from mobility obtained by capillary electrophoresis. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2010, 53, 1267-1271.	1.4	13
48	Dual CD system in capillary electrophoresis for direct separation of the four stereoisomers of agonist and antagonist melatonergic ligands. <i>Electrophoresis</i> , 2010, 31, 1529-1532.	1.3	19
49	Chiral Capillary Electrophoresis with Highly Sulfated Cyclodextrins; Resolution of Benzoxazolinone Aminoalcohols, and Aminoketon Precursors, Potential Adrenergic Ligands. <i>Analytical Letters</i> , 2010, 43, 2356-2371.	1.0	4
50	Enantioseparation of benzoxazolinone aminoalcohols and their aminoketone precursors, potential adrenergic ligands, by analytical and preparative liquid chromatography on amylose chiral stationary phases and characterization of the enantiomers. <i>Chirality</i> , 2009, 21, 769-776.	1.3	8
51	Enantioseparation of chiral benzimidazole derivatives by electrokinetic chromatography using sulfated cyclodextrins. <i>Journal of Separation Science</i> , 2009, 32, 1907-1915.	1.3	11
52	Enantiomeric Resolution of Benzoxazolinonic Aminoalcohols, Potential Adrenergic Ligands, by LC and CE. <i>Chromatographia</i> , 2009, 69, 465-472.	0.7	4
53	Chiral Separation of Four Piperidinic Benzoxazolinone Compounds by CE and LC. <i>Chromatographia</i> , 2009, 69, 903-909.	0.7	3
54	LC Using Two Different Cellulose Chiral Stationary Phases for Direct Enantioseparation of Benzoxazolinone Aminoalcohols and Aminoketones. <i>Chromatographia</i> , 2008, 68, 1053-1057.	0.7	1

#	ARTICLE	IF	CITATIONS
55	Chiral resolution of melatonergic ligands by EKC using highly sulfated CDs. <i>Electrophoresis</i> , 2007, 28, 3915-3921.	1.3	5
56	Synthesis of diastereoisomeric pairs of novel analogues of d4T having an isochroman glycon moiety; their enzymatic kinetic resolution, their enantiopure synthesis, molecular modeling and NMR structural study. <i>Tetrahedron</i> , 2005, 61, 10583-10595.	1.0	4
57	Enantiomeric excess determination, purification and biological evaluation of (3S) and (3R) β , β' -butenolide analogues of isobenzofuranone. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2005, 15, 501-504.	1.0	9
58	Preparative HPLC separation of methoxytetralins, ligands for melatonin receptors, containing two chiral centers with polysaccharide chiral stationary phases. Determination of enantiomeric purity. <i>Journal of Proteomics</i> , 2005, 64, 46-58.	2.4	7
59	Enantiomeric analysis of baclofen analogs by capillary zone electrophoresis, using highly sulfated cyclodextrins: Inclusion ionization constant pKa determination. <i>Electrophoresis</i> , 2005, 26, 2974-2983.	1.3	20
60	Enantioseparation of chiral N-imidazole derivatives by electrokinetic chromatography using highly sulfated cyclodextrins: Mechanism of enantioselective recognition. <i>Electrophoresis</i> , 2005, 26, 3824-3832.	1.3	15
61	HPLC Separation and Determination of Enantiomeric Purity of Novel Nucleoside Analogs, on Cyclodextrin Chiral Stationary Phases, Using Reversed and Polar Organic Modes. <i>Analytical Letters</i> , 2004, 37, 385-398.	1.0	6
62	Enantioseparation of new nucleoside analogs, related to d4T and acyclovir, by chiral capillary electrophoresis using highly sulfated β -cyclodextrins. <i>Electrophoresis</i> , 2004, 25, 444-453.	1.3	15
63	Design and synthesis of 3-phenyl tetrahydronaphthalenic derivatives as new selective MT2 melatonergic ligands. <i>Bioorganic and Medicinal Chemistry</i> , 2003, 11, 753-759.	1.4	36
64	Determination of the enantiomeric purity of nucleoside analogs related to d4T and acyclovir, new potential antiviral agents, using liquid chromatography on cellulose chiral stationary phases. <i>Journal of Chromatography A</i> , 2002, 972, 211-219.	1.8	10
65	Diastereomeric resolution of nucleoside analogues, new potential antiviral agents, using high-performance liquid chromatography on polysaccharide-type chiral stationary phases. <i>Journal of Chromatography A</i> , 2002, 943, 91-100.	1.8	22
66	Melatonin receptor agents: Synthesis, resolution by HPLC on polysaccharides chiral stationary phases, absolute configuration, and pharmacology of the enantiomers of (R)-N-[2-(7-fluoro-1,2,3,4-tetrahydroisochroman-5-yl)ethyl]acetamide. <i>Journal of Medicinal Chemistry</i> , 2002, 45, 1000-1010.	1.8	10