

Mariano Monzo

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8292840/mariano-monzo-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

100
papers

4,364
citations

32
h-index

65
g-index

107
ext. papers

4,746
ext. citations

4.3
avg, IF

4.83
L-index

#	Paper	IF	Citations
100	Human embryonic mesenchymal lung-conditioned medium promotes differentiation to myofibroblast and loss of stemness phenotype in lung adenocarcinoma cell lines.. <i>Journal of Experimental and Clinical Cancer Research</i> , 2022 , 41, 37	12.8	0
99	Role of the epithelial-mesenchymal transition-related circular RNA, circ-10720, in non-small-cell lung cancer. <i>Translational Lung Cancer Research</i> , 2021 , 10, 1804-1818	4.4	1
98	Lincp21-RNA as Predictive Response Marker for Preoperative Chemoradiotherapy in Rectal Cancer. <i>Journal of Personalized Medicine</i> , 2021 , 11,	3.6	1
97	LincRNA-p21 as predictive response marker for preoperative chemoradiotherapy in rectal cancer.. <i>Journal of Clinical Oncology</i> , 2021 , 39, e15534-e15534	2.2	
96	The tibioacaneal bundle of the deltoid ligament - Prevalence and variations. <i>Foot and Ankle Surgery</i> , 2021 , 27, 138-142	3.1	5
95	LncRNA Quantification from Extracellular Vesicles Isolated from Blood Plasma or Conditioned Media. <i>Methods in Molecular Biology</i> , 2021 , 2348, 285-304	1.4	1
94	Monitoring HOTTIP levels on extracellular vesicles for predicting recurrence in surgical non-small cell lung cancer patients. <i>Translational Oncology</i> , 2021 , 14, 101144	4.9	0
93	Extracellular Vesicle lincRNA-p21 Expression in Tumor-Draining Pulmonary Vein Defines Prognosis in NSCLC and Modulates Endothelial Cell Behavior. <i>Cancers</i> , 2020 , 12,	6.6	22
92	KRAS mutations by digital PCR in circulating tumor cells isolated from the mesenteric vein are associated with residual disease and overall survival in resected colorectal cancer patients. <i>International Journal of Colorectal Disease</i> , 2020 , 35, 805-813	3	1
91	Long Non-Coding RNA NANCI/NKX2-1 Duplex Impacts Prognosis in Stage I Non-Small-Cell Lung Cancer 2020 , 56, 630-630		
90	Long Non-Coding RNA NANCI/NKX2-1 Duplex Impacts Prognosis in Stage I Non-Small-Cell Lung Cancer. <i>Archivos De Bronconeumologia</i> , 2020 , 56, 630-636	0.7	3
89	LincRNA-p21 Levels Relates to Survival and Post-Operative Radiotherapy Benefit in Rectal Cancer Patients. <i>Life</i> , 2020 , 10,	3	2
88	Clinical significance of long non-coding RNA HOTTIP in early-stage non-small-cell lung cancer. <i>BMC Pulmonary Medicine</i> , 2019 , 19, 55	3.5	17
87	Exosome Analysis in Tumor-Draining Pulmonary Vein Identifies NSCLC Patients with Higher Risk of Relapse after Curative Surgery. <i>Cancers</i> , 2019 , 11,	6.6	18
86	Proteomic Analysis of Liquid Biopsy from Tumor-Draining Vein Indicates that High Expression of Exosomal ECM1 Is Associated with Relapse in Stage I-III Colon Cancer. <i>Translational Oncology</i> , 2018 , 11, 715-721	4.9	15
85	miR-328 mediates a metabolic shift in colon cancer cells by targeting SLC2A1/GLUT1. <i>Clinical and Translational Oncology</i> , 2018 , 20, 1161-1167	3.6	29
84	Prognostic Impact of miR-200 Family Members in Plasma and Exosomes from Tumor-Draining versus Peripheral Veins of Colon Cancer Patients. <i>Oncology</i> , 2018 , 95, 309-318	3.6	26

83	Rotational Instability after Anterior Talofibular and Calcaneofibular Ligament Section: The Experimental Basis for the Ankle Pivot Test. <i>Journal of Foot and Ankle Surgery</i> , 2018 , 57, 1087-1091	1.6	15
82	miR-21, miR-99b and miR-375 combination as predictive response signature for preoperative chemoradiotherapy in rectal cancer. <i>PLoS ONE</i> , 2018 , 13, e0206542	3.7	23
81	NKX2-1 expression as a prognostic marker in early-stage non-small-cell lung cancer. <i>BMC Pulmonary Medicine</i> , 2017 , 17, 197	3.5	8
80	Exosomal microRNAs isolated from plasma of mesenteric veins linked to liver metastases in resected patients with colon cancer. <i>Oncotarget</i> , 2017 , 8, 30859-30869	3.3	27
79	Epigenetic regulation mechanisms of microRNA expression. <i>Biomolecular Concepts</i> , 2017 , 8, 203-212	3.7	78
78	Non-Coding RNAs in Hodgkin Lymphoma. <i>International Journal of Molecular Sciences</i> , 2017 , 18,	6.3	10
77	LincRNA-p21 Impacts Prognosis in Resected Non-Small Cell Lung Cancer Patients through Angiogenesis Regulation. <i>Journal of Thoracic Oncology</i> , 2016 , 11, 2173-2182	8.9	65
76	miR-203 and miR-221 regulate SOCS1 and SOCS3 in essential thrombocythemia. <i>Blood Cancer Journal</i> , 2016 , 6, e406	7	16
75	PiwiRNA-651 as marker of treatment response and survival in classical Hodgkin lymphoma. <i>Oncotarget</i> , 2016 , 7, 46002-46013	3.3	32
74	YKT6 expression, exosome release, and survival in non-small cell lung cancer. <i>Oncotarget</i> , 2016 , 7, 51515-51524	3.5	46
73	Talonavicular joint arthroscopic portals: A cadaveric study of feasibility and safety. <i>Foot and Ankle Surgery</i> , 2016 , 22, 205-209	3.1	3
72	Identifying High-Risk Stage II Colon Cancer Patients: A Three-MicroRNA-Based Score as a Prognostic Biomarker. <i>Clinical Colorectal Cancer</i> , 2016 , 15, e175-e182	3.8	31
71	Differential MIR-21 expression in plasma from mesenteric versus peripheral veins: an observational study of disease-free survival in surgically resected colon cancer patients. <i>Medicine (United States)</i> , 2015 , 94, e145	1.8	10
70	The expression level of BAALC-associated microRNA miR-3151 is an independent prognostic factor in younger patients with cytogenetic intermediate-risk acute myeloid leukemia. <i>Blood Cancer Journal</i> , 2015 , 5, e352	7	15
69	The significance of PIWI family expression in human lung embryogenesis and non-small cell lung cancer. <i>Oncotarget</i> , 2015 , 6, 31544-56	3.3	37
68	Epigenetic regulation of microRNA expression in Hodgkin lymphoma. <i>Leukemia and Lymphoma</i> , 2015 , 56, 2683-9	1.9	11
67	piRNA-651 as a prognostic marker in surgically resected non-small-cell lung cancer.. <i>Journal of Clinical Oncology</i> , 2015 , 33, 11043-11043	2.2	1
66	A serum microRNA signature associated with complete remission and progression after autologous stem-cell transplantation in patients with multiple myeloma. <i>Oncotarget</i> , 2015 , 6, 1874-83	3.3	36

65	The lincRNA HOTAIRM1, located in the HOXA genomic region, is expressed in acute myeloid leukemia, impacts prognosis in patients in the intermediate-risk cytogenetic category, and is associated with a distinctive microRNA signature. <i>Oncotarget</i> , 2015 , 6, 31613-27	3.3	69
64	miR-141 and miR-200c as markers of overall survival in early stage non-small cell lung cancer adenocarcinoma. <i>PLoS ONE</i> , 2014 , 9, e101899	3.7	103
63	MicroRNA expression at diagnosis adds relevant prognostic information to molecular categorization in patients with intermediate-risk cytogenetic acute myeloid leukemia. <i>Leukemia</i> , 2014 , 28, 804-12	10.7	51
62	Role of miR-200 family members in survival of colorectal cancer patients treated with fluoropyrimidines. <i>Journal of Surgical Oncology</i> , 2014 , 109, 676-83	2.8	55
61	miR-200 family in CRC primary tumors and metastases. <i>Journal of Surgical Oncology</i> , 2014 , 110, 486	2.8	2
60	High levels of global DNA methylation are an independent adverse prognostic factor in a series of 90 patients with de novo myelodysplastic syndrome. <i>Leukemia Research</i> , 2014 , 38, 874-81	2.7	15
59	Piwirna-651 Expression Influences Treatment Response and Impacts Survival in Classical Hodgkin Lymphoma Patients through Regulation of ABCC5. <i>Blood</i> , 2014 , 124, 134-134	2.2	2
58	The LincRNA HOTAIRM1, Located in the HOXA genomic Region, impacts Prognosis in Acute Myeloid Leukemia and Is Associated with a Distinctive microRNA Signature. <i>Blood</i> , 2014 , 124, 1003-1003 ^{2,2}		
57	Acute myeloid leukemia with translocation (8;16)(p11;p13) and MYST3-CREBBP rearrangement harbors a distinctive microRNA signature targeting RET proto-oncogene. <i>Leukemia</i> , 2013 , 27, 595-603	10.7	39
56	Enalapril and carvedilol for preventing chemotherapy-induced left ventricular systolic dysfunction in patients with malignant hemopathies: the OVERCOME trial (prevention of left Ventricular dysfunction with Enalapril and carvedilol in patients submitted to intensive Chemotherapy for the treatment of Malignant hEmopathies). <i>Journal of the American College of Cardiology</i> , 2013 , 61, 2355-62	15.1	403
55	Impact of global and gene-specific DNA methylation pattern in relapsed multiple myeloma patients treated with bortezomib. <i>Leukemia Research</i> , 2013 , 37, 641-6	2.7	10
54	Low miR-145 and high miR-367 are associated with unfavourable prognosis in resected nonsmall cell lung cancer. <i>European Respiratory Journal</i> , 2013 , 41, 1172-8	13.6	61
53	Pint lincRNA connects the p53 pathway with epigenetic silencing by the Polycomb repressive complex 2. <i>Genome Biology</i> , 2013 , 14, R104	18.3	182
52	MicroRNA expression in chronic lymphocytic leukemia developing autoimmune hemolytic anemia. <i>Leukemia and Lymphoma</i> , 2013 , 54, 2016-22	1.9	24
51	MiR-SNPs as markers of toxicity and clinical outcome in Hodgkin lymphoma patients. <i>PLoS ONE</i> , 2013 , 8, e64716	3.7	21
50	Prognostic Significance Of a 4-Microrna Signature Targeting JAK2 In Classical Hodgkin Lymphoma. <i>Blood</i> , 2013 , 122, 629-629	2.2	1
49	Association of miR-141 and miR-200c with time to recurrence (TTR) and overall survival (OS) in resected non-small-cell lung cancer (NSCLC) adenocarcinoma (ADC) patients (p).. <i>Journal of Clinical Oncology</i> , 2013 , 31, 7547-7547	2.2	
48	Impact of MiRSNPs on survival and progression in patients with multiple myeloma undergoing autologous stem cell transplantation. <i>Clinical Cancer Research</i> , 2012 , 18, 3697-704	12.9	49

47	Single nucleotide polymorphisms analysis of BRAC and ERCC1 as predictor of recurrence after chemoradiation for cervical cancer patients.. <i>Journal of Clinical Oncology</i> , 2012 , 30, e13540-e13540	2.2	
46	Prevention of chemotherapy-induced left ventricular dysfunction with enalapril and carvedilol: rationale and design of the OVERCOME trial. <i>Journal of Cardiac Failure</i> , 2011 , 17, 643-8	3.3	22
45	Single nucleotide polymorphisms in tobacco metabolism and DNA repair genes and prognosis in resected non-small-cell lung cancer. <i>Journal of Surgical Research</i> , 2011 , 167, e5-12	2.5	6
44	Lestaurtinib inhibition of the Jak/STAT signaling pathway in hodgkin lymphoma inhibits proliferation and induces apoptosis. <i>PLoS ONE</i> , 2011 , 6, e18856	3.7	53
43	Impact on response and survival of DNA repair single nucleotide polymorphisms in relapsed or refractory multiple myeloma patients treated with thalidomide. <i>Leukemia Research</i> , 2011 , 35, 1178-83	2.7	19
42	Expression of Myosin Heavy Chain Isoforms in the Supraspinatus Muscle of Different Primate Species: Implications for the Study of the Adaptation of Primate Shoulder Muscles to Different Locomotor Modes. <i>International Journal of Primatology</i> , 2011 , 32, 931-944	2	10
41	Prognostic implications of miR-16 expression levels in resected non-small-cell lung cancer. <i>Journal of Surgical Oncology</i> , 2011 , 103, 411-5	2.8	48
40	Ultrastructural and immunohistochemical analysis of intestinal myofibroblasts during the early organogenesis of the human small intestine. <i>Anatomical Record</i> , 2011 , 294, 462-71	2.1	13
39	A dual role for KRT81: a miR-SNP associated with recurrence in non-small-cell lung cancer and a novel marker of squamous cell lung carcinoma. <i>PLoS ONE</i> , 2011 , 6, e22509	3.7	45
38	MicroRNAs in human embryonic and cancer stem cells. <i>Yonsei Medical Journal</i> , 2010 , 51, 622-32	3	33
37	MicroRNAs expressed during lung cancer development are expressed in human pseudoglandular lung embryogenesis. <i>Oncology</i> , 2009 , 76, 162-9	3.6	47
36	Regulation of JAK2 by miR-135a: prognostic impact in classic Hodgkin lymphoma. <i>Blood</i> , 2009 , 114, 2945-51	5.1	141
35	miR-34a as a prognostic marker of relapse in surgically resected non-small-cell lung cancer. <i>Carcinogenesis</i> , 2009 , 30, 1903-9	4.6	282
34	Gene single nucleotide polymorphism accumulation improves survival in advanced head and neck cancer patients treated with weekly paclitaxel. <i>Laryngoscope</i> , 2009 , 119, 1484-90	3.6	15
33	Hematopoiesis-related microRNA expression in myelodysplastic syndromes. <i>Leukemia and Lymphoma</i> , 2009 , 50, 1854-9	1.9	74
32	Overlapping expression of microRNAs in human embryonic colon and colorectal cancer. <i>Cell Research</i> , 2008 , 18, 823-33	24.7	159
31	Common variants in NLRP2 and NLRP3 genes are strong prognostic factors for the outcome of HLA-identical sibling allogeneic stem cell transplantation. <i>Blood</i> , 2008 , 112, 4337-42	2.2	30
30	Epigenetic events in normal colonic mucosa surrounding colorectal cancer lesions. <i>European Journal of Cancer</i> , 2008 , 44, 2689-95	7.5	25

29	A single-nucleotide polymorphism in the aromatase gene is associated with the efficacy of the aromatase inhibitor letrozole in advanced breast carcinoma. <i>Clinical Cancer Research</i> , 2008 , 14, 811-6	12.9	104
28	MicroRNA expression profiling in classic Hodgkin lymphoma. <i>Blood</i> , 2008 , 111, 2825-32	2.2	148
27	Pharmacogenomics: a tool for improving cancer chemotherapy. <i>Clinical and Translational Oncology</i> , 2008 , 10, 628-37	3.6	6
26	Regulation of Jak2 Kinase Gene by the Mir-135 Family in Classical Hodgkin Lymphoma. <i>Blood</i> , 2008 , 112, 518-518	2.2	
25	Single nucleotide polymorphisms in nucleotide excision repair genes XPA, XPD, XPG and ERCC1 in advanced colorectal cancer patients treated with first-line oxaliplatin/fluoropyrimidine. <i>Oncology</i> , 2007 , 72, 364-70	3.6	51
24	Hematopoietic miRNAs Expression in Myelodysplastic Syndromes.. <i>Blood</i> , 2007 , 110, 2422-2422	2.2	
23	Genetic Polymorphisms in the Inflammasomes Are Associated with Relapse and Survival in HLA-Identical Sibling Donor Allogeneic Stem Cell Transplantation.. <i>Blood</i> , 2007 , 110, 1075-1075	2.2	
22	Mir-135a Expression Is Associated with Relapse in Hodgkin Lymphoma.. <i>Blood</i> , 2007 , 110, 2270-2270	2.2	
21	MicroRNA Analysis by In Situ Hybridization in Hodgkin Lymphoma.. <i>Blood</i> , 2007 , 110, 2271-2271	2.2	
20	Genomic polymorphisms provide prognostic information in intermediate-risk acute myeloblastic leukemia. <i>Blood</i> , 2006 , 107, 4871-9	2.2	58
19	Identification by Real-time PCR of 13 mature microRNAs differentially expressed in colorectal cancer and non-tumoral tissues. <i>Molecular Cancer</i> , 2006 , 5, 29	42.1	668
18	Time-related improvement of survival in resectable gastric cancer: the role of Japanese-style gastrectomy with D2 lymphadenectomy and adjuvant chemotherapy. <i>World Journal of Surgical Oncology</i> , 2006 , 4, 53	3.4	6
17	Sonic hedgehog mRNA expression by real-time quantitative PCR in normal and tumor tissues from colorectal cancer patients. <i>Cancer Letters</i> , 2006 , 233, 117-23	9.9	33
16	Adra2A gene expression levels during the recovery of a2D-adrenoceptors after chronic lithium treatment in rat cerebral cortex. <i>FASEB Journal</i> , 2006 , 20, LB113	0.9	
15	Effect of Germline Polymorphisms on Clinical Outcome in Hodgkin Lymphoma (HL).. <i>Blood</i> , 2006 , 108, 2267-2267	2.2	
14	Analysis of microRNA Patterns in Hodgkin Lymphoma (HL).. <i>Blood</i> , 2006 , 108, 474-474	2.2	
13	Effect of NOD2/CARD15 variants in T-cell depleted allogeneic stem cell transplantation. <i>Haematologica</i> , 2006 , 91, 1372-6	6.6	47
12	Letrozole efficacy is related to human aromatase CYP19 single nucleotide polymorphisms (SNPs) in metastatic breast cancer patients. <i>Journal of Clinical Oncology</i> , 2004 , 22, 507-507	2.2	2

11	Expression of cyclooxygenase-2 mRNA (COX2-mRNA) in peripheral blood (PB) of head and neck cancer patients (HNC) and healthy controls (HC). <i>Journal of Clinical Oncology</i> , 2004 , 22, 5521-5521	2.2	2
10	Letrozole efficacy is related to human aromatase CYP19 single nucleotide polymorphisms (SNPs) in metastatic breast cancer patients. <i>Journal of Clinical Oncology</i> , 2004 , 22, 507-507	2.2	3
9	Expression of cyclooxygenase-2 mRNA (COX2-mRNA) in peripheral blood (PB) of head and neck cancer patients (HNC) and healthy controls (HC). <i>Journal of Clinical Oncology</i> , 2004 , 22, 5521-5521	2.2	2
8	Translational oncogenomics: toward rational therapeutic decision-making. <i>Current Opinion in Oncology</i> , 2002 , 14, 171-9	4.2	18
7	A novel anti-apoptosis gene: Re-expression of survivin messenger RNA as a prognosis marker in non-small-cell lung cancers. <i>Journal of Clinical Oncology</i> , 1999 , 17, 2100-4	2.2	43 ⁰
6	k-FGF Protoncogene Expression is Associated With Murine Testicular Teratogenesis, But is Not Involved During Mouse Testicular Development. <i>Journal of Urology</i> , 1998 , 159, 2240-2241	2.5	
5	TP53 mutational pattern in Spanish and Polish non-small cell lung cancer patients: null mutations are associated with poor prognosis. <i>Oncogene</i> , 1997 , 15, 2951-8	9.2	43
4	Growth control of embryonic stem cells injected into mouse uterus on fifth day of pregnancy. <i>International Journal of Cancer</i> , 1994 , 56, 387-92	7.5	7
3	Laminin and fibronectin expression during in vivo growth of embryoid bodies derived from teratocarcinoma. <i>Virchows Archiv A, Pathological Anatomy and Histopathology</i> , 1991 , 418, 509-14		
2	Alpha-fetoprotein in tumours derived from cystic and simple embryoid bodies. <i>Archivio Italiano Di Anatomia E Di Embriologia Italian Journal of Anatomy and Embryology</i> , 1991 , 96, 113-20		
1	Cytodifferentiation of mouse visceral endoderm: study with a teratocarcinoma antiserum. <i>Acta Anatomica</i> , 1988 , 132, 66-8		