

# Filipe Pinto

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8296959/publications.pdf>

Version: 2024-02-01

24  
papers

495  
citations

759055

12  
h-index

752573

20  
g-index

24  
all docs

24  
docs citations

24  
times ranked

672  
citing authors

#	ARTICLE	IF	CITATIONS
1	The Extracellular Small Leucine-Rich Proteoglycan Biglycan Is a Key Player in Gastric Cancer Aggressiveness. <i>Cancers</i> , 2021, 13, 1330.	1.7	26
2	Clinical impact of brachyury expression in Ewing sarcoma patients. <i>Advances in Medical Sciences</i> , 2021, 66, 321-325.	0.9	0
3	Loss of SPINT2 expression frequently occurs in glioma, leading to increased growth and invasion via MMP2. <i>Cellular Oncology (Dordrecht)</i> , 2020, 43, 107-121.	2.1	8
4	Brachyury Is Associated with Glioma Differentiation and Response to Temozolomide. <i>Neurotherapeutics</i> , 2020, 17, 2015-2027.	2.1	7
5	The Tâ€ transcription factor brachyury behaves as a tumor suppressor in gliomas. <i>Journal of Pathology</i> , 2020, 251, 87-99.	2.1	10
6	Impact of Truncated O-glycans in Gastric-Cancer-Associated CD44v9 Detection. <i>Cells</i> , 2020, 9, 264.	1.8	11
7	High-Throughput Sequencing Identifies 3 Novel Susceptibility Genes for Hereditary Melanoma. <i>Genes</i> , 2020, 11, 403.	1.0	14
8	Hypoxia and Macrophages Act in Concert Towards a Beneficial Outcome in Colon Cancer. <i>Cancers</i> , 2020, 12, 818.	1.7	9
9	Carcinoembryonic antigen carrying SLe <sup>X</sup> as a new biomarker of more aggressive gastric carcinomas. <i>Theranostics</i> , 2019, 9, 7431-7446.	4.6	35
10	O-glycans truncation modulates gastric cancer cell signaling and transcription leading to a more aggressive phenotype. <i>EBioMedicine</i> , 2019, 40, 349-362.	2.7	63
11	Genetic variants of vascular endothelial growth factor predict risk and survival of gliomas. <i>Tumor Biology</i> , 2018, 40, 101042831876627.	0.8	9
12	Brachyury oncogene is a prognostic factor in high-risk testicular germ cell tumors. <i>Andrology</i> , 2018, 6, 597-604.	1.9	11
13	Hypoxia and serum deprivation induces glycan alterations in triple negative breast cancer cells. <i>Biological Chemistry</i> , 2018, 399, 661-672.	1.2	11
14	Brachyury, a driver of epithelial mesenchymal transition, as an independent prognostic factor in high-grade testicular germ cell tumors.. <i>Journal of Clinical Oncology</i> , 2017, 35, e16039-e16039.	0.8	0
15	The embryonic Brachyury transcription factor is a novel biomarker of GIST aggressiveness and poor survival. <i>Gastric Cancer</i> , 2016, 19, 651-659.	2.7	18
16	SPINT2 Deregulation in Prostate Carcinoma. <i>Journal of Histochemistry and Cytochemistry</i> , 2016, 64, 32-41.	1.3	13
17	Brachyury identifies a class of enteroendocrine cells in normal human intestinal crypts and colorectal cancer. <i>Oncotarget</i> , 2016, 7, 11478-11486.	0.8	47
18	Brachyury as a potential modulator of androgen receptor activity and a key player in therapy resistance in prostate cancer. <i>Oncotarget</i> , 2016, 7, 28891-28902.	0.8	19

#	ARTICLE	IF	CITATIONS
19	Drivers of neuroendocrine prostate cancer. <i>Translational Cancer Research</i> , 2016, 5, S551-S553.	0.4	0
20	Silencing of WNK2 is associated with upregulation of MMP2 and JNK in gliomas. <i>Oncotarget</i> , 2015, 6, 1422-1434.	0.8	21
21	T-box Transcription Factor Brachyury Is Associated with Prostate Cancer Progression and Aggressiveness. <i>Clinical Cancer Research</i> , 2014, 20, 4949-4961.	3.2	67
22	Loss of WNK2 expression by promoter gene methylation occurs in adult gliomas and triggers Rac1-mediated tumour cell invasiveness. <i>Human Molecular Genetics</i> , 2013, 22, 84-95.	1.4	44
23	RKIP Inhibition in Cervical Cancer Is Associated with Higher Tumor Aggressive Behavior and Resistance to Cisplatin Therapy. <i>PLoS ONE</i> , 2013, 8, e59104.	1.1	52
24	295 Protein Kinase WNK2 Was Correlated With Poor Outcome and Malignant Behavior in Glioma Cell Lines. <i>European Journal of Cancer</i> , 2012, 48, S72.	1.3	0