

# Zi-Hua Zhao

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8307657/publications.pdf>

Version: 2024-02-01

42  
papers

702  
citations

623734  
14  
h-index

610901  
24  
g-index

44  
all docs

44  
docs citations

44  
times ranked

720  
citing authors

#	ARTICLE	IF	CITATIONS
1	Effects of agricultural intensification on ability of natural enemies to control aphids. Scientific Reports, 2015, 5, 8024.	3.3	58
2	Laboratory and field efficacy of entomopathogenic fungi for the management of the sweetpotato weevil, <i>Cylas formicarius</i> (Coleoptera: Brentidae). Journal of Invertebrate Pathology, 2014, 122, 10-15.	3.2	53
3	Climate change impacts on the global potential geographical distribution of the agricultural invasive pest, <i>Bactrocera dorsalis</i> (Hendel) (Diptera: Tephritidae). Climatic Change, 2019, 155, 145-156.	3.6	52
4	Global distribution and invasion pattern of oriental fruit fly, <i>Bactrocera dorsalis</i> (Diptera: Tephritidae). Journal of Economic Entomology, 2019, 112, 10-15.	1.8	44
5	Effects of inter-annual landscape change on interactions between cereal aphids and their natural enemies. Basic and Applied Ecology, 2013, 14, 472-479.	2.7	40
6	Solving the pitfalls of pitfall trapping: a two-circle method for density estimation of ground-dwelling arthropods. Methods in Ecology and Evolution, 2013, 4, 865-871.	5.2	33
7	Responses of Cereal Aphids and Their Parasitic Wasps to Landscape Complexity. Journal of Economic Entomology, 2014, 107, 630-637.	1.8	31
8	Approaches and mechanisms for ecologically based pest management across multiple scales. Agriculture, Ecosystems and Environment, 2016, 230, 199-209.	5.3	30
9	Effects of position within wheat field and adjacent habitats on the density and diversity of cereal aphids and their natural enemies. BioControl, 2013, 58, 765-776.	2.0	28
10	<i>Bactrocera dorsalis</i> (Hendel) (Diptera: Tephritidae) is not invasive through Asia: It's been there all along. Journal of Applied Entomology, 2019, 143, 797-801.	1.8	28
11	A transcriptional and functional analysis of heat hardening in two invasive fruit fly species, <i>Bactrocera dorsalis</i> and <i>Bactrocera correcta</i> . Evolutionary Applications, 2019, 12, 1147-1163.	3.1	26
12	Population dynamics and associated factors of cereal aphids and armyworms under global change. Scientific Reports, 2016, 5, 18801.	3.3	23
13	From the inverse density-area relationship to the minimum patch size of a host-parasitoid system. Ecological Research, 2012, 27, 303-309.	1.5	19
14	Effects of crop species richness on pest-natural enemy systems based on an experimental model system using a microlandscape. Science China Life Sciences, 2013, 56, 758-766.	4.9	18
15	Landscape changes have greater effects than climate changes on six insect pests in China. Science China Life Sciences, 2016, 59, 627-633.	4.9	14
16	Comparative Transcriptome Analyses Uncover Key Candidate Genes Mediating Flight Capacity in <i>Bactrocera dorsalis</i> (Hendel) and <i>Bactrocera correcta</i> (Bezzi) (Diptera: Tephritidae). International Journal of Molecular Sciences, 2018, 19, 396.	4.1	14
17	The Combined Effect of Elevated O <sub>3</sub> Levels and TYLCV Infection Increases the Fitness of <i>Bemisia tabaci</i> Mediterranean on Tomato Plants. Environmental Entomology, 2019, 48, 1425-1433.	1.4	14
18	Crop Diversity and Land Simplification Effects on Pest Damage in Northern China. Annals of the Entomological Society of America, 2017, 110, 91-96.	2.5	13

#	ARTICLE	IF	CITATIONS
19	A potential genetic control by suppression of the wing developmental gene wingless in a global invasive pest <i>Bactrocera dorsalis</i> . <i>Journal of Pest Science</i> , 2021, 94, 517-529.	3.7	13
20	Shifts in natural enemy assemblages resulting from landscape simplification account for biocontrol loss in wheat fields. <i>Ecological Research</i> , 2015, 30, 493-498.	1.5	12
21	Habitat heterogeneity stabilizes the spatial and temporal interactions between cereal aphids and parasitic wasps. <i>Basic and Applied Ecology</i> , 2015, 16, 510-518.	2.7	12
22	UV radiation increases mortality and decreases the antioxidant activity in a tephritid fly. <i>Food and Energy Security</i> , 2021, 10, e297.	4.3	12
23	Cropland expansion facilitated the outbreak of cereal aphids during 1951–2010 in China. <i>Science Bulletin</i> , 2015, 60, 1036-1037.	9.0	11
24	The failure of success: cyclic recurrences of a globally invasive pest. <i>Ecological Applications</i> , 2019, 29, e01991.	3.8	10
25	Life table invasion models: spatial progression and species-specific partitioning. <i>Ecology</i> , 2019, 100, e02682.	3.2	10
26	Plant cover associated with aboveground net primary productivity (ANPP) mediates insect community composition in steppes of Northwest China. <i>Journal of Asia-Pacific Entomology</i> , 2018, 21, 361-366.	0.9	8
27	Semi-natural habitats mediate influence of inter-annual landscape variation on cereal aphid-parasitic wasp system in an agricultural landscape. <i>Biological Control</i> , 2019, 128, 17-23.	3.0	8
28	Asymmetric response of different functional insect groups to low grazing pressure in Eurasian steppe in Ningxia. <i>Ecology and Evolution</i> , 2018, 8, 11609-11618.	1.9	7
29	Optimization of Nitrogen Fertilizer Application Enhances Biocontrol Function and Net Income. <i>Journal of Economic Entomology</i> , 2020, 113, 2035-2038.	1.8	7
30	The synergy between climate change and transportation activities drives the propagation of an invasive fruit fly. <i>Journal of Pest Science</i> , 2020, 93, 615-625.	3.7	6
31	Increased nitrogen fertilization inhibits the biocontrol activity promoted by the intercropping partner plant. <i>Insect Science</i> , 2020, 28, 1179-1190.	3.0	6
32	Mild Drought Facilitates the Increase in Wheat Aphid Abundance by Changing Host Metabolism. <i>Annals of the Entomological Society of America</i> , 2021, 114, 79-83.	2.5	6
33	Including climate change to predict the global suitable area of an invasive pest: <i>Bactrocera correcta</i> (Diptera: Tephritidae). <i>Global Ecology and Conservation</i> , 2022, 34, e02021.	2.1	6
34	Tracing dietary origins of aphids and the predatory beetle <i>Propylea japonica</i> in agricultural systems using stable isotope analyses. <i>Entomologia Experimentalis Et Applicata</i> , 2015, 155, 87-94.	1.4	5
35	Effects of seed mixture sowing with resistant and susceptible rice on population dynamics of target planthoppers and non-target stemborers and leafhoppers. <i>Pest Management Science</i> , 2018, 74, 1664-1676.	3.4	5
36	Cultivar Mixture Enhances Crop Yield by Decreasing Aphids. <i>Agronomy</i> , 2022, 12, 335.	3.0	5

#	ARTICLE	IF	CITATIONS
37	Plant Species Richness Controls Arthropod Food Web: Evidence From an Experimental Model System. <i>Annals of the Entomological Society of America</i> , 2019, 112, 27-32.	2.5	4
38	The potential geographic distribution of <i>Bactrocera correcta</i> (Diptera: Tephritidae) in China based on eclosion rate model. <i>Applied Entomology and Zoology</i> , 2015, 50, 371-381.	1.2	3
39	Differential temperature responses between <i>Plutella xylostella</i> and its specialist endoparasitoid <i>Diadegma semiclausum</i> —Implications for biological control. <i>Insect Science</i> , 2022, 29, 855-864.	3.0	3
40	The asymmetric responses of carabid beetles to steppe fragmentation in Northwest China. <i>Global Ecology and Conservation</i> , 2020, 23, e01058.	2.1	2
41	Potential pest invasion risk posed by international sweet cherry trade. <i>Food and Energy Security</i> , 2021, 10, e257.	4.3	2
42	Landscape pattern affects species composition and abundance of ground-dwelling predator. <i>Journal of Asia-Pacific Entomology</i> , 2015, 18, 331-334.	0.9	1