

# Anthony D Heymann, Mbbs Mha

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/8307997/anthony-d-heyman-mbbs-mha-publications-by-year.pdf>  
**Version:** 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.  
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

62 papers	555 citations	13 h-index	22 g-index
68 ext. papers	720 ext. citations	4.2 avg, IF	3.53 L-index

#	Paper	IF	Citations
62	Real-world safety data for the Pfizer BNT162b2 SARS-CoV-2 vaccine, historical cohort study - Author's reply.. <i>Clinical Microbiology and Infection</i> , <b>2021</b> ,	9.5	2
61	Glycemic Control and Risk of Cellulitis. <i>Diabetes Care</i> , <b>2021</b> , 44, 367-372	14.6	3
60	BNT162b2 Vaccine Effectiveness in Preventing Asymptomatic Infection With SARS-CoV-2 Virus: A Nationwide Historical Cohort Study. <i>Open Forum Infectious Diseases</i> , <b>2021</b> , 8, ofab262	1	14
59	Trajectories of depression symptoms over time differ by APOE4 genotype in older adults with type 2 diabetes. <i>International Journal of Geriatric Psychiatry</i> , <b>2021</b> , 36, 1567-1575	3.9	0
58	Preferences and barriers to the utilization of primary health care by sick physicians: a nationwide survey. <i>Family Practice</i> , <b>2021</b> , 38, 109-114	1.9	0
57	Increase in Number of Depression Symptoms Over Time is Related to Worse Cognitive Outcomes in Older Adults With Type 2 Diabetes. <i>American Journal of Geriatric Psychiatry</i> , <b>2021</b> , 29, 1-11	6.5	1
56	Association of the CD2AP locus with cognitive functioning among middle-aged individuals with a family history of Alzheimer's disease. <i>Neurobiology of Aging</i> , <b>2021</b> , 101, 50-56	5.6	1
55	Specific Dimensions of Depression Have Different Associations With Cognitive Decline in Older Adults With Type 2 Diabetes. <i>Diabetes Care</i> , <b>2021</b> , 44, 655-662	14.6	1
54	Real-world safety data for the Pfizer BNT162b2 SARS-CoV-2 vaccine: historical cohort study. <i>Clinical Microbiology and Infection</i> , <b>2021</b> ,	9.5	11
53	The COMT rs4680 polymorphism is associated with rate of cognitive decline in older adults with type 2 diabetes. <i>Alzheimer's and Dementia</i> , <b>2020</b> , 16, e045227	1.2	
52	Trajectories of depression symptoms over time differ by APOE4 carriership in older adults with type 2 diabetes. <i>Alzheimer's and Dementia</i> , <b>2020</b> , 16, e045304	1.2	
51	Greater apathy is associated with faster rate of cognitive decline in older adults with type 2 diabetes. <i>Alzheimer's and Dementia</i> , <b>2020</b> , 16, e045641	1.2	
50	Deterioration in Motor Function Over Time in Older Adults With Type 2 Diabetes is Associated with Accelerated Cognitive Decline. <i>Endocrine Practice</i> , <b>2020</b> , 26, 1143-1152	3.2	6
49	Distinct trajectories in HbA1c are associated with different all-cause mortality and morbidity in newly diagnosed patients with type 2 diabetes. <i>Primary Care Diabetes</i> , <b>2020</b> , 14, 413-419	2.4	5
48	The Israel Registry for Alzheimer's Prevention (IRAP) Study: Design and Baseline Characteristics. <i>Journal of Alzheimer's Disease</i> , <b>2020</b> , 78, 777-788	4.3	3
47	Higher BMI is associated with smaller regional brain volume in older adults with type 2 diabetes. <i>Diabetologia</i> , <b>2020</b> , 63, 2446-2451	10.3	2
46	Long-term trajectories of BMI predict carotid stiffness and plaque volume in type 2 diabetes older adults: a cohort study. <i>Cardiovascular Diabetology</i> , <b>2020</b> , 19, 138	8.7	2

45	Computerized Cognitive Training for Older Adults at Higher Dementia Risk due to Diabetes: Findings From a Randomized Controlled Trial. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , <b>2020</b> , 75, 747-754	6.4	11
44	Vitamin E Intake Is Associated with Lower Brain Volume in Haptoglobin 1-1 Elderly with Type 2 Diabetes. <i>Journal of Alzheimer's Disease</i> , <b>2020</b> , 74, 649-658	4.3	1
43	The Association of Depressive Symptoms With Brain Volume Is Stronger Among Diabetic Elderly Carriers of the Haptoglobin 1-1 Genotype Compared to Non-carriers. <i>Frontiers in Endocrinology</i> , <b>2019</b> , 10, 68	5.7	1
42	TCF7L2 polymorphisms are associated with amygdalar volume in elderly individuals with Type 2 Diabetes. <i>Scientific Reports</i> , <b>2019</b> , 9, 15818	4.9	3
41	Depression is more strongly associated with cognition in elderly women than men with type 2 diabetes. <i>International Psychogeriatrics</i> , <b>2019</b> , 31, 591-595	3.4	
40	The gene is associated with processing speed performance - evidence among elderly with type 2 diabetes. <i>World Journal of Biological Psychiatry</i> , <b>2019</b> , 20, 577-583	3.8	5
39	Depressive Symptoms Are Associated with Cognitive Function in the Elderly with Type 2 Diabetes. <i>Journal of Alzheimer's Disease</i> , <b>2018</b> , 65, 683-692	4.3	10
38	Comparing assessment of diabetes-related quality of life between patients and their physicians. <i>Health and Quality of Life Outcomes</i> , <b>2018</b> , 16, 214	3	3
37	Adherence to quality of care measurements among 58,182 patients with new onset diabetes and its association with mortality. <i>PLoS ONE</i> , <b>2018</b> , 13, e0208539	3.7	1
36	A paradigm shift: from doctor-patient to payer-patient relationship. <i>British Journal of General Practice</i> , <b>2018</b> , 68, 438-439	1.6	0
35	Hemoglobin A Variability Predicts Symptoms of Depression in Elderly Individuals With Type 2 Diabetes. <i>Diabetes Care</i> , <b>2017</b> , 40, 1187-1193	14.6	18
34	Haptoglobin 1-1 Genotype Modulates the Association of Glycemic Control With Hippocampal Volume in Elderly Individuals With Type 2 Diabetes. <i>Diabetes</i> , <b>2017</b> , 66, 2927-2932	0.9	9
33	Impact of the Holocaust on the Rehabilitation Outcome of Older Patients Sustaining a Hip Fracture. <i>Israel Medical Association Journal</i> , <b>2017</b> , 19, 207-210	0.9	1
32	Potential contribution of the Alzheimer's disease risk locus BIN1 to episodic memory performance in cognitively normal Type 2 diabetes elderly. <i>European Neuropsychopharmacology</i> , <b>2016</b> , 26, 787-95	1.2	9
31	Unsupportive Social Interactions are Associated with Poorer Self-reported Health in Israeli Patients with Diabetes. <i>American Journal of Health Behavior</i> , <b>2016</b> , 40, 645-51	1.9	2
30	Ethnicity/culture modulates the relationships of the haptoglobin (Hp) 1-1 phenotype with cognitive function in older individuals with type 2 diabetes. <i>International Journal of Geriatric Psychiatry</i> , <b>2016</b> , 31, 494-501	3.9	3
29	P2-252: Structural Brain Abnormalities Associated with Depression in Elderly with Type 2 Diabetes Differ by Haptoglobin Genotype <b>2016</b> , 12, P722-P722		1
28	P3-375: Neuropsychiatric Symptoms are Associated with Cognitive Function in Elderly with Type 2 Diabetes <b>2016</b> , 12, P993-P993		

27	P4-155: The Relationship of Depressive Symptoms with Cognitive Function is Different for Men and Women with Type 2 Diabetes <b>2016</b> , 12, P1074-P1074		
26	O4-12-06: Adherence to Type 2 Diabetes Medications is Associated with Better Cognitive Function <b>2016</b> , 12, P365-P365		
25	Long-term Variability in Glycemic Control Is Associated With White Matter Hyperintensities in APOE4 Genotype Carriers With Type 2 Diabetes. <i>Diabetes Care</i> , <b>2016</b> , 39, 1056-9	14.6	16
24	Waist circumference is correlated with poorer cognition in elderly type 2 diabetes women. <i>Alzheimer's and Dementia</i> , <b>2016</b> , 12, 925-9	1.2	16
23	P4-011: The Alzheimer's disease risk loci bin1 is associated with poorer episodic memory performance in cognitively normal type 2 diabetes elderly <b>2015</b> , 11, P769-P770		
22	P1-266: Trajectories of glycemic control are associated with depressive symptoms beyond cognitive function in nondemented very elderly with type 2 diabetes <b>2015</b> , 11, P457-P457		
21	P3-250: Haptoglobin genotype modulates the relationships of glycaemic control with cognitive function in elderly individuals with type 2 diabetes <b>2015</b> , 11, P726-P726		
20	The Israel Diabetes and Cognitive Decline (IDCD) study: Design and baseline characteristics. <i>Alzheimer's and Dementia</i> , <b>2014</b> , 10, 769-78	1.2	40
19	The TOMM40 poly-T rs10524523 variant is associated with cognitive performance among non-demented elderly with type 2 diabetes. <i>European Neuropsychopharmacology</i> , <b>2014</b> , 24, 1492-9	1.2	18
18	Efficacy of liraglutide in a real-life cohort. <i>Diabetes Therapy</i> , <b>2014</b> , 5, 193-206	3.6	15
17	Decreased motor function is associated with poorer cognitive function in elderly with type 2 diabetes. <i>Dementia and Geriatric Cognitive Disorders Extra</i> , <b>2014</b> , 4, 103-12	2.5	12
16	O2-09-04: TRAJECTORIES IN GLYCEMIC CONTROL OVER TIME ARE ASSOCIATED WITH COGNITIVE PERFORMANCE IN ELDERLY SUBJECTS WITH TYPE 2 DIABETES <b>2014</b> , 10, P184-P185		1
15	The association of duration of type 2 diabetes with cognitive performance is modulated by long-term glycemic control. <i>American Journal of Geriatric Psychiatry</i> , <b>2014</b> , 22, 1055-9	6.5	46
14	The ApoE4 genotype modifies the relationship of long-term glycemic control with cognitive functioning in elderly with type 2 diabetes. <i>European Neuropsychopharmacology</i> , <b>2014</b> , 24, 1303-8	1.2	14
13	Trajectories in glycemic control over time are associated with cognitive performance in elderly subjects with type 2 diabetes. <i>PLoS ONE</i> , <b>2014</b> , 9, e97384	3.7	43
12	The high cost of celiac disease in an Israeli Health Maintenance Organization. <i>Health Economics Review</i> , <b>2013</b> , 3, 23	2	3
11	Rites of passage. <i>Family Medicine</i> , <b>2013</b> , 45, 436-7	0.8	1
10	Exposed. <i>Patient Education and Counseling</i> , <b>2012</b> , 88, 350-1	3.1	

9	Quality of life and patient-perceived difficulties in the treatment of type 2 diabetes. <i>Review of Diabetic Studies</i> , <b>2012</b> , 9, 46-54	3.6	8
8	A matter of faith. <i>Patient Education and Counseling</i> , <b>2011</b> , 85, e6-7	3.1	
7	Factors associated with hypertensive patients' compliance with recommended lifestyle behaviors. <i>Israel Medical Association Journal</i> , <b>2011</b> , 13, 553-7	0.9	19
6	The perceived performance, barriers and solutions for the good preventive care of elderly people in Israel. <i>Quality in Primary Care</i> , <b>2010</b> , 18, 173-9		3
5	Pemphigus variant associated with penicillin use: a case-cohort study of 363 patients from Israel. <i>Archives of Dermatology</i> , <b>2007</b> , 143, 704-7		13
4	Differences between soldiers, with and without emotional distress, in number of primary care medical visits and type of presenting complaints. <i>Israel Medical Association Journal</i> , <b>2007</b> , 9, 90-3	0.9	2
3	The implementation of managed care for diabetes using medical informatics in a large Preferred Provider Organization. <i>Diabetes Research and Clinical Practice</i> , <b>2006</b> , 71, 290-8	7.4	46
2	Influence of school closure on the incidence of viral respiratory diseases among children and on health care utilization. <i>Pediatric Infectious Disease Journal</i> , <b>2004</b> , 23, 675-7	3.4	109
1	Systematic Inventive Thinking: a new tool for the analysis of complex problems in medical management. <i>Israel Medical Association Journal</i> , <b>2004</b> , 6, 67-9	0.9	1