

Arsenio Muoz de la Pea

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8311286/arsenio-munoz-de-la-pena-publications-by-year.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

172
papers

4,573
citations

39
h-index

57
g-index

174
ext. papers

4,789
ext. citations

5.2
avg, IF

5.31
L-index

#	Paper	IF	Citations
172	Binding the gap between experiments, statistics, and method comparison: A tutorial for computing limits of detection and quantification in univariate calibration for complex samples.. <i>Analytica Chimica Acta</i> , 2022 , 1209, 339342	6.6	2
171	Multi-way calibration for the quantification of polycyclic aromatic hydrocarbons in samples of environmental impact. <i>Microchemical Journal</i> , 2021 , 164, 106016	4.8	2
170	Untargeted classification for paprika powder authentication using visible [Near infrared spectroscopy (VIS-NIRS). <i>Food Control</i> , 2021 , 121, 107564	6.2	7
169	Second-order calibration in combination with fluorescence fibre-optic data modelling as a novel approach for monitoring the maturation stage of plums. <i>Chemometrics and Intelligent Laboratory Systems</i> , 2020 , 199, 103980	3.8	2
168	Non-destructive fluorescence spectroscopy combined with second-order calibration as a new strategy for the analysis of the illegal Sudan I dye in paprika powder. <i>Microchemical Journal</i> , 2020 , 154, 104539	4.8	5
167	Evolution of polyphenols content in plum fruits (<i>Prunus salicina</i>) with harvesting time by second-order excitation-emission fluorescence multivariate calibration. <i>Microchemical Journal</i> , 2020 , 158, 105299	4.8	7
166	Combination of fluorescence excitation emission matrices in polar and non-polar solvents to obtain three- and four- way arrays for classification of Tempranillo grapes according to maturation stage and hydric status. <i>Talanta</i> , 2019 , 199, 652-661	6.2	5
165	Four- and five-way excitation-emission luminescence-based data acquisition and modeling for analytical applications. A review. <i>Analytica Chimica Acta</i> , 2019 , 1083, 41-57	6.6	10
164	Front-Face Fluorescence Combined with Second-Order Multiway Classification, Based on Polyphenol and Chlorophyll Compounds, for Virgin Olive Oil Monitoring Under Different Photo- and Thermal-Oxidation Procedures. <i>Food Analytical Methods</i> , 2019 , 12, 1399-1411	3.4	10
163	Non-destructive Raman spectroscopy as a tool for measuring ASTA color values and Sudan I content in paprika powder. <i>Food Chemistry</i> , 2019 , 274, 187-193	8.5	24
162	Classification of pre-dyed textile fibers exposed to weathering and photodegradation by non-destructive excitation-emission fluorescence spectroscopy paired with discriminant unfolded-partial least squares. <i>Forensic Chemistry</i> , 2019 , 12, 25-32	2.8	6
161	Determination of pungency in spicy food by means of excitation-emission fluorescence coupled with second-order chemometric calibration. <i>Journal of Food Composition and Analysis</i> , 2018 , 67, 10-18	4.1	9
160	Detection and quantification of extra virgin olive oil adulteration by means of autofluorescence excitation-emission profiles combined with multi-way classification. <i>Talanta</i> , 2018 , 178, 751-762	6.2	50
159	Phenanthrene metabolites determination in human breast and cow milk by combining elution time-emission fluorescence data with multiway calibration. <i>Talanta</i> , 2018 , 188, 299-307	6.2	6
158	Modeling second-order data for classification issues: Data characteristics, algorithms, processing procedures and applications. <i>TrAC - Trends in Analytical Chemistry</i> , 2018 , 107, 151-168	14.6	22
157	Turn-on Fluorescence Hg ²⁺ Chemosensing Based in a Rhodamine 6G Derivative and Different Sensing Immobilization Approaches. <i>Advanced Engineering Forum</i> , 2018 , 29, 57-61	0.2	
156	Chemometric Discrimination Between Smoked and Non-Smoked Paprika Samples. Quantification of PAHs in Smoked Paprika by Fluorescence-U-PLS/RBL. <i>Food Analytical Methods</i> , 2017 , 10, 1128-1137	3.4	9

155	Nondestructive Total Excitation-Emission Fluorescence Microscopy Combined with Multi-Way Chemometric Analysis for Visually Indistinguishable Single Fiber Discrimination. <i>Analytical Chemistry</i> , 2016 , 88, 2967-75	7.8	18
154	Isocratic LC-DAD-FLD method for the determination of flavonoids in paprika samples by using a rapid resolution column and post-column pH change. <i>Talanta</i> , 2016 , 152, 15-22	6.2	8
153	Fluorescence properties of flavonoid compounds. Quantification in paprika samples using spectrofluorimetry coupled to second order chemometric tools. <i>Food Chemistry</i> , 2016 , 196, 1058-65	8.5	30
152	Highly Selective and Ultrasensitive Turn-on Luminescence Chemosensor for Mercury (II) Determination Based on the Rhodamine 6G Derivative FC1 and Au Nanoparticles. <i>Sensors</i> , 2016 , 16,	3.8	6
151	Rapid ultrasensitive chemometrics-fluorescence methodology to quantify fluoroquinolones antibiotics residues in surface water. <i>Journal of Water Chemistry and Technology</i> , 2016 , 38, 280-286	0.4	3
150	Excitation-Emission matrix fluorescence spectroscopy combined with MCR-ALS as a tool for the forensic analysis of similar and dissimilar sets of textile fiber extracts. <i>Analytical Methods</i> , 2016 , 8, 8314-8321	3.2	8
149	Combination of Liquid Chromatography with Multivariate Curve Resolution-Alternating Least-Squares (MCR-ALS) in the Quantitation of Polycyclic Aromatic Hydrocarbons Present in Paprika Samples. <i>Journal of Agricultural and Food Chemistry</i> , 2016 , 64, 8254-8262	5.7	14
148	Unfolded and Multiway Partial Least-Squares with Residual Multilinearization: Fundamentals. <i>Data Handling in Science and Technology</i> , 2015 , 29, 347-363	2.7	4
147	Unfolded and Multiway Partial Least-Squares with Residual Multilinearization. <i>Data Handling in Science and Technology</i> , 2015 , 29, 365-397	2.7	6
146	Novel augmented parallel factor model for four-way calibration of high-performance liquid chromatography-fluorescence excitation-Emission data. <i>Chemometrics and Intelligent Laboratory Systems</i> , 2015 , 141, 1-11	3.8	21
145	Modeling four and three-way fast high-performance liquid chromatography with fluorescence detection data for quantitation of fluoroquinolones in water samples. <i>Analytica Chimica Acta</i> , 2014 , 809, 37-46	6.6	38
144	Second- and higher-order data generation and calibration: a tutorial. <i>Analytica Chimica Acta</i> , 2014 , 806, 8-26	6.6	139
143	A second-order fluorimetric approach based on a boron dipyrromethene tetraamide derivative for Hg(II) chemosensing in water and fish samples. <i>Analytical Methods</i> , 2014 , 6, 8535-8543	3.2	3
142	A sensing microfibre mat produced by electrospinning for the turn-on luminescence determination of Hg ²⁺ in water samples. <i>Sensors and Actuators B: Chemical</i> , 2014 , 195, 8-14	8.5	20
141	Hg ²⁺ -selective sensing film based on the incorporation of a rhodamine 6G derivative into a novel hydrophilic water-insoluble copolymer. <i>Analytical Methods</i> , 2013 , 5, 6642	3.2	13
140	Rhodamine and BODIPY chemodosimeters and chemosensors for the detection of Hg ²⁺ , based on fluorescence enhancement effects. <i>Analytical Methods</i> , 2013 , 5, 30-49	3.2	136
139	Photoinduced electron transfer fluorometric Hg(II) chemosensor based on a BODIPY armed with a tetrapod receptor. <i>Talanta</i> , 2013 , 117, 288-96	6.2	20
138	Automatic Web-Based Grading System: Application in an Advanced Instrumental Analysis Chemistry Laboratory. <i>Journal of Chemical Education</i> , 2013 , 90, 308-314	2.4	4

137	Four-way multivariate calibration using ultra-fast high-performance liquid chromatography with fluorescence excitation-emission detection. Application to the direct analysis of chlorophylls a and b and pheophytins a and b in olive oils. <i>Chemometrics and Intelligent Laboratory Systems</i> , 2013 , 125, 121-131	3.8	38
136	Modelling ambient ozone in an urban area using an objective model and geostatistical algorithms. <i>Atmospheric Environment</i> , 2012 , 63, 86-93	5.3	6
135	Multiresidue method for the control of pesticide residues in tomatoes and derived products. <i>Analytical Methods</i> , 2012 , 4, 2543	3.2	1
134	A novel nylon membrane-rhodamine 6G spirocyclic phenylthiosemicarbazide derivative system as a fluorimetric probe for mercury(II) ion. <i>Analytical Methods</i> , 2012 , 4, 2002	3.2	18
133	Enhanced MCR-ALS modeling of HPLC with fast scan fluorimetric detection second-order data for quantitation of metabolic disorder marker pteridines in urine. <i>Talanta</i> , 2011 , 85, 2368-74	6.2	30
132	Unfolded partial least-squares with residual quadrilinearization: A new multivariate algorithm for processing five-way data achieving the second-order advantage. Application to fourth-order excitation-emission-kinetic-pH fluorescence analytical data. <i>Chemometrics and Intelligent Laboratory Systems</i> , 2011 , 119, 176-185	3.8	39
131	Determination of marker pteridines in urine by HPLC with fluorimetric detection and second-order multivariate calibration using MCR-ALS. <i>Analytical and Bioanalytical Chemistry</i> , 2011 , 399, 2123-35	4.4	37
130	Separation and determination of 11 marker pteridines in human urine by liquid chromatography and fluorimetric detection. <i>Journal of Separation Science</i> , 2011 , 34, 1283-92	3.4	16
129	Second-order and higher-order multivariate calibration methods applied to non-multilinear data using different algorithms. <i>TrAC - Trends in Analytical Chemistry</i> , 2011 , 30, 607-617	14.6	86
128	Fluorescent determination of Hg ²⁺ in water and fish samples using a chemodosimeter based in a Rhodamine 6G derivative and a portable fiber-optic spectrofluorimeter. <i>Applied Spectroscopy</i> , 2010 , 64, 520-7	3.1	20
127	Second-order data obtained by time-resolved room temperature phosphorescence. A new approach for PARAFAC multicomponent analysis. <i>Journal of Fluorescence</i> , 2009 , 19, 345-52	2.4	6
126	Analysis of antibiotics in fish samples. <i>Analytical and Bioanalytical Chemistry</i> , 2009 , 395, 987-1008	4.4	92
125	Determination of danofloxacin in milk combining second-order calibration and standard addition method using excitation-emission fluorescence data. <i>Food Chemistry</i> , 2009 , 113, 1260-1265	8.5	22
124	Second-order multivariate calibration procedures applied to high-performance liquid chromatography coupled to fast-scanning fluorescence detection for the determination of fluoroquinolones. <i>Journal of Chromatography A</i> , 2009 , 1216, 4868-76	4.5	52
123	Determination of marker pteridins and biopterin reduced forms, tetrahydrobiopterin and dihydrobiopterin, in human urine, using a post-column photoinduced fluorescence liquid chromatographic derivatization method. <i>Analytica Chimica Acta</i> , 2009 , 648, 113-22	6.6	31
122	LC determination of biopterin reduced forms by UV-photogeneration of biopterin and fluorimetric detection. <i>Talanta</i> , 2008 , 77, 844-851	6.2	18
121	Flow-through photochemically induced fluorescence optosensor for the determination of linuron. <i>Talanta</i> , 2008 , 77, 852-857	6.2	30
120	Nonlinear four-way kinetic-excitation-emission fluorescence data processed by a variant of parallel factor analysis and by a neural network model achieving the second-order advantage: malonaldehyde determination in olive oil samples. <i>Analytical Chemistry</i> , 2008 , 80, 7248-56	7.8	39

119	Optimization of Verapamil drug analysis by excitation-emission fluorescence in combination with second-order multivariate calibration. <i>Journal of Fluorescence</i> , 2008 , 18, 1065-76	2.4	12
118	Photoinduced fluorimetric determination of folic acid and 5-methyltetrahydrofolic acid in serum using the kinetic evolution of the emission spectra accomplished with multivariate second-order calibration methods. <i>Analytical and Bioanalytical Chemistry</i> , 2008 , 391, 827-35	4.4	17
117	On line photochemically induced excitation-emission-kinetic four-way data: analytical application for the determination of folic acid and its two main metabolites in serum by U-PLS and N-PLS/residual trilinearization (RTL) calibration. <i>Analytica Chimica Acta</i> , 2008 , 622, 94-103	6.6	29
116	Separation of fifteen quinolones by high performance liquid chromatography: application to pharmaceuticals and ofloxacin determination in urine. <i>Journal of Separation Science</i> , 2007 , 30, 1242-9	3.4	33
115	Second- and third-order multivariate calibration: data, algorithms and applications. <i>TrAC - Trends in Analytical Chemistry</i> , 2007 , 26, 752-765	14.6	272
114	Fluorimetric determination of sulphaguanidine and sulphamethoxazole by host-guest complexation in beta-cyclodextrin and partial least squares calibration. <i>Journal of Fluorescence</i> , 2007 , 17, 309-18	2.4	18
113	Resolution of ofloxacin-ciprofloxacin and ofloxacin-norfloxacin binary mixtures by flow-injection chemiluminescence in combination with partial least squares multivariate calibration. <i>Journal of Fluorescence</i> , 2007 , 17, 481-91	2.4	29
112	Evaluation of unfolded-partial least-squares coupled to residual trilinearization for four-way calibration of folic acid and methotrexate in human serum samples. <i>Talanta</i> , 2007 , 72, 1261-8	6.2	26
111	A chemometric sensor for determining sulphaguanidine residues in honey samples. <i>Talanta</i> , 2007 , 73, 304-13	6.2	15
110	Multiway partial least-squares coupled to residual trilinearization: a genuine multidimensional tool for the study of third-order data. Simultaneous analysis of procaine and its metabolite p-aminobenzoic acid in equine serum. <i>Analytical Chemistry</i> , 2007 , 79, 6949-58	7.8	53
109	Four-way calibration applied to the simultaneous determination of folic acid and methotrexate in urine samples. <i>Analytical and Bioanalytical Chemistry</i> , 2006 , 385, 1289-97	4.4	29
108	HPLC determination of ciprofloxacin, cloxacillin, and ibuprofen drugs in human urine samples. <i>Journal of Separation Science</i> , 2006 , 29, 1969-76	3.4	25
107	Second-order advantage achieved by unfolded-partial least-squares/residual bilinearization modeling of excitation-emission fluorescence data presenting inner filter effects. <i>Analytical Chemistry</i> , 2006 , 78, 8051-8	7.8	63
106	Second-order calibration of excitation-emission matrix fluorescence spectra for the determination of N-phenylanthranilic acid derivatives. <i>Applied Spectroscopy</i> , 2006 , 60, 330-8	3.1	22
105	Determination of fluoroquinolones in urine and serum by using high performance liquid chromatography and multiemission scan fluorimetric detection. <i>Talanta</i> , 2006 , 68, 1215-21	6.2	47
104	Stopped-flow and kinetic-fluorimetric determination of quinalphos in water samples. <i>Talanta</i> , 2006 , 69, 397-402	6.2	6
103	Simultaneous determination of flufenamic and meclofenamic acids in human urine samples by second-order multivariate parallel factor analysis (PARAFAC) calibration of micellar-enhanced excitation-emission fluorescence data. <i>Analytica Chimica Acta</i> , 2006 , 569, 250-259	6.6	32
102	Trilinear least-squares and unfolded-PLS coupled to residual trilinearization: New chemometric tools for the analysis of four-way instrumental data. <i>Chemometrics and Intelligent Laboratory Systems</i> , 2006 , 80, 77-86	3.8	82

101	Strategies for solving matrix effects in the analysis of sulfathiazole in honey samples using three-way photochemically induced fluorescence data. <i>Talanta</i> , 2005 , 65, 806-13	6.2	27
100	HPLC determination of enoxacin, ciprofloxacin, norfloxacin and ofloxacin with photoinduced fluorimetric (PIF) detection and multiemission scanning: application to urine and serum. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2005 , 822, 185-93	3.2	79
99	Determinations of fluoroquinolones and nonsteroidal anti-inflammatory drugs in urine by extractive spectrophotometry and photoinduced spectrofluorimetry using multivariate calibration. <i>Analytical Biochemistry</i> , 2005 , 347, 275-86	3.1	36
98	Determination of triamterene in pharmaceutical formulations and of triamterene and its main metabolite hydroxytriamterene sulfate in urine using solid-phase and aqueous solution luminescence. <i>Analytica Chimica Acta</i> , 2005 , 538, 77-84	6.6	20
97	Photoinduced spectrofluorimetric determination of fluoroquinolones in human urine by using three- and two-way spectroscopic data and multivariate calibration. <i>Analytica Chimica Acta</i> , 2005 , 531, 257-266	6.6	42
96	Complexation Study and Spectrofluorimetric Determination of Pipemidic Acid with β -Cyclodextrin. <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 2005 , 51, 137-142		5
95	Determination of carbamazepine in serum and pharmaceutical preparations using immobilization on a nylon support and fluorescence detection. <i>Analytica Chimica Acta</i> , 2004 , 506, 161-170	6.6	72
94	Second-order advantage achieved with four-way fluorescence excitation-emission-kinetic data processed by parallel factor analysis and trilinear least-squares. Determination of methotrexate and leucovorin in human urine. <i>Analytical Chemistry</i> , 2004 , 76, 5657-66	7.8	99
93	Partial least squares multicomponent fluorimetric determination of fluoroquinolones in human urine samples. <i>Talanta</i> , 2004 , 62, 853-60	6.2	48
92	Two multivariate strategies applied to three-way kinetic spectrophotometric data for the determination of mixtures of the pesticides carbaryl and chlorpyrifos. <i>Applied Spectroscopy</i> , 2004 , 58, 83-90	3.1	41
91	Interference-free analysis using three-way fluorescence data and the parallel factor model. Determination of fluoroquinolone antibiotics in human serum. <i>Analytical Chemistry</i> , 2003 , 75, 2640-6	7.8	88
90	Determination of carbendazim, thiabendazole and fuberidazole using a net analyte signal-based method. <i>Talanta</i> , 2003 , 59, 1107-16	6.2	36
89	Monitoring of phenylurea and propanil herbicides in river water by solid-phase-extraction high performance liquid chromatography with photoinduced-fluorimetric detection. <i>Talanta</i> , 2003 , 60, 279-85	6.2	48
88	Determination of antitubercular drugs by micellar electrokinetic capillary chromatography (MEKC). <i>Analytical and Bioanalytical Chemistry</i> , 2002 , 374, 432-6	4.4	24
87	Comparative study of net analyte signal-based methods and partial least squares for the simultaneous determination of amoxicillin and clavulanic acid by stopped-flow kinetic analysis. <i>Analytica Chimica Acta</i> , 2002 , 463, 75-88	6.6	43
86	High-performance liquid chromatographic determination of phenylureas by photochemically-induced fluorescence detection. <i>Journal of Chromatography A</i> , 2002 , 950, 287-91	4.5	17
85	An Investigation of Inclusion Complexes of Cyclodextrins with Phenylurea Herbicides by Photochemically-Induced Fluorescence. Analytical Applications. <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 2002 , 42, 61-70		22
84	Stopped-flow fluorimetric determination of amoxicillin and clavulanic acid by partial least-squares multivariate calibration. <i>Talanta</i> , 2002 , 56, 635-42	6.2	14

83	Determination of antitubercular drugs in urine and pharmaceuticals by LC using a gradient flow combined with programmed diode array photometric detection. <i>Talanta</i> , 2002 , 58, 273-80	6.2	30
82	Selection of the wavelength range and spectrophotometric determination of leucovorin and methotrexate in human serum by a net analyte signal based method. <i>Talanta</i> , 2002 , 58, 255-63	6.2	25
81	Comparative study of partial least squares and a modification of hybrid linear analysis calibration in the simultaneous spectrophotometric determination of rifampicin, pyrazinamide and isoniazid. <i>Analytica Chimica Acta</i> , 2001 , 427, 129-136	6.6	40
80	HOST-GUEST ROOM TEMPERATURE PHOSPHORESCENCE OF 1-NAPHTHALENACETIC ACID INCLUDED IN β -CYCLODEXTRIN IN PRESENCE OF 1,3-DIBROMOPROPANE. <i>Analytical Letters</i> , 2001 , 34, 2391-2401	2.2	
79	Room-Temperature Phosphorescence of 1-Bromonaphthalene upon Formation of Beta-Cyclodextrin Ternary Complexes with Alcohols and Surfactants: Optimization of Analytical Figures of Merit by Rigorous Equilibrium Studies. <i>Applied Spectroscopy</i> , 2001 , 55, 496-503	3.1	9
78	Optimization of the room-temperature phosphorescence of the 6-bromo-2-naphthol β -cyclodextrin system in aqueous solution. <i>Talanta</i> , 2000 , 51, 949-955	6.2	9
77	Complexation of antibacterial quinolonic acid and cinolonic derivatives with Zn(II) and Al(III): application to their determination in human urine. <i>Analyst, The</i> , 2000 , 125, 1471-6	5	15
76	Room temperature phosphorescence in cyclodextrins. Analytical applications. <i>Analisis - European Journal of Analytical Chemistry</i> , 2000 , 28, 670-678		15
75	Determination of theophylline in blood serum by UV spectrophotometry and partial least-squares (PLS-1) calibration. <i>Analytica Chimica Acta</i> , 1999 , 384, 95-103	6.6	38
74	Room temperature phosphorescence of 1-naphtalenacetamide included in beta-cyclodextrin in presence of 1,3-dibromopropane. <i>Talanta</i> , 1999 , 48, 15-21	6.2	10
73	Stopped-flow determination of dipyridamole in pharmaceutical preparations by micellar-stabilized room temperature phosphorescence. <i>Talanta</i> , 1999 , 48, 1061-73	6.2	22
72	Usefulness of micellar media for the quantitative analysis of phenylurea herbicides in water by photochemically-induced fluorescence. <i>Analisis - European Journal of Analytical Chemistry</i> , 1999 , 27, 857-863		19
71	Room-temperature phosphorescence of acenaphthene in aerated solutions in the presence of bromoalcohols and β -cyclodextrin. <i>Analytica Chimica Acta</i> , 1998 , 370, 199-205	6.6	40
70	Simultaneous fluorometric determination of nalidixic acid and 7-hydroxymethylnalidixic acid by partial least squares calibration. <i>Talanta</i> , 1998 , 45, 899-907	6.2	21
69	Cross-sections of spectrochromatograms for the resolution of overlapping peaks in diode-array high-performance liquid-chromatography. <i>Talanta</i> , 1998 , 46, 1329-40	6.2	7
68	Determination of nafronyl in pharmaceutical preparations by means of stopped-flow micellar-stabilized room temperature phosphorescence. <i>Analyst, The</i> , 1998 , 123, 2285-90	5	14
67	Simultaneous determination of nafcillin and methicillin by different fluorimetric techniques using partial least-squares calibration. <i>Analyst, The</i> , 1998 , 123, 1073-7	5	15
66	Kinetic Fluorometric Determination of Malonaldehyde Based on the Hantzsch Reaction: Application to Olive Oil Analysis. <i>Journal of Agricultural and Food Chemistry</i> , 1997 , 45, 172-177	5.7	3

65	Spectroscopic Studies of the Interaction of 1,4-Diphenyl-1,3-Butadiene with β and γ -Cyclodextrins. <i>Applied Spectroscopy</i> , 1997 , 51, 153-159	3.1	4
64	Spectrofluorimetric Study of the Inclusion Complex of 7-Hydroxymethylnalidixic Acid with β -Cyclodextrin in Aqueous Solution. <i>Applied Spectroscopy</i> , 1997 , 51, 684-688	3.1	13
63	Resolution of overlapping peaks in HPLC with diode array detection by application of partial least squares calibration to cross-sections of spectrochromatograms. <i>Analytica Chimica Acta</i> , 1997 , 348, 177-185	6.6	25
62	Synchronous fluorimetric determination of salicylic acid and diflunisal in human serum using partial least-squares calibration. <i>Talanta</i> , 1996 , 43, 1349-56	6.2	21
61	Synthesis of a water-soluble chiral N-acylcalix(4)arene amino acid derivative. <i>Tetrahedron Letters</i> , 1996 , 37, 5841-5844	2	49
60	Photometric and fluorimetric study of the acid-base behavior of 2,2'-diquinolyl and 2,2'-terpyridyl. <i>Journal of Fluorescence</i> , 1996 , 6, 97-102	2.4	1
59	pH-Induced Difference Spectrophotometry in the Analysis of Binary Mixtures.. <i>Analytical Letters</i> , 1996 , 29, 2525-2540	2.2	6
58	Photochemically induced fluorescence investigation of α -Cyclodextrin: Azure A inclusion complex and determination of analytical parameters. <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 1995 , 22, 235-247		20
57	Resolution of ternary mixtures of salicylic, salicylic and gentisic acids by partial least squares and principal component regression: Optimization of the scanning path in the excitation-emission matrices. <i>Fresenius Journal of Analytical Chemistry</i> , 1995 , 351, 571-576		19
56	Influence of Triethanolamine on Aqueous β -Cyclodextrin/Pyrene Complexes. <i>Applied Spectroscopy</i> , 1995 , 49, 520-525	3.1	1
55	Simultaneous fluorimetric determination of acetylsalicylic acid metabolites in urine by partial least squares multivariate calibration. <i>Fresenius Journal of Analytical Chemistry</i> , 1995 , 353, 211-214		10
54	Simultaneous fluorimetric determination of acetylsalicylic acid metabolites in urine by partial least squares multivariate calibration. <i>Analytical and Bioanalytical Chemistry</i> , 1995 , 353, 211-4	4.4	1
53	Basic studies of the influence of β -Cyclodextrin and 2-hydroxypropyl- β -Cyclodextrin on the photo-oxidation reaction of phenothiazine in inclusion complexes, using fluorescence detection. <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 1994 , 18, 69-79		4
52	Determination of binary mixtures of sulfonamides by photochemically induced fluorescence using partial least squares multivariate calibration. <i>Analyst, The</i> , 1994 , 119, 1177-1181	5	25
51	Fluorimetric Determination of Sulfamethoxazole in Pharmaceutical Preparations in Combination with Trimethoprim by Inclusion in β -Cyclodextrin/Urea. <i>Analytical Letters</i> , 1994 , 27, 1893-1906	2.2	20
50	Determination of 1-naphthylamine and the related pesticides, naptalam and antu, in river-water by high-performance liquid chromatography. Application to the study of the degradation processes of naptalam. <i>Analyst, The</i> , 1994 , 119, 1151-1155	5	15
49	Spectrophotometric determination of mixtures of iron(III) and manganese(II) by complexation with 3-indolylacetohydroxamic acid and principal component regression multivariate calibration. <i>Analyst, The</i> , 1994 , 119, 1537-1540	5	6
48	Spectrofluorimetric determination of nalidixic acid based on host-guest complexation with β -Cyclodextrin. <i>Analyst, The</i> , 1994 , 119, 1215-1219	5	40

47	Influence of alcohols on the .beta.-cyclodextrin/acridine complex. <i>Journal of the American Chemical Society</i> , 1993 , 115, 292-298	16.4	62
46	Multicomponent determination of flavour enhancers in food preparations by partial least squares and principal component regression modelling of spectrophotometric data. <i>Analyst, The</i> , 1993 , 118, 807-813	5.1	55
45	Host-guest stabilized room temperature phosphorescence in beta-cyclodextrin/ bromoalcohol solutions from 2-naphthyl-oxy-acetic acid and 1-naphthyl-acetic acid. <i>Talanta</i> , 1993 , 40, 1657-64	6.2	19
44	Simultaneous Determination of Flavor Enhancers Inosine 5'-Monophosphate and Guanosine 5'-Monophosphate in Food Preparations by Derivative Spectrophotometry. <i>Journal of AOAC INTERNATIONAL</i> , 1993 , 76, 754-759	1.7	3
43	Simultaneous determination of 2-furfuraldehyde, 5-hydroxymethylfurfuraldehyde and malonaldehyde in mixtures by derivative spectrophotometry and partial least-squares analysis. <i>Analytica Chimica Acta</i> , 1993 , 276, 141-149	6.6	48
42	Absorptiometric and spectrofluorimetric study of the inclusion complexes of 2-naphthyloxyacetic acid and 1-naphthylacetic acid with β -cyclodextrin in aqueous solution. <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 1993 , 15, 131-143		61
41	Elimination of spectral interferences in the reaction of 2-thiobarbituric acid with malonaldehyde, 2-furfuraldehyde and 5-hydroxymethyl-2-furfuraldehyde by partial least squares multivariate calibration (PLS). <i>Fresenius Journal of Analytical Chemistry</i> , 1993 , 347, 371-375		4
40	Spectroscopic Studies of the Interaction of Tert-Butylamine and n-Propylamine with the β -Cyclodextrin: Pyrene Complex. <i>Applied Spectroscopy</i> , 1993 , 47, 277-282	3.1	31
39	Analytical Chemistry in the EC. Research in Analytical Chemistry in Spain today, through an evaluation of the scientific literature. <i>Analytical Proceedings</i> , 1992 , 29, 517-520		5
38	Simultaneous determination of 2-furfuraldehyde and 5-(hydroxymethyl)-2-furfuraldehyde by derivative spectrophotometry. <i>Journal of Agricultural and Food Chemistry</i> , 1992 , 40, 1022-1025	5.7	28
37	Determination of formation constants for .beta.-cyclodextrin complexes of anthracene and pyrene using reversed-phase liquid chromatography. <i>Analytical Chemistry</i> , 1992 , 64, 484-489	7.8	41
36	Complexation of Doxorubicin with α and β -Cyclodextrins. <i>Applied Spectroscopy</i> , 1992 , 46, 652-658	3.1	44
35	Retention behavior of β -cyclodextrin complexes of anthracene and pyrene using reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 1992 , 594, 37-43	4.5	13
34	Simultaneous determination of pesticides by multivariate spectral analysis and derivative spectrophotometry. <i>Analytica Chimica Acta</i> , 1992 , 258, 47-53	6.6	37
33	Simultaneous determination of propranolol and hydralazine by derivative synchronous spectrofluorimetry. <i>Analytica Chimica Acta</i> , 1991 , 255, 317-323	6.6	24
32	Cyclodextrin-induced fluid solution room-temperature phosphorescence from acenaphthene in the presence of 2-bromoethanol. <i>Analytica Chimica Acta</i> , 1991 , 255, 351-357	6.6	53
31	Spectral characterization of β -Cyclodextrin : Triton X-100 complexes. <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 1991 , 10, 471-484		57
30	Characterization of the .beta.-cyclodextrin/acridine complex. <i>The Journal of Physical Chemistry</i> , 1991 , 95, 4897-4902		56

29	Analysis of mixtures of oxytetracycline and riboflavine by first-derivative synchronous spectrofluorimetry. <i>Analyst, The</i> , 1991 , 116, 291-296	5	6
28	Alcohol size as a factor in the ternary complexes formed with pyrene and .beta.-cyclodextrin. <i>Journal of the American Chemical Society</i> , 1991 , 113, 1572-1577	16.4	150
27	Solution studies of .beta.-cyclodextrin-pyrene complexes under reversed-phase liquid chromatographic conditions: effect of alcohols as mobile-phase comodifiers. <i>Analytical Chemistry</i> , 1991 , 63, 1018-1023	7.8	36
26	Influence of alcohol addition on the .gamma.-CD:pyrene complex. <i>The Journal of Physical Chemistry</i> , 1991 , 95, 6701-6706		55
25	Application of time-domain differentiation of chromatographic peaks in liquid chromatography. <i>Analytica Chimica Acta</i> , 1990 , 234, 263-267	6.6	13
24	Analysis of Mixtures of Doxycycline and Oxytetracycline in Pharmaceutical Preparations by First Derivative Fluorimetry. <i>Analytical Letters</i> , 1990 , 23, 863-876	2.2	26
23	Determination of salicylic acid and its metabolites in urine by derivative synchronous spectrofluorimetry. <i>Analyst, The</i> , 1990 , 115, 1007-11	5	43
22	Studies on Ground and Excited State Prototropic Equilibria Sodium 1,3-Dihydroxy-4-aminoanthraquinone-2-sulphonate. <i>Analytical Letters</i> , 1989 , 22, 2371-2380	2.2	1
21	Simultaneous determination of molybdenum and tungsten by first-derivative synchronous spectrofluorimetry. <i>Analyst, The</i> , 1989 , 114, 1297-1301	5	13
20	Development of a low-cost interface for coupling a microcomputer with a fluorescence spectrophotometer. Computer-assisted fluorimetry. <i>Computers & Chemistry</i> , 1988 , 12, 213-217		7
19	Determination of Aluminium with 8-Hydroxyquinoline-5-Sulfonic Acid in Presence of A Cationic Surfactant by First and Second Derivative Synchronous Fluorimetry. <i>Analytical Letters</i> , 1988 , 21, 1457-1468	2.2	9
18	Simultaneous determination of salicylic and salicyluric acids in urine by first-derivative synchronous fluorescence spectroscopy. <i>Analytical Chemistry</i> , 1988 , 60, 2493-6	7.8	40
17	Simultaneous determination of cobalt and nickel by first-derivative spectrophotometry. <i>Analyst, The</i> , 1988 , 113, 1439-1442	5	19
16	Spectrofluorimetric determination of vanadium(V) in petroleum crudes and basic slag with Nuclear Fast Red. <i>Analyst, The</i> , 1988 , 113, 987-990	5	5
15	Simultaneous determination of aluminium and zinc by first- and second-derivative synchronous fluorimetry. <i>Analyst, The</i> , 1988 , 113, 1435-1438	5	14
14	Spectrofluorimetric determination of beryllium in rocks, alloys and steels with nuclear fast red. <i>Analyst, The</i> , 1987 , 112, 645-648	5	6
13	Simultaneous determination of beryllium and magnesium with 1-hydroxy-2-carboxyanthraquinone by first-derivative spectrophotometry. <i>Analyst, The</i> , 1987 , 112, 1391-1394	5	14
12	Spectrofluorimetric determination of boron in plants with quinizarin-2-sulphonic acid. <i>Analyst, The</i> , 1987 , 112, 913-915	5	12

11	1-Hydroxy-2-carboxyanthraquinone as a fluorometric reagent for lanthanum. <i>Microchemical Journal</i> , 1987 , 36, 79-83	4.8	2
10	Spectrophotometric and First Derivative Spectrophotometric Determination of Magnesium with 1-Hydroxy-2-Carboxyanthraquinone. <i>Analytical Letters</i> , 1986 , 19, 1349-1358	2.2	3
9	Phototautomerism in the lowest excited singlet state of 1-hydroxy-2-carboxyanthraquinone. <i>Talanta</i> , 1986 , 33, 923-6	6.2	1
8	Determination of magnesium by spectrofluorimetry and synchronous scanning first and second derivative spectrofluorimetry with 2-quinizarinsulphonate. <i>Mikrochimica Acta</i> , 1985 , 87, 361-368	5.8	7
7	Kinetics and Equilibria of Proton Exchange in the Lowest Excited Singlet State of 1-Hydroxyfluorene in Aqueous Solutions. <i>Analytical Letters</i> , 1985 , 18, 2489-2495	2.2	1
6	Spectrophotometric study of pseudopurpurin-Pd(II) dimer complex. Spectrophotometric determination of traces of Pd(II). <i>Microchemical Journal</i> , 1984 , 29, 275-281	4.8	1
5	Spectrophotometric determination of BrO ₃ ⁻ using the 1,2,4-trihydroxyanthraquinone-3-carboxylic acid as reagent. <i>Microchemical Journal</i> , 1984 , 29, 19-25	4.8	4
4	Spectrofluorimetric determination of Ga(III) with 1-hydroxy-2-carboxyanthraquinone. <i>Mikrochimica Acta</i> , 1984 , 84, 79-86	5.8	3
3	Spectrofluorimetric determination of BrO ₃ ⁻ by oxidation of 1,2,4-trihydroxyanthraquinone-3-carboxylic acid. <i>Mikrochimica Acta</i> , 1984 , 83, 75-83	5.8	1
2	Spectrofluorimetric determination of aluminium(III) in Portland cement and aluminium bronze with 1-hydroxy-2-carboxyanthraquinone. <i>Analyst, The</i> , 1984 , 109, 1135-1137	5	4
1	Spectrofluorimetric Determination of Y(III) with 1-Hydroxy-2-Carboxyanthraquinone. <i>Analytical Letters</i> , 1984 , 17, 497-505	2.2	5