

Francesco Perticone

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8330958/francesco-perticone-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

239
papers

8,336
citations

45
h-index

83
g-index

251
ext. papers

9,333
ext. citations

5.3
avg, IF

5.4
L-index

#	Paper	IF	Citations
239	Effects of Sacubitril-Valsartan on Clinical, Echocardiographic, and Polygraphic Parameters in Patients Affected by Heart Failure With Reduced Ejection Fraction and Sleep Apnea.. <i>Frontiers in Cardiovascular Medicine</i> , 2022 , 9, 861663	5.4	0
238	Insulin-like growth factor-1 (IGF-1) as predictor of cardiovascular mortality in heart failure patients: data from the T.O.S.CA. registry.. <i>Internal and Emergency Medicine</i> , 2022 , 1	3.7	0
237	Continuous Positive Airway Pressure Improves Renal Function in Obese Patients With Obstructive Sleep Apnea Syndrome. <i>Frontiers in Medicine</i> , 2021 , 8, 642086	4.9	1
236	COPD significantly increases cerebral and cardiovascular events in hypertensives. <i>Scientific Reports</i> , 2021 , 11, 7884	4.9	2
235	Hyperglycemia at admission, comorbidities, and in-hospital mortality in elderly patients hospitalized in internal medicine wards: data from the RePoSI Registry. <i>Acta Diabetologica</i> , 2021 , 58, 1225-1236	3.9	2
234	Pattern of comorbidities and 1-year mortality in elderly patients with COPD hospitalized in internal medicine wards: data from the RePoSI Registry. <i>Internal and Emergency Medicine</i> , 2021 , 16, 389-400	3.7	7
233	The multifaceted spectrum of liver cirrhosis in older hospitalised patients: analysis of the REPOSI registry. <i>Age and Ageing</i> , 2021 , 50, 498-504	3	0
232	Endothelial dysfunction and C-reactive protein predict the incidence of heart failure in hypertensive patients. <i>ESC Heart Failure</i> , 2021 , 8, 399-407	3.7	3
231	One Hour-Post-load Plasma Glucose ≥ 55 mg/dl in Healthy Glucose Normotolerant Subjects Is Associated With Subcortical Brain MRI Alterations and Impaired Cognition: A Pilot Study. <i>Frontiers in Aging Neuroscience</i> , 2021 , 13, 608736	5.3	0
230	Multiple hormonal and metabolic deficiency syndrome predicts outcome in heart failure: the T.O.S.CA. Registry. <i>European Journal of Preventive Cardiology</i> , 2021 ,	3.9	6
229	Predictors of Renal Function Worsening in Patients with Chronic Obstructive Pulmonary Disease (COPD): A Multicenter Observational Study. <i>Nutrients</i> , 2021 , 13,	6.7	3
228	Impact of implementing a Choosing Wisely educational intervention into clinical practice: The CW-SIMI study (a multicenter-controlled study). <i>European Journal of Internal Medicine</i> , 2021 , 93, 71-77	3.9	2
227	Endothelial progenitor cells predict vascular damage progression in naive hypertensive patients according to sex. <i>Hypertension Research</i> , 2021 , 44, 1451-1461	4.7	1
226	Non-alcoholic fatty liver disease is associated with cardiovascular disease in subjects with different glucose tolerance. <i>Diabetes/Metabolism Research and Reviews</i> , 2020 , 36, e3333	7.5	9
225	Reduction in Global Myocardial Glucose Metabolism in Subjects With 1-Hour Postload Hyperglycemia and Impaired Glucose Tolerance. <i>Diabetes Care</i> , 2020 , 43, 669-676	14.6	7
224	Association of different oral anticoagulants use with renal function worsening in patients with atrial fibrillation: A multicentre cohort study. <i>British Journal of Clinical Pharmacology</i> , 2020 , 86, 2455-2463	3.8	14
223	Association between Serum Mg Concentrations and Cardiovascular Organ Damage in a Cohort of Adult Subjects. <i>Nutrients</i> , 2020 , 12,	6.7	2

222	Alkaline phosphatase affects renal function in never-treated hypertensive patients: effect modification by age. <i>Scientific Reports</i> , 2020 , 10, 9847	4.9	4
221	Metabolic and Cognitive Effects of Ranolazine in Type 2 Diabetes Mellitus: Data from an in vivo Model. <i>Nutrients</i> , 2020 , 12,	6.7	10
220	Different Patterns of Left Ventricular Hypertrophy in Metabolically Healthy and Insulin-Resistant Obese Subjects. <i>Nutrients</i> , 2020 , 12,	6.7	8
219	HDL cholesterol is an independent predictor of β-cell function decline and incident type 2 diabetes: A longitudinal study. <i>Diabetes/Metabolism Research and Reviews</i> , 2020 , 36, e3289	7.5	9
218	Comorbidity does not mean clinical complexity: evidence from the RePoSI register. <i>Internal and Emergency Medicine</i> , 2020 , 15, 621-628	3.7	7
217	Cardiometabolic risk prevention strategies: the importance of sharing experiences between Mediterranean countries. <i>Internal and Emergency Medicine</i> , 2020 , 15, 543-548	3.7	1
216	Prevalence and Impact of Nonalcoholic Fatty Liver Disease in Atrial Fibrillation. <i>Mayo Clinic Proceedings</i> , 2020 , 95, 513-520	6.4	5
215	Pain and Frailty in Hospitalized Older Adults. <i>Pain and Therapy</i> , 2020 , 9, 727-740	3.6	8
214	Serum γ-Glutamyltransferase Concentration Predicts Endothelial Dysfunction in Naïve Hypertensive Patients. <i>Biomedicines</i> , 2020 , 8,	4.8	2
213	Uric Acid and Vascular Damage in Essential Hypertension: Role of Insulin Resistance. <i>Nutrients</i> , 2020 , 12,	6.7	9
212	Management of chronic ischemic heart disease into internal medicine and geriatric departments in Italy. <i>Internal and Emergency Medicine</i> , 2020 , 15, 151-154	3.7	
211	Incidence and Recurrence of Portal Vein Thrombosis in Cirrhotic Patients. <i>Thrombosis and Haemostasis</i> , 2019 , 119, 496-499	7	18
210	Individuals With Prediabetes Display Different Age-Related Pathophysiological Characteristics. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019 , 104, 2911-2924	5.6	8
209	Antibiotic use and associated factors in a large sample of hospitalised older people. <i>Journal of Global Antimicrobial Resistance</i> , 2019 , 19, 167-172	3.4	3
208	Serum Albumin Is Inversely Associated With Portal Vein Thrombosis in Cirrhosis. <i>Hepatology Communications</i> , 2019 , 3, 504-512	6	31
207	Elevated 1-h post-load plasma glucose is associated with right ventricular morphofunctional parameters in hypertensive patients. <i>Endocrine</i> , 2019 , 64, 525-535	4	2
206	Prevalence of use and appropriateness of antidepressants prescription in acutely hospitalized elderly patients. <i>European Journal of Internal Medicine</i> , 2019 , 68, e7-e11	3.9	1
205	A functional variant of the dimethylarginine dimethylaminohydrolase-2 gene is associated with myocardial infarction in type 2 diabetic patients. <i>Cardiovascular Diabetology</i> , 2019 , 18, 102	8.7	5

204	Ketogenic Diet-Induced Weight Loss is Associated with an Increase in Vitamin D Levels in Obese Adults. <i>Molecules</i> , 2019 , 24,	4.8	20
203	Are Circulating Mg Levels Associated with Glucose Tolerance Profiles and Incident Type 2 Diabetes?. <i>Nutrients</i> , 2019 , 11,	6.7	9
202	Immunity, Inflammation and Heart Failure: Their Role on Cardiac Function and Iron Status. <i>Frontiers in Immunology</i> , 2019 , 10, 2315	8.4	8
201	Double Infection in a Patient with Psoriatic Arthritis Under TNF-alpha Blockers Therapy: A Case Report. <i>Current Drug Safety</i> , 2019 , 14, 147-150	1.4	0
200	Mortality rate and risk factors for gastrointestinal bleeding in elderly patients. <i>European Journal of Internal Medicine</i> , 2019 , 61, 54-61	3.9	34
199	Sex-Differences in the Pattern of Comorbidities, Functional Independence, and Mortality in Elderly Inpatients: Evidence from the RePoSI Register. <i>Journal of Clinical Medicine</i> , 2019 , 8,	5.1	13
198	Low Rate of Intrahospital Deep Venous Thrombosis in Acutely Ill Medical Patients: Results From the AURELIO Study. <i>Mayo Clinic Proceedings</i> , 2019 , 94, 37-43	6.4	8
197	One-hour post-load hyperglycemia combined with HbA1c identifies individuals with higher risk of cardiovascular diseases: Cross-sectional data from the CATAMERI study. <i>Diabetes/Metabolism Research and Reviews</i> , 2019 , 35, e3096	7.5	16
196	Short- vs long-course antibiotic therapy for pneumonia: a comparison of systematic reviews and guidelines for the SIMI Choosing Wisely Campaign. <i>Internal and Emergency Medicine</i> , 2019 , 14, 377-394	3.7	3
195	Cow Milk Consumption and Health: A Health Professional Guide. <i>Journal of the American College of Nutrition</i> , 2019 , 38, 197-208	3.5	40
194	Metabolic and vascular effects of silybin in hypertensive patients with high 1-h post-load plasma glucose. <i>Internal and Emergency Medicine</i> , 2019 , 14, 77-84	3.7	3
193	Position paper of the Italian Society of Internal Medicine (SIMI) on prophylaxis and treatment of venous thromboembolism in patients with cancer. <i>Internal and Emergency Medicine</i> , 2019 , 14, 21-38	3.7	8
192	Use of oral anticoagulant drugs in older patients with atrial fibrillation in internal medicine wards. <i>European Journal of Internal Medicine</i> , 2018 , 52, e12-e14	3.9	6
191	Major adverse cardiovascular events in non-valvular atrial fibrillation with chronic obstructive pulmonary disease: the ARAPACIS study. <i>Internal and Emergency Medicine</i> , 2018 , 13, 651-660	3.7	9
190	Data on incidence of bleeding in patients with atrial fibrillation and advanced liver fibrosis on treatment with vitamin K or non-vitamin K antagonist oral anticoagulants. <i>Data in Brief</i> , 2018 , 17, 830-836 ^{1,2}		5
189	Italian intersociety consensus on prevention, diagnosis, and treatment of delirium in hospitalized older persons. <i>Internal and Emergency Medicine</i> , 2018 , 13, 113-121	3.7	15
188	Uric Acid Impairs Insulin Signaling by Promoting Enpp1 Binding to Insulin Receptor in Human Umbilical Vein Endothelial Cells. <i>Frontiers in Endocrinology</i> , 2018 , 9, 98	5.7	21
187	A novel flash glucose monitoring system in patients with diabetes mellitus type 1 and 2: which are the pieces of evidence for payer decision-makers?. <i>Internal and Emergency Medicine</i> , 2018 , 13, 947-949	3.7	

186	Prevalence of Delirium in a Population of Elderly Outpatients with Dementia: A Retrospective Study. <i>Journal of Alzheimer's Disease</i> , 2018 , 61, 251-257	4.3	5
185	Competitive interaction between chronic obstructive pulmonary disease and CHADS-VASc score in predicting incident atrial fibrillation. <i>International Journal of Cardiology</i> , 2018 , 255, 74-79	3.2	4
184	Management of hepatitis C virus infection in patients with chronic kidney disease: position statement of the joint committee of Italian association for the study of the liver (AISF), Italian society of internal medicine (SIMI), Italian society of infectious and tropical disease (SIMIT) and Italian society of nephrology (SIN). <i>Digestive and Liver Disease</i> , 2018 , 50, 1133-1152	3.3	2
183	Serum IgG2 levels are specifically associated with whole-body insulin-mediated glucose disposal in non-diabetic offspring of type 2 diabetic individuals: a cross-sectional study. <i>Scientific Reports</i> , 2018 , 8, 13616	4.9	5
182	One-Hour Postload Hyperglycemia: Implications for Prediction and Prevention of Type 2 Diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018 , 103, 3131-3143	5.6	23
181	The polymorphism rs35767 at IGF1 locus is associated with serum urate levels. <i>Scientific Reports</i> , 2018 , 8, 12255	4.9	1
180	Incidence of bleeding in patients with atrial fibrillation and advanced liver fibrosis on treatment with vitamin K or non-vitamin K antagonist oral anticoagulants. <i>International Journal of Cardiology</i> , 2018 , 264, 58-63	3.2	50
179	Higher serum levels of uric acid are associated with a reduced insulin clearance in non-diabetic individuals. <i>Acta Diabetologica</i> , 2018 , 55, 835-842	3.9	17
178	Nutraceuticals and functional foods for the control of plasma cholesterol levels. An intersociety position paper. <i>Pharmacological Research</i> , 2018 , 134, 51-60	10.2	79
177	Carotid plaque detection improves the predictive value of CHADS-VASc score in patients with non-valvular atrial fibrillation: The ARAPACIS Study. <i>International Journal of Cardiology</i> , 2017 , 231, 143-149	3.2	16
176	Uric Acid Is Associated With Inflammatory Biomarkers and Induces Inflammation Via Activating the NF- κ B Signaling Pathway in HepG2 Cells. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2017 , 37, 1241-1249	9.4	88
175	Risk reclassification ability of uric acid for cardiovascular outcomes in essential hypertension. <i>International Journal of Cardiology</i> , 2017 , 243, 473-478	3.2	20
174	Elevated 1-h post-challenge plasma glucose levels in subjects with normal glucose tolerance or impaired glucose tolerance are associated with whole blood viscosity. <i>Acta Diabetologica</i> , 2017 , 54, 775-784	3.9	12
173	Severe reduction of blood lysosomal acid lipase activity in cryptogenic cirrhosis: A nationwide multicentre cohort study. <i>Atherosclerosis</i> , 2017 , 262, 179-184	3.1	14
172	Duodenal Sodium/Glucose Cotransporter 1 Expression Under Fasting Conditions Is Associated With Postload Hyperglycemia. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017 , 102, 3979-3989	5.6	20
171	Association between hemoglobin glycation index and hepatic steatosis in non-diabetic individuals. <i>Diabetes Research and Clinical Practice</i> , 2017 , 134, 53-61	7.4	17
170	Growth Hormone Deficiency Is Associated with Worse Cardiac Function, Physical Performance, and Outcome in Chronic Heart Failure: Insights from the T.O.S.CA. GHD Study. <i>PLoS ONE</i> , 2017 , 12, e0170058	3.7	37
169	Association between hemoglobin glycation index with insulin resistance and carotid atherosclerosis in non-diabetic individuals. <i>PLoS ONE</i> , 2017 , 12, e0175547	3.7	28

168	Plasma kisspeptin levels are associated with insulin secretion in nondiabetic individuals. <i>PLoS ONE</i> , 2017 , 12, e0179834	3.7	21
167	Elevated 1-h post-load plasma glucose levels in subjects with normal glucose tolerance are associated with a pro-atherogenic lipid profile. <i>Atherosclerosis</i> , 2017 , 256, 15-20	3.1	14
166	A polymorphism at IGF1 locus is associated with anemia. <i>Oncotarget</i> , 2017 , 8, 32398-32406	3.3	6
165	Elevated hemoglobin glycation index identify non-diabetic individuals at increased risk of kidney dysfunction. <i>Oncotarget</i> , 2017 , 8, 79576-79586	3.3	10
164	Sympathovagal balance and 1-h postload plasma glucose in normoglycose tolerant hypertensive patients. <i>Acta Diabetologica</i> , 2016 , 53, 41-7	3.9	3
163	Aspirin and renal insufficiency progression in patients with atrial fibrillation and chronic kidney disease. <i>International Journal of Cardiology</i> , 2016 , 223, 619-624	3.2	11
162	One-Hour Postload Hyperglycemia Confers Higher Risk of Hepatic Steatosis to HbA1c-Defined Prediabetic Subjects. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2016 , 101, 4030-4038	5.6	21
161	Clinical Severity, Age, and Sex Overcome Cardiometabolic Morbidities but Not Stroke as Predictors of Mortality in Elderly Inpatients: Data From the REgistro POLiterapie Societ�Italiana di Medicina Interna Registry. <i>Journal of the American Geriatrics Society</i> , 2016 , 64, 1737-9	5.6	2
160	The Italian Society of Internal Medicine choosing wisely campaign. <i>Internal and Emergency Medicine</i> , 2016 , 11, 1125-1130	3.7	21
159	Response to the Letter: Comment to the letter by Marathe CS, Rayne CK, Jones KL, Horowitz M. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2016 , 101, L35	5.6	
158	Ankle-Brachial Index and cardiovascular events in atrial fibrillation. The ARAPACIS Study. <i>Thrombosis and Haemostasis</i> , 2016 , 115, 856-63	7	26
157	Multiplicative effect of serum phosphorus levels and insulin resistance on hypertensive vascular stiffness. <i>Thrombosis and Haemostasis</i> , 2016 , 115, 227-9	7	7
156	Additive Effect of Non-Alcoholic Fatty Liver Disease on Metabolic Syndrome-Related Endothelial Dysfunction in Hypertensive Patients. <i>International Journal of Molecular Sciences</i> , 2016 , 17, 456	6.3	18
155	Cardiovascular Events in Rheumatoid Arthritis-Time to See Beyond Articular Involvement in "Real World" Clinical Practice: Comment on the Article by Mackey et al. <i>Arthritis and Rheumatology</i> , 2016 , 68, 554-5	9.5	
154	Portal vein thrombosis relevance on liver cirrhosis: Italian Venous Thrombotic Events Registry. <i>Internal and Emergency Medicine</i> , 2016 , 11, 1059-1066	3.7	62
153	Serum phosphorus levels are associated with endothelial dysfunction in hypertensive patients. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2016 , 26, 683-8	4.5	6
152	Renal function is impaired in normotensive chronic HCV patients: role of insulin resistance. <i>Internal and Emergency Medicine</i> , 2016 , 11, 553-9	3.7	4
151	Multiple hormone deficiency syndrome in heart failure with preserved ejection fraction. <i>International Journal of Cardiology</i> , 2016 , 225, 1-3	3.2	32

150	One-hour post-load hyperglycemia combined with HbA1c identifies pre-diabetic individuals with a higher cardio-metabolic risk burden. <i>Atherosclerosis</i> , 2016 , 253, 61-69	3.1	27
149	Relationship between carotid intima-media thickness and non valvular atrial fibrillation type. <i>Atherosclerosis</i> , 2015 , 238, 350-5	3.1	23
148	Reducing the risk of hospital admission: a call to action from the Italian Society of Internal Medicine. <i>European Journal of Internal Medicine</i> , 2015 , 26, 476-7	3.9	2
147	Lights and shadows in the management of old and new oral anticoagulants in the real world of atrial fibrillation by Italian internists. A survey from the Atrial Fibrillation Registry for Ankle-Brachial Index Prevalence Assessment-Collaborative Italian Study. <i>European Journal of Internal Medicine</i> , 2015 , 26, e31-3	3.9	2
146	Uric acid is an independent predictor of cardiovascular events in post-menopausal women. <i>International Journal of Cardiology</i> , 2015 , 197, 271-5	3.2	19
145	Frequency of Left Ventricular Hypertrophy in Non-Valvular Atrial Fibrillation. <i>American Journal of Cardiology</i> , 2015 , 116, 877-82	3	21
144	Low dose of acetylsalicylic acid and oxidative stress-mediated endothelial dysfunction in diabetes: a short-term evaluation. <i>Acta Diabetologica</i> , 2015 , 52, 249-56	3.9	9
143	Rheumatoid arthritis affects left ventricular mass: Systematic review and meta-analysis. <i>European Journal of Internal Medicine</i> , 2015 , 26, 259-67	3.9	16
142	One-Hour Postload Hyperglycemia Is a Stronger Predictor of Type 2 Diabetes Than Impaired Fasting Glucose. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015 , 100, 3744-51	5.6	78
141	Insulin-resistance HCV infection-related affects vascular stiffness in normotensives. <i>Atherosclerosis</i> , 2015 , 238, 108-12	3.1	16
140	Characterization of left ventricular mass in individuals at risk for type 2 diabetes identified by HbA1c levels according to the American Diabetes Association criteria. <i>International Journal of Cardiology</i> , 2015 , 179, 211-3	3.2	6
139	Atrial fibrillation is not associated with increased risk of venous thromboembolism: Results from ARAPACIS study. <i>Thrombosis and Haemostasis</i> , 2015 , 114, 655	7	1
138	SRT1720 counteracts glucosamine-induced endoplasmic reticulum stress and endothelial dysfunction. <i>Cardiovascular Research</i> , 2015 , 107, 295-306	9.9	19
137	CHADS2 and CHA2DS2-VASc scores are independently associated with incident atrial fibrillation: the Catanzaro Atrial Fibrillation Project. <i>Internal and Emergency Medicine</i> , 2015 , 10, 815-21	3.7	21
136	Relationship between low Ankle-Brachial Index and rapid renal function decline in patients with atrial fibrillation: a prospective multicentre cohort study. <i>BMJ Open</i> , 2015 , 5, e008026	3	17
135	Serum alkaline phosphatase negatively affects endothelium-dependent vasodilation in naïve hypertensive patients. <i>Hypertension</i> , 2015 , 66, 874-80	8.5	20
134	Vitamin D and 1-hour post-load plasma glucose in hypertensive patients. <i>Cardiovascular Diabetology</i> , 2014 , 13, 48	8.7	14
133	Differences in insulin clearance between metabolically healthy and unhealthy obese subjects. <i>Acta Diabetologica</i> , 2014 , 51, 257-61	3.9	37

132	Elevated 1-h post-load plasma glucose levels in subjects with normal glucose tolerance are associated with unfavorable inflammatory profile. <i>Acta Diabetologica</i> , 2014 , 51, 927-32	3.9	26
131	Dietary patterns and 1-h post-load glucose in essential hypertension. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2014 , 24, 547-53	4.5	4
130	A polymorphism at IGF1 locus is associated with carotid intima media thickness and endothelium-dependent vasodilatation. <i>Atherosclerosis</i> , 2014 , 232, 25-30	3.1	20
129	Chronic HCV infection increases cardiac left ventricular mass index in normotensive patients. <i>Journal of Hepatology</i> , 2014 , 61, 755-60	13.4	15
128	Association between noninvasive fibrosis markers and cardio-vascular organ damage among adults with hepatic steatosis. <i>PLoS ONE</i> , 2014 , 9, e104941	3.7	23
127	Elevated 1 h postload plasma glucose levels identify adults with normal glucose tolerance but increased risk of non-alcoholic fatty liver disease. <i>BMJ Open Diabetes Research and Care</i> , 2014 , 2, e000145	16.5	28
126	Genome-wide association study identifies CAMKID variants involved in blood pressure response to losartan: the SOPHIA study. <i>Pharmacogenomics</i> , 2014 , 15, 1643-52	2.6	24
125	Clinical risk factors and subclinical target organ damage as predictors of new-onset of atrial fibrillation: the Catanzaro atrial fibrillation project. <i>International Journal of Cardiology</i> , 2014 , 177, 666-8	3.2	2
124	Medication prescription and adherence disparities in non valvular atrial fibrillation patients: an Italian portrait from the ARAPACIS study. <i>Internal and Emergency Medicine</i> , 2014 , 9, 861-70	3.7	21
123	Renal disease and left atrial remodeling predict atrial fibrillation in patients with cardiovascular risk factors. <i>International Journal of Cardiology</i> , 2014 , 175, 90-5	3.2	21
122	Reply: ankle-brachial index in patients with nonvalvular atrial fibrillation. <i>Journal of the American College of Cardiology</i> , 2014 , 63, 1457-8	15.1	1
121	3PUTR OLR1/LOX-1 gene polymorphism and endothelial dysfunction: molecular and vascular data in never-treated hypertensive patients. <i>Internal and Emergency Medicine</i> , 2014 , 9, 273-81	3.7	2
120	Association between noninvasive fibrosis markers and chronic kidney disease among adults with nonalcoholic fatty liver disease. <i>PLoS ONE</i> , 2014 , 9, e88569	3.7	33
119	Prevalence of peripheral artery disease by abnormal ankle-brachial index in atrial fibrillation: implications for risk and therapy. <i>Journal of the American College of Cardiology</i> , 2013 , 62, 2255-6	15.1	44
118	Heart rate affects endothelial function in essential hypertension. <i>Internal and Emergency Medicine</i> , 2013 , 8, 211-9	3.7	5
117	Endothelial dysfunction predicts regression of hypertensive cardiac mass. <i>International Journal of Cardiology</i> , 2013 , 167, 1188-92	3.2	12
116	Creatinine and insulin predict cardiac mass in drug-naïve hypertensive patients. <i>International Journal of Cardiology</i> , 2013 , 167, 519-24	3.2	6
115	Phenotypic characterization of normotolerant hypertensive patients. <i>International Journal of Cardiology</i> , 2013 , 165, 322-6	3.2	7

114	Interaction between uric acid and endothelial dysfunction predicts new onset of diabetes in hypertensive patients. <i>International Journal of Cardiology</i> , 2013 , 167, 232-6	3.2	31
113	Insulin clearance is associated with carotid artery intima-media thickness. <i>Atherosclerosis</i> , 2013 , 229, 453-8	3.1	7
112	A functional variant of the dimethylarginine dimethylaminohydrolase-2 gene is associated with chronic kidney disease. <i>Atherosclerosis</i> , 2013 , 231, 141-4	3.1	6
111	Angiotensin (1-7) counteracts the negative effect of angiotensin II on insulin signalling in HUVECs. <i>Cardiovascular Research</i> , 2013 , 99, 129-36	9.9	29
110	Insulin-like growth factor-I, inflammatory proteins, and fibrosis in subjects with nonalcoholic fatty liver disease. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013 , 98, E304-8	5.6	48
109	Association of insulin resistance indexes to carotid intima-media thickness. <i>PLoS ONE</i> , 2013 , 8, e53968	3.7	11
108	Decreased insulin clearance in individuals with elevated 1-h post-load plasma glucose levels. <i>PLoS ONE</i> , 2013 , 8, e77440	3.7	21
107	A fasting insulin-raising allele at IGF1 locus is associated with circulating levels of IGF-1 and insulin sensitivity. <i>PLoS ONE</i> , 2013 , 8, e85483	3.7	26
106	Oxidative stress impairs endothelial function in nondipper hypertensive patients. <i>Cardiovascular Therapeutics</i> , 2012 , 30, 85-92	3.3	21
105	Cardiometabolic risk profiles and carotid atherosclerosis in individuals with prediabetes identified by fasting glucose, postchallenge glucose, and hemoglobin A1c criteria. <i>Diabetes Care</i> , 2012 , 35, 1144-9	14.6	67
104	One-hour post-load plasma glucose and IGF-1 in hypertensive patients. <i>European Journal of Clinical Investigation</i> , 2012 , 42, 1325-31	4.6	9
103	A functional variant of the dimethylarginine dimethylaminohydrolase-2 gene is associated with insulin sensitivity. <i>PLoS ONE</i> , 2012 , 7, e36224	3.7	16
102	Association between one-hour post-load plasma glucose levels and vascular stiffness in essential hypertension. <i>PLoS ONE</i> , 2012 , 7, e44470	3.7	50
101	Asymmetric dimethylarginine plasma levels and endothelial function in newly diagnosed type 2 diabetic patients. <i>International Journal of Molecular Sciences</i> , 2012 , 13, 13804-15	6.3	24
100	Serum uric acid and 1-h postload glucose in essential hypertension. <i>Diabetes Care</i> , 2012 , 35, 153-7	14.6	33
99	Insulin sensitivity, β cell function, and incretin effect in individuals with elevated 1-hour postload plasma glucose levels. <i>Diabetes Care</i> , 2012 , 35, 868-72	14.6	58
98	Angiotensin II type 1 receptor, but no type 2 receptor, interferes with the insulin-induced nitric oxide production in HUVECs. <i>Atherosclerosis</i> , 2011 , 219, 463-7	3.1	7
97	Low insulin-like growth factor-1 levels are associated with anaemia in adult non-diabetic subjects. <i>Thrombosis and Haemostasis</i> , 2011 , 105, 365-70	7	20

96	Cardiovascular and noncardiovascular comorbidities in patients with chronic heart failure. <i>Journal of Cardiovascular Medicine</i> , 2011 , 12, 76-84	1.9	45
95	Usefulness of hemoglobin A1c as a criterion to define the metabolic syndrome in a cohort of italian nondiabetic white subjects. <i>American Journal of Cardiology</i> , 2011 , 107, 1650-5	3	22
94	Nonalcoholic fatty liver disease is associated with low circulating levels of insulin-like growth factor-I. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2011 , 96, E1640-4	5.6	70
93	One-hour postload plasma glucose levels and diastolic function in hypertensive patients. <i>Diabetes Care</i> , 2011 , 34, 2291-6	14.6	37
92	The TRIB3 R84 variant is associated with increased carotid intima-media thickness in vivo and with enhanced MAPK signalling in human endothelial cells. <i>Cardiovascular Research</i> , 2011 , 89, 184-92	9.9	23
91	One-hour postload plasma glucose levels and left ventricular mass in hypertensive patients. <i>Diabetes Care</i> , 2011 , 34, 1406-11	14.6	68
90	Association between hemoglobin level and endothelial function in uncomplicated, untreated hypertensive patients. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2011 , 6, 648-55	6.9	11
89	One-hour postload plasma glucose levels are associated with kidney dysfunction. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2010 , 5, 1922-7	6.9	61
88	Association between carotid atherosclerosis and metabolic syndrome: results from the ISMIR study. <i>Angiology</i> , 2010 , 61, 443-8	2.1	13
87	Positive association between plasma IGF1 and high-density lipoprotein cholesterol levels in adult nondiabetic subjects. <i>European Journal of Endocrinology</i> , 2010 , 163, 75-80	6.5	23
86	Endothelial dysfunction and subsequent decline in glomerular filtration rate in hypertensive patients. <i>Circulation</i> , 2010 , 122, 379-84	16.7	85
85	Endothelial dysfunction, ADMA and insulin resistance in essential hypertension. <i>International Journal of Cardiology</i> , 2010 , 142, 236-41	3.2	60
84	Carotid intima-media thickness in asymptomatic patients with arterial hypertension without clinical cardiovascular disease: relation with left ventricular geometry and mass and coexisting risk factors. <i>Angiology</i> , 2009 , 60, 705-13	2.1	15
83	ENPP1 Q121 variant, increased pulse pressure and reduced insulin signaling, and nitric oxide synthase activity in endothelial cells. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2009 , 29, 1678-83	9.4	22
82	Renal function predicts cardiovascular outcomes in southern Italian postmenopausal women. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2009 , 16, 481-6		12
81	The TRIB3 Q84R polymorphism and risk of early-onset type 2 diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2009 , 94, 190-6	5.6	49
80	Relation of carotid intima-media thickness and aortic valve sclerosis (from the ISMIR study ["Ispessimento Medio Intimale e Rischio Cardiovascolare"]) of the Italian Society of Cardiovascular Echography). <i>American Journal of Cardiology</i> , 2009 , 103, 1556-61	3	12
79	Differences in cardiovascular risk profile based on relationship between post-load plasma glucose and fasting plasma levels. <i>Diabetes/Metabolism Research and Reviews</i> , 2009 , 25, 351-6	7.5	6

78	Insulin-like growth factor-1 and glomerular filtration rate in hypertensive patients. <i>Journal of Hypertension</i> , 2009 , 27, 613-7	1.9	17
77	Insulin secretion in metabolically obese, but normal weight, and in metabolically healthy but obese individuals. <i>Obesity</i> , 2008 , 16, 1881-6	8	110
76	Kidney function and risk factors for left ventricular hypertrophy in untreated uncomplicated essential hypertension. <i>American Journal of Kidney Diseases</i> , 2008 , 52, 74-84	7.4	25
75	Large brachial and common carotid artery diameter in postmenopausal women with carotid atherosclerosis. <i>Atherosclerosis</i> , 2008 , 196, 443-448	3.1	17
74	The CCR2 promoter polymorphism T-960A, but not the serum MCP-1 level, is associated with endothelial function in prediabetic individuals. <i>Atherosclerosis</i> , 2008 , 198, 338-46	3.1	7
73	Reciprocal association of plasma IGF-1 and interleukin-6 levels with cardiometabolic risk factors in nondiabetic subjects. <i>Diabetes Care</i> , 2008 , 31, 1886-8	14.6	43
72	Endothelial dysfunction and C-reactive protein are risk factors for diabetes in essential hypertension. <i>Diabetes</i> , 2008 , 57, 167-71	0.9	67
71	Low-plasma insulin-like growth factor-I levels are associated with impaired endothelium-dependent vasodilatation in a cohort of untreated, hypertensive Caucasian subjects. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008 , 93, 2806-10	5.6	33
70	Exercise prescription for the prevention and treatment of cardiovascular diseases: part II. <i>Journal of Cardiovascular Medicine</i> , 2008 , 9, 641-52	1.9	9
69	Exercise prescription for the prevention and treatment of cardiovascular diseases: part I. <i>Journal of Cardiovascular Medicine</i> , 2008 , 9, 529-44	1.9	21
68	Plasma interleukin-6 levels are increased in subjects with impaired glucose tolerance but not in those with impaired fasting glucose in a cohort of Italian Caucasians. <i>Diabetes/Metabolism Research and Reviews</i> , 2007 , 23, 141-5	7.5	27
67	Metabolic and cardiovascular risk factors in subjects with impaired fasting glucose: the 100 versus 110 mg/dL threshold. <i>Diabetes/Metabolism Research and Reviews</i> , 2007 , 23, 547-50	7.5	21
66	Glutamine to arginine substitution at amino acid 84 of mammalian tribbles homolog TRIB3 and CKD in whites with type 2 diabetes. <i>American Journal of Kidney Diseases</i> , 2007 , 50, 688-9	7.4	14
65	Interleukin-6 impairs the insulin signaling pathway, promoting production of nitric oxide in human umbilical vein endothelial cells. <i>Molecular and Cellular Biology</i> , 2007 , 27, 2372-83	4.8	86
64	Metabolically healthy but obese women have an intermediate cardiovascular risk profile between healthy nonobese women and obese insulin-resistant women. <i>Diabetes Care</i> , 2007 , 30, 2145-7	14.6	122
63	Successful radiofrequency ablation determines atrio-ventricular remodelling and improves systo-diastolic function at tissue Doppler-imaging. <i>European Journal of Cardio-thoracic Surgery</i> , 2007 , 31, 414-21; discussion 421-2	3	7
62	Gly460Trp alpha-adducin gene polymorphism and endothelial function in untreated hypertensive patients. <i>Journal of Hypertension</i> , 2007 , 25, 2234-9	1.9	8
61	Endogenous testosterone and endothelial function in postmenopausal women. <i>Coronary Artery Disease</i> , 2007 , 18, 9-13	1.4	56

60	Effects of growth hormone and insulin-like growth factor-1 on cardiac hypertrophy of hypertensive patients. <i>Journal of Hypertension</i> , 2007 , 25, 471-7	1.9	23
59	Role of PC-1 and ACE genes on insulin resistance and cardiac mass in never-treated hypertensive patients. Suggestive evidence for a digenic additive modulation. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2007 , 17, 181-7	4.5	12
58	Prognostic value of serial electrocardiographic voltage and repolarization changes in essential hypertension: the HEART Survey study. <i>American Journal of Hypertension</i> , 2007 , 20, 997-1004	2.3	29
57	Carotid atherosclerosis associated to metabolic syndrome but not BMI in healthy menopausal women. <i>Diabetes Research and Clinical Practice</i> , 2007 , 76, 378-82	7.4	22
56	Serum magnesium and lipids: More clarity is needed. <i>Atherosclerosis</i> , 2007 , 192, 233-4; author reply 231-3	3.1	6
55	Effect of interaction between left ventricular dysfunction and endothelial function in hypertension. <i>Current Hypertension Reports</i> , 2006 , 8, 212-8	4.7	9
54	Plasma interleukin-6 levels are independently associated with insulin secretion in a cohort of Italian-Caucasian nondiabetic subjects. <i>Diabetes</i> , 2006 , 55, 2021-4	0.9	44
53	The E23K variant of KCNJ11 encoding the pancreatic beta-cell adenosine 5P triphosphate-sensitive potassium channel subunit Kir6.2 is associated with an increased risk of secondary failure to sulfonylurea in patients with type 2 diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2006 , 91, 2334-9	5.6	139
52	Uric acid and endothelial dysfunction in essential hypertension. <i>Journal of the American Society of Nephrology: JASN</i> , 2006 , 17, 1466-71	12.7	180
51	Variants of the interleukin-10 promoter gene are associated with obesity and insulin resistance but not type 2 diabetes in caucasian italian subjects. <i>Diabetes</i> , 2006 , 55, 1529-33	0.9	73
50	Inflammation as a mediator of the link between mild to moderate renal insufficiency and endothelial dysfunction in essential hypertension. <i>Journal of the American Society of Nephrology: JASN</i> , 2006 , 17, S64-8	12.7	32
49	Serum ionized magnesium levels in relation to metabolic syndrome in type 2 diabetic patients. <i>Journal of the American College of Nutrition</i> , 2006 , 25, 210-5	3.5	74
48	The prognostic value of big endothelin-1 in more than 2,300 patients with heart failure enrolled in the Valsartan Heart Failure Trial (Val-HeFT). <i>Journal of Cardiac Failure</i> , 2006 , 12, 375-80	3.3	54
47	Betamyosin mutations and phenotypic heterogeneity in hypertrophic cardiomyopathy. <i>International Journal of Cardiology</i> , 2006 , 110, 119-21	3.2	2
46	Interaction between vascular dysfunction and cardiac mass increases the risk of cardiovascular outcomes in essential hypertension. <i>European Heart Journal</i> , 2005 , 26, 921-7	9.5	39
45	Asymmetric dimethylarginine, L-arginine, and endothelial dysfunction in essential hypertension. <i>Journal of the American College of Cardiology</i> , 2005 , 46, 518-23	15.1	214
44	Association of LDL cholesterol with carotid atherosclerosis in menopausal women affected by the metabolic syndrome. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2005 , 15, 368-72	4.5	21
43	Plasma concentration of IGF-I is independently associated with insulin sensitivity in subjects with different degrees of glucose tolerance. <i>Diabetes Care</i> , 2005 , 28, 120-5	14.6	135

42	Impaired endothelial function in never-treated hypertensive subjects carrying the Arg972 polymorphism in the insulin receptor substrate-1 gene. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2004 , 89, 3606-9	5.6	34
41	The -866A/A genotype in the promoter of the human uncoupling protein 2 gene is associated with insulin resistance and increased risk of type 2 diabetes. <i>Diabetes</i> , 2004 , 53, 1905-10	0.9	94
40	The Arg972 variant in insulin receptor substrate-1 is associated with an increased risk of secondary failure to sulfonylurea in patients with type 2 diabetes. <i>Diabetes Care</i> , 2004 , 27, 1394-8	14.6	62
39	Angiotensin II impairs the insulin signaling pathway promoting production of nitric oxide by inducing phosphorylation of insulin receptor substrate-1 on Ser312 and Ser616 in human umbilical vein endothelial cells. <i>Circulation Research</i> , 2004 , 94, 1211-8	15.7	167
38	Endothelial dysfunction and mild renal insufficiency in essential hypertension. <i>Circulation</i> , 2004 , 110, 821-5	16.7	84
37	Trends in prescribing antihypertensive drugs in the Gruppo Italiano di Farmacovigilanza nell'Anziano 1988-1997. <i>European Journal of Clinical Pharmacology</i> , 2003 , 58, 851-2	2.8	
36	Relation of fasting insulin related to insertion/deletion polymorphism of angiotensin-converting enzyme-gene and cardiac mass in never-treated patients with systemic hypertension. <i>American Journal of Cardiology</i> , 2003 , 92, 1234-7	3	15
35	Pulse pressure and endothelial dysfunction in never-treated hypertensive patients. <i>Journal of the American College of Cardiology</i> , 2003 , 41, 1753-8	15.1	84
34	Comparative effect of lercanidipine, felodipine, and nifedipine GITS on blood pressure and heart rate in patients with mild to moderate arterial hypertension: the Lercanidipine in Adults (LEAD) Study. <i>Journal of Clinical Hypertension</i> , 2003 , 5, 249-53	2.3	29
33	Renin-Angiotensin system and endothelial function. <i>International Congress Series</i> , 2003 , 1253, 205-208		
32	Seasonal hypertension: a clue to explain the high prevalence of unrecognized hypertension in the elderly? Gruppo Italiano di Farmacovigilanza nell'Anziano (GIFA). <i>Aging Clinical and Experimental Research</i> , 2003 , 15, 296-300	4.8	4
31	Weight loss in combination with physical activity improves endothelial dysfunction in human obesity. <i>Diabetes Care</i> , 2003 , 26, 1673-8	14.6	121
30	Protection of human endothelial cells from oxidative stress: role of Ras-ERK1/2 signaling. <i>Circulation</i> , 2002 , 105, 968-74	16.7	84
29	Inhibition of neutrophil apoptosis after coronary bypass operation with cardiopulmonary bypass. <i>Annals of Thoracic Surgery</i> , 2002 , 73, 123-9; discussion 129-30	2.7	54
28	Effects of leptin on platelet function in obese patients. <i>JAMA - Journal of the American Medical Association</i> , 2002 , 288, 313-4; author reply 314	27.4	4
27	Post-prandial hypertriglyceridemia and endothelium-dependent vasorelaxation. <i>European Journal of Clinical Investigation</i> , 2001 , 31, 1013-4	4.6	3
26	Relationship between angiotensin-converting enzyme gene polymorphism and insulin resistance in never-treated hypertensive patients. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2001 , 86, 172-8	5.6	31
25	Obesity and body fat distribution induce endothelial dysfunction by oxidative stress: protective effect of vitamin C. <i>Diabetes</i> , 2001 , 50, 159-65	0.9	284

24	Comparison of endothelial function evaluated by strain gauge plethysmography and brachial artery ultrasound. <i>Atherosclerosis</i> , 2001 , 158, 53-9	3.1	35
23	Prognostic significance of endothelial dysfunction in hypertensive patients. <i>Circulation</i> , 2001 , 104, 191-6	16.7	901
22	Continuous relation between left ventricular mass and cardiovascular risk in essential hypertension. <i>Hypertension</i> , 2000 , 35, 580-6	8.5	396
21	Effects of atorvastatin and vitamin C on endothelial function of hypercholesterolemic patients. <i>Atherosclerosis</i> , 2000 , 152, 511-8	3.1	66
20	Hypertensive left ventricular remodeling and ACE-gene polymorphism. <i>Cardiovascular Research</i> , 1999 , 43, 192-9	9.9	21
19	Relationship between left ventricular mass and endothelium-dependent vasodilation in never-treated hypertensive patients. <i>Circulation</i> , 1999 , 99, 1991-6	16.7	86
18	ACE-gene polymorphism and endothelial dysfunction in normal humans. <i>Hypertension</i> , 1999 , 34, e20-1	8.5	1
17	Nitric oxide modulation of neutrophil-endothelium interaction: difference between arterial and venous coronary bypass grafts. <i>Journal of the American College of Cardiology</i> , 1998 , 31, 823-6	15.1	42
16	Prolongation of the QT interval and the sudden infant death syndrome. <i>New England Journal of Medicine</i> , 1998 , 338, 1709-14	59.2	580
15	Neutrophil-endothelial cells modulation in diabetic patients undergoing coronary artery bypass grafting. <i>European Journal of Cardio-thoracic Surgery</i> , 1998 , 14, 373-9	3	11
14	Side-Branch of Internal Mammary Artery Bypass Graft Causing Coronary Steal. <i>Asian Cardiovascular and Thoracic Annals</i> , 1998 , 6, 219-220	0.6	4
13	Angiotensin-converting enzyme gene polymorphism is associated with endothelium-dependent vasodilation in never treated hypertensive patients. <i>Hypertension</i> , 1998 , 31, 900-5	8.5	59
12	Prolonged Magnesium Sulfate Infusion in the Treatment of Ventricular Tachycardia in Acquired Long QT Syndrome. <i>Clinical Drug Investigation</i> , 1997 , 13, 229-236	3.2	2
11	Deletion polymorphism of angiotensin-converting enzyme gene and left ventricular hypertrophy in southern Italian patients. <i>Journal of the American College of Cardiology</i> , 1997 , 29, 365-9	15.1	54
10	Elevated circulating levels of von Willebrand factor and D-dimer in patients with heart failure and mechanical prosthesis. <i>Scandinavian Journal of Thoracic and Cardiovascular Surgery</i> , 1996 , 30, 77-81		2
9	Evaluation of antihypertensive effects of once-a-day isradipine and fosinopril: a double-blind crossover study by means of ambulatory blood pressure monitoring. <i>Clinical Cardiology</i> , 1995 , 18, 401-7	3.3	2
8	Prolonged QT interval: a marker of sudden infant death syndrome?. <i>Clinical Cardiology</i> , 1991 , 14, 417-21	3.3	12
7	Heart rate variability and sudden infant death syndrome. <i>PACE - Pacing and Clinical Electrophysiology</i> , 1990 , 13, 2096-9	1.6	15

6	Efficacy of magnesium sulfate in the treatment of torsade de pointes. <i>American Heart Journal</i> , 1986 , 112, 847-9	4.9	68
5	Transvenous catheter ablation of the accessory atrioventricular pathway in the permanent form of junctional reciprocating tachycardia. <i>American Journal of Cardiology</i> , 1985 , 55, 1639-41	3	20
4	Evaluation of noninvasive tests for identifying patients with preexcitation syndrome at risk of rapid ventricular response. <i>American Heart Journal</i> , 1984 , 108, 905-9	4.9	37
3	Transesophageal pacing for prognostic evaluation of preexcitation syndrome and assessment of protective therapy. <i>American Journal of Cardiology</i> , 1983 , 51, 513-8	3	61
2	Programmed electrical stimulation and Amiodarone therapy for the control of persistent junctional tachycardia. <i>Cardiovascular Diseases</i> , 1981 , 8, 232-237		1
1	Automatic "scanning" by radiofrequency in the long-term electrical treatment of arrhythmias. <i>PACE - Pacing and Clinical Electrophysiology</i> , 1979 , 2, 289-96	1.6	25