Richard M Pope

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/8337990/richard-m-pope-publications-by-year.pdf

Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

111	7,387	52	84
papers	citations	h-index	g-index
115	8,013 ext. citations	7.1	5.69
ext. papers		avg, IF	L-index

#	Paper	IF	Citations
111	Critical role of synovial tissue-resident macrophage niche in joint homeostasis and suppression of chronic inflammation. <i>Science Advances</i> , 2021 , 7,	14.3	5
110	Transcriptional Profiling of Synovial Macrophages Using Minimally Invasive Ultrasound-Guided Synovial Biopsies in Rheumatoid Arthritis. <i>Arthritis and Rheumatology</i> , 2018 , 70, 841-854	9.5	27
109	The Role of Macrophages in the Response to TNF Inhibition in Experimental Arthritis. <i>Journal of Immunology</i> , 2018 , 200, 130-138	5.3	15
108	Association of Rheumatoid Factors With Subclinical and Clinical Atherosclerosis in African American Women: The Multiethnic Study of Atherosclerosis. <i>Arthritis Care and Research</i> , 2017 , 69, 166-174	4.7	19
107	Association of Increased F4/80 Macrophages With Suppression of Serum-Transfer Arthritis in Mice With Reduced FLIP in Myeloid Cells. <i>Arthritis and Rheumatology</i> , 2017 , 69, 1762-1771	9.5	9
106	SNAPIN is critical for lysosomal acidification and autophagosome maturation in macrophages. <i>Autophagy</i> , 2017 , 13, 285-301	10.2	17
105	The inflammatory role of phagocyte apoptotic pathways in rheumatic diseases. <i>Nature Reviews Rheumatology</i> , 2016 , 12, 543-58	8.1	34
104	Deletion of calponin 2 in macrophages attenuates the severity of inflammatory arthritis in mice. <i>American Journal of Physiology - Cell Physiology</i> , 2016 , 311, C673-C685	5.4	11
103	CD11c-mediated deletion of Flip promotes autoreactivity and inflammatory arthritis. <i>Nature Communications</i> , 2015 , 6, 7086	17.4	15
102	Fas signaling in macrophages promotes chronicity in K/BxN serum-induced arthritis. <i>Arthritis and Rheumatology</i> , 2014 , 66, 68-77	9.5	12
101	Possible roles of IL-12-family cytokines in rheumatoid arthritis. <i>Nature Reviews Rheumatology</i> , 2013 , 9, 252-6	8.1	68
100	The role of glycoprotein 96 in the persistent inflammation of rheumatoid arthritis. <i>Archives of Biochemistry and Biophysics</i> , 2013 , 530, 1-6	4.1	20
99	TLR2 deletion promotes arthritis through reduction of IL-10. <i>Journal of Leukocyte Biology</i> , 2013 , 93, 757	1- 0 .5	10
98	Toll-like receptor 4 signaling augments transforming growth factor-Iresponses: a novel mechanism for maintaining and amplifying fibrosis in scleroderma. <i>American Journal of Pathology</i> , 2013 , 182, 192-205	5.8	184
97	Antiphospholipid antibodies and sub-clinical atherosclerosis in the Coronary Artery Risk Development in Young Adults (CARDIA) cohort. <i>Inflammation Research</i> , 2013 , 62, 919-27	7.2	17
96	Ligation of TLR7 by rheumatoid arthritis synovial fluid single strand RNA induces transcription of TNFIIn monocytes. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 418-26	2.4	51
95	Requirement of myeloid cell-specific Fas expression for prevention of systemic autoimmunity in mice. <i>Arthritis and Rheumatism</i> , 2012 , 64, 808-20		18

(2009-2012)

94	Glycoprotein 96 perpetuates the persistent inflammation of rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2012 , 64, 3638-48		21
93	TLR5, a novel and unidentified inflammatory mediator in rheumatoid arthritis that correlates with disease activity score and joint TNF-levels. <i>Journal of Immunology</i> , 2012 , 189, 475-83	5.3	59
92	Role of the CCL21 and CCR7 pathways in rheumatoid arthritis angiogenesis. <i>Arthritis and Rheumatism</i> , 2012 , 64, 2471-81		53
91	SNAPIN: an endogenous Toll-like receptor ligand in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 1411-7	2.4	31
90	Portability of an algorithm to identify rheumatoid arthritis in electronic health records. <i>Journal of the American Medical Informatics Association: JAMIA</i> , 2012 , 19, e162-9	8.6	164
89	Anti-CXCL5 therapy ameliorates IL-17-induced arthritis by decreasing joint vascularization. <i>Angiogenesis</i> , 2011 , 14, 443-55	10.6	34
88	Characterization of CCL19 and CCL21 in rheumatoid arthritis. Arthritis and Rheumatism, 2011, 63, 914-2	2	91
87	Characterization of interleukin-7 and interleukin-7 receptor in the pathogenesis of rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2011 , 63, 2884-93		57
86	Activated TLR signaling in atherosclerosis among women with lower Framingham risk score: the multi-ethnic study of atherosclerosis. <i>PLoS ONE</i> , 2011 , 6, e21067	3.7	16
85	Toll-like receptor signaling: a potential link among rheumatoid arthritis, systemic lupus, and atherosclerosis. <i>Journal of Leukocyte Biology</i> , 2010 , 88, 253-62	6.5	54
84	IL-17 contributes to angiogenesis in rheumatoid arthritis. <i>Journal of Immunology</i> , 2010 , 184, 3233-41	5.3	140
83	IL-17-mediated monocyte migration occurs partially through CC chemokine ligand 2/monocyte chemoattractant protein-1 induction. <i>Journal of Immunology</i> , 2010 , 184, 4479-87	5.3	116
82	Prostate secretions from men with chronic pelvic pain syndrome inhibit proinflammatory mediators. <i>Journal of Urology</i> , 2010 , 184, 1536-42	2.5	13
81	Innate immunity and rheumatoid arthritis. Rheumatic Disease Clinics of North America, 2010, 36, 271-96	2.4	82
80	FLIP: a novel regulator of macrophage differentiation and granulocyte homeostasis. <i>Blood</i> , 2010 , 116, 4968-77	2.2	25
79	Bim-Bcl-2 homology 3 mimetic therapy is effective at suppressing inflammatory arthritis through the activation of myeloid cell apoptosis. <i>Arthritis and Rheumatism</i> , 2010 , 62, 441-51		34
78	The synovial lining micromass system: toward rheumatoid arthritis in a dish?. <i>Arthritis and Rheumatism</i> , 2010 , 62, 643-6		14
77	Activation-induced degradation of FLIP(L) is mediated via the phosphatidylinositol 3-kinase/Akt signaling pathway in macrophages. <i>Journal of Biological Chemistry</i> , 2009 , 284, 14513-23	5.4	17

76	IL-17 induces monocyte migration in rheumatoid arthritis. <i>Journal of Immunology</i> , 2009 , 182, 3884-91	5.3	133
75	Heat shock protein 96 is elevated in rheumatoid arthritis and activates macrophages primarily via TLR2 signaling. <i>Journal of Immunology</i> , 2009 , 182, 4965-73	5.3	118
74	The CDK domain of p21 is a suppressor of IL-1beta-mediated inflammation in activated macrophages. <i>European Journal of Immunology</i> , 2009 , 39, 820-5	6.1	52
73	TNFalpha-induced macrophage death via caspase-dependent and independent pathways. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2009 , 14, 320-32	5.4	36
72	The role of toll-like receptors in rheumatoid arthritis. Current Rheumatology Reports, 2009, 11, 357-64	4.9	173
71	Co-stimulatory pathways in the therapy of rheumatoid arthritis 2009 , 27-43		1
70	TH-17 cells in rheumatoid arthritis. Arthritis Research and Therapy, 2008, 10, R93	5.7	162
69	Apigenin, a dietary flavonoid, sensitizes human T cells for activation-induced cell death by inhibiting PKB/Akt and NF-kappaB activation pathway. <i>Immunology Letters</i> , 2008 , 121, 74-83	4.1	38
68	Monocyte chemoattractant protein-1 and macrophage inflammatory protein-1alpha as possible biomarkers for the chronic pelvic pain syndrome. <i>Journal of Urology</i> , 2008 , 179, 1857-61; discussion 186	1 ² -2 ⁵	86
67	Role of H2-calponin in regulating macrophage motility and phagocytosis. <i>Journal of Biological Chemistry</i> , 2008 , 283, 25887-99	5.4	45
66	Inhibition of monocyte chemoattractant protein-1 ameliorates rat adjuvant-induced arthritis. <i>Journal of Immunology</i> , 2008 , 180, 3447-56	5.3	83
65	Transcriptional diversity during monocyte to macrophage differentiation. <i>Immunology Letters</i> , 2008 , 117, 70-80	4.1	59
64	Increased macrophage activation mediated through toll-like receptors in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2007 , 56, 2192-201		150
63	The role of interleukin-1 and the inflammasome in gout: implications for therapy. <i>Arthritis and Rheumatism</i> , 2007 , 56, 3183-8		139
62	ICOS and B7 costimulatory molecule expression identifies activated cellular subsets in rheumatoid arthritis. Cytometry Part A: the Journal of the International Society for Analytical Cytology, 2007, 71, 317-	26 ⁶	21
61	Myeloid cell leukemia-1 as a therapeutic target. <i>Expert Opinion on Therapeutic Targets</i> , 2007 , 11, 363-73	6.4	27
60	The CD95 receptor: apoptosis revisited. <i>Cell</i> , 2007 , 129, 447-50	56.2	327
59	Regulation of Mcl-1 expression in rheumatoid arthritis synovial macrophages. <i>Arthritis and Rheumatism</i> , 2006 , 54, 3174-81		55

(2003-2006)

Inhibition of ADP/ATP exchange in receptor-interacting protein-mediated necrosis. <i>Molecular and Cellular Biology</i> , 2006 , 26, 2215-25	4.8	165
RANTES modulates TLR4-induced cytokine secretion in human peripheral blood monocytes. <i>Journal of Immunology</i> , 2006 , 177, 5077-87	5.3	36
Drug Insight: abatacept for the treatment of rheumatoid arthritis. <i>Nature Clinical Practice Rheumatology</i> , 2006 , 2, 654-60		21
The evolving clinical profile of abatacept (CTLA4-Ig): a novel co-stimulatory modulator for the treatment of rheumatoid arthritis. <i>Arthritis Research</i> , 2005 , 7 Suppl 2, S21-5		66
Amelioration of rat adjuvant-induced arthritis by Met-RANTES. Arthritis and Rheumatism, 2005, 52, 1907	7-19	94
Mcl-1 is essential for the survival of synovial fibroblasts in rheumatoid arthritis. <i>Journal of Immunology</i> , 2005 , 175, 8337-45	5.3	68
NF-kappaB protects macrophages from lipopolysaccharide-induced cell death: the role of caspase 8 and receptor-interacting protein. <i>Journal of Biological Chemistry</i> , 2005 , 280, 41827-34	5.4	73
The role of macrophages in rheumatoid arthritis. Current Pharmaceutical Design, 2005, 11, 569-80	3.3	112
TNF-alpha-induced apoptosis of macrophages following inhibition of NF-kappa B: a central role for disruption of mitochondria. <i>Journal of Immunology</i> , 2004 , 172, 1907-15	5.3	69
Fas ligation on macrophages enhances IL-1R1-Toll-like receptor 4 signaling and promotes chronic inflammation. <i>Nature Immunology</i> , 2004 , 5, 380-7	19.1	113
Retinoblastoma suppression of matrix metalloproteinase 1, but not interleukin-6, through a p38-dependent pathway in rheumatoid arthritis synovial fibroblasts. <i>Arthritis and Rheumatism</i> , 2004 , 50, 78-87		16
NF-kappaB-regulated expression of cellular FLIP protects rheumatoid arthritis synovial fibroblasts from tumor necrosis factor alpha-mediated apoptosis. <i>Arthritis and Rheumatism</i> , 2004 , 50, 3844-55		56
Apoptosis in rheumatoid arthritis: friend or foe. <i>Rheumatic Disease Clinics of North America</i> , 2004 , 30, 603-25, x	2.4	30
Phagocytes: mechanisms of inflammation and tissue destruction. <i>Rheumatic Disease Clinics of North America</i> , 2004 , 30, 19-39, v	2.4	48
Serine phosphorylation of STAT3 is essential for Mcl-1 expression and macrophage survival. <i>Blood</i> , 2003 , 102, 344-52	2.2	119
Rheumatoid arthritis synovial fluid macrophages express decreased tumor necrosis factor-related apoptosis-inducing ligand R2 and increased decoy receptor tumor necrosis factor-related apoptosis-inducing ligand R3. <i>Arthritis and Rheumatism</i> , 2003 , 48, 3096-101		24
The role of apoptosis in rheumatoid arthritis. Current Opinion in Pharmacology, 2003, 3, 317-22	5.1	150
IL-6 and matrix metalloproteinase-1 are regulated by the cyclin-dependent kinase inhibitor p21 in synovial fibroblasts. <i>Journal of Immunology</i> , 2003 , 170, 838-45	5.3	82
	RANTES modulates TLR4-induced cytokine secretion in human peripheral blood monocytes. <i>Journal of Immunology</i> , 2006, 177, 5077-87 Drug Insight: abatacept for the treatment of rheumatoid arthritis. <i>Nature Clinical Practice Rheumatology</i> , 2006, 2, 654-60 The evolving clinical profile of abatacept (CTLA4-Ig): a novel co-stimulatory modulator for the treatment of rheumatoid arthritis. <i>Arthritis Research</i> , 2005, 7 Suppl 2, 521-5 Amelioration of rat adjuvant-induced arthritis by Met-RANTES. <i>Arthritis and Rheumatism</i> , 2005, 52, 190: Mcl-1 is essential for the survival of synovial fibroblasts in rheumatoid arthritis. <i>Journal of Immunology</i> , 2005, 175, 8337-45 NF-kappaB protects macrophages from lipopolysaccharide-induced cell death: the role of caspase 8 and receptor-interacting protein. <i>Journal of Biological Chemistry</i> , 2005, 280, 41827-34 The role of macrophages in rheumatoid arthritis. <i>Current Pharmaceutical Design</i> , 2005, 11, 569-80 TNF-alpha-induced apoptosis of macrophages following inhibition of NF-kappa B: a central role for disruption of mitochondria. <i>Journal of Immunology</i> , 2004, 172, 1907-15 Fas ligation on macrophages enhances IL-1R1-Toll-like receptor 4 signaling and promotes chronic inflammation. <i>Nature Immunology</i> , 2004, 5, 380-7 Retinoblastoma suppression of matrix metalloproteinase 1, but not interleukin-6, through a p38-dependent pathway in rheumatoid arthritis synovial fibroblasts. <i>Arthritis and Rheumatism</i> , 2004, 50, 78-87 NF-kappaB-regulated expression of cellular FLIP protects rheumatoid arthritis synovial fibroblasts. <i>Pharmaceus Clinics of North America</i> , 2004, 30, 30, 603-25, x Phagocytes: mechanisms of inflammation and tissue destruction. <i>Rheumatis of North America</i> , 2004, 30, 19-39, v Serine phosphorylation of STAT3 is essential for Mcl-1 expression and macrophage survival. <i>Blood</i> , 2003, 102, 344-52 Rheumatoid arthritis synovial fluid macrophages express decreased tumor necrosis factor-related apoptosis-inducing ligand R2 and increased decoy receptor tumor necrosis fact	RANTES modulates TLR4-induced cytokine secretion in human peripheral blood monocytes. <i>Journal of Immunology</i> , 2006, 177, 5077-87 Drug Insight: abatacept for the treatment of rheumatoid arthritis. <i>Nature Clinical Practice Rheumatology</i> , 2006, 2, 654-60 The evolving clinical profile of abatacept (CTLA4-Ig): a novel co-stimulatory modulator for the treatment of rheumatoid arthritis. <i>Arthritis Research</i> , 2005, 7 Suppl 2, 521-5 Amelioration of rat adjuvant-induced arthritis by Met-RANTES. <i>Arthritis and Rheumatism</i> , 2005, 52, 1907-19 Mcl-1 is essential for the survival of synovial fibroblasts in rheumatoid arthritis. <i>Journal of Immunology</i> , 2005, 175, 8337-45 NF-kappaB protects macrophages from lipopolysaccharide-induced cell death: the role of caspase 8 and receptor-interacting protein. <i>Journal of Biological Chemistry</i> , 2005, 280, 41827-34 The role of macrophages in rheumatoid arthritis. <i>Current Pharmaceutical Design</i> , 2005, 11, 569-80 33 TNF-alpha-induced apoptosis of macrophages following inhibition of NF-kappa B: a central role for disruption of mitochondria. <i>Journal of Immunology</i> , 2004, 172, 1907-15 Fas ligation on macrophages enhances IL-1R1-Toll-like receptor 4 signaling and promotes chronic inflammation. <i>Nature Immunology</i> , 2004, 5, 380-7 Retinoblastoma suppression of matrix metalloproteinase 1, but not interleukin-6, through a p38-dependent pathway in rheumatoid arthritis synovial fibroblasts. <i>Arthritis and Rheumatism</i> , 2004, 50, 3844-55 Apoptosis in rheumatoid arthritis: friend or foe. <i>Rheumatic Disease Clinics of North America</i> , 2004, 30, 603-25, x Phagocytes: mechanisms of inflammation and tissue destruction. <i>Rheumatic Disease Clinics of North America</i> , 2004, 30, 19-39, v Serine phosphorylation of STAT3 is essential for Mcl-1 expression and macrophage survival. <i>Blood</i> , 22, 2003, 102, 344-52 Rheumatoid arthritis synovial fluid macrophages express decreased tumor necrosis factor-related apoptosis-inducing ligand R2 and increased decoy receptor tumor necrosis factor-related apopto

40	Apoptosis as a therapeutic tool in rheumatoid arthritis. <i>Nature Reviews Immunology</i> , 2002 , 2, 527-35	36.5	301
39	Rheumatoid arthritis synovial macrophages express the Fas-associated death domain-like interleukin-1beta-converting enzyme-inhibitory protein and are refractory to Fas-mediated apoptosis. <i>Arthritis and Rheumatism</i> , 2001 , 44, 21-30		84
38	Differential expression of chemokine receptors on peripheral blood, synovial fluid, and synovial tissue monocytes/macrophages in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2001 , 44, 1022-32		238
37	Selective lymphocyte chemokine receptor expression in the rheumatoid joint. <i>Arthritis and Rheumatism</i> , 2001 , 44, 2750-60		148
36	Differential expression pattern of the antiapoptotic proteins, Bcl-2 and FLIP, in experimental arthritis. <i>Arthritis and Rheumatism</i> , 2001 , 44, 2899-908		59
35	The Fas-FasL death receptor and PI3K pathways independently regulate monocyte homeostasis. <i>European Journal of Immunology</i> , 2001 , 31, 2421-30	6.1	31
34	Constitutively activated Akt-1 is vital for the survival of human monocyte-differentiated macrophages. Role of Mcl-1, independent of nuclear factor (NF)-kappaB, Bad, or caspase activation. <i>Journal of Experimental Medicine</i> , 2001 , 194, 113-26	16.6	188
33	TNF-alpha gene expression in macrophages: regulation by NF-kappa B is independent of c-Jun or C/EBP beta. <i>Journal of Immunology</i> , 2000 , 164, 4277-85	5.3	179
32	Regulation of IL-6 and IL-8 expression in rheumatoid arthritis synovial fibroblasts: the dominant role for NF-kappa B but not C/EBP beta or c-Jun. <i>Journal of Immunology</i> , 2000 , 165, 7199-206	5.3	176
31	Bcl-2 expression in synovial fibroblasts is essential for maintaining mitochondrial homeostasis and cell viability. <i>Journal of Immunology</i> , 2000 , 164, 5227-35	5.3	97
30	Macrophages require constitutive NF-kappaB activation to maintain A1 expression and mitochondrial homeostasis. <i>Molecular and Cellular Biology</i> , 2000 , 20, 8855-65	4.8	133
29	FLICE-inhibitory protein expression during macrophage differentiation confers resistance to fas-mediated apoptosis. <i>Journal of Experimental Medicine</i> , 1999 , 190, 1679-88	16.6	212
28	C/EBP beta in rheumatoid arthritis: correlation with inflammation, not disease specificity. <i>Clinical Immunology</i> , 1999 , 91, 271-82	9	29
27	Autocrine regulation of collagenase gene expression by TNF-alpha in U937 cells. <i>Journal of Leukocyte Biology</i> , 1996 , 59, 125-32	6.5	20
26	Soluble forms of P-selectin and intercellular adhesion molecule-3 in synovial fluids. <i>Clinical Immunology and Immunopathology</i> , 1996 , 78, 276-82		18
25	Recombinant human interleukin-1 receptor type I in the treatment of patients with active rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 1996 , 39, 257-65		85
24	Hepatocyte growth factor. A cytokine mediating endothelial migration in inflammatory arthritis. <i>Arthritis and Rheumatism</i> , 1996 , 39, 1566-75		62
23	Macrophage inflammatory protein-1 beta: a C-C chemokine in osteoarthritis. <i>Clinical Immunology and Immunopathology</i> , 1995 , 77, 307-14		63

22	Soluble intercellular adhesion molecule-1 in arthritis. <i>Clinical Immunology and Immunopathology</i> , 1994 , 71, 208-15	22
21	Soluble E-selectin in arthritis. <i>Clinical Immunology and Immunopathology</i> , 1993 , 69, 29-35	46
20	Clonal heterogeneity of synovial fluid T lymphocytes in inflammatory synovitis. <i>Clinical Immunology and Immunopathology</i> , 1992 , 63, 28-33	4
19	Activation of synovial fluid T lymphocytes by 60-kd heat-shock proteins in patients with inflammatory synovitis. <i>Arthritis and Rheumatism</i> , 1992 , 35, 43-8	53
18	Gamma/delta T cell receptor positive T cells in the inflammatory joint: lack of association with response to soluble antigens. <i>Cellular Immunology</i> , 1991 , 137, 127-38	5
17	T cell activation by mycobacterial antigens in inflammatory synovitis. <i>Cellular Immunology</i> , 1991 , 4-4	22
16	Identification of a population of large granular lymphocytes obtained from the rheumatoid joint coexpressing the CD3 and CD16 antigens. <i>Clinical Immunology and Immunopathology</i> , 1991 , 58, 409-18	5
15	Immunoregulatory mechanisms present in the maternal circulation during pregnancy. <i>Baillierets Clinical Rheumatology</i> , 1990 , 4, 33-52	23
14	Distribution of CD45RA and CD45RO T-lymphocyte subsets in rheumatoid arthritis synovial tissue. <i>Journal of Clinical Immunology</i> , 1990 , 10, 192-9	33
13	Enhanced cytotoxicity in the rheumatoid joint. Clinical Immunology and Immunopathology, 1990 , 54, 431-41	3
12	Antigenic specificity of rheumatoid synovial fluid lymphocytes. <i>Arthritis and Rheumatism</i> , 1989 , 32, 1371-80	41
11	Increased helper inducer and decreased suppressor inducer phenotypes in the rheumatoid joint. <i>Arthritis and Rheumatism</i> , 1988 , 31, 52-9	78
10	Modulation of spontaneous immunoglobulin production by natural killer cells in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 1986 , 29, 1435-9	6
9	Characterization of the defective autologous mixed lymphocyte response in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 1984 , 27, 1234-44	39
8	Effect of pregnancy on immune complexes and rheumatoid factors in patients with rheumatoid arthritis. <i>American Journal of Medicine</i> , 1983 , 74, 973-9	36
7	The detection of circulating immune complexes and IgG and IgM rheumatoid factors in normal human pregnancy. American Journal of Reproductive Immunology: AJRI: Official Journal of the American Society for the Immunology of Reproduction and the International Coordination Committee	7
6	THE EFFECT OF PREGNANCY ON IMMUNE COMPLEXES AND RHEUMATOID FACTORS IN NORMAL WOMEN AND IN PATIENTS WITH RHEUMATOID ARTHRITIS. <i>The Journal of the Japanese Society of Internal Medicine</i> , 1982 , 71, 1397-1409	
5	Fatal thrombotic thrombocytopenic purpura in a patient with systemic lupus erythematosus. Relationship to circulating immune complexes. <i>Arthritis and Rheumatism</i> , 1981 , 24, 550-3	25

4	IgG rheumatoid factor. Relationship to seropositive rheumatoid arthritis and absence in seronegative disorders. <i>Arthritis and Rheumatism</i> , 1979 , 22, 988-98		85
3	Intermediate complexes formed by self-association of IgG-rheumatoid factors. <i>Annals of the New York Academy of Sciences</i> , 1975 , 256, 82-7	6.5	35
2	The hyperviscosity syndrome in rheumatoid arthritis due to intermediate complexes formed by self-association of IgG-rheumatoid factors. <i>Arthritis and Rheumatism</i> , 1975 , 18, 97-106		58
1	Rheumatoid Arthritis Associated With Hyperviscosity Syndrome and Intermediate Complex Formation. <i>Archives of Internal Medicine</i> , 1975 , 135, 281		18