

Bonnie Ch Kwan

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/833968/publications.pdf>

Version: 2024-02-01

163
papers

5,890
citations

57758

44
h-index

91884

69
g-index

163
all docs

163
docs citations

163
times ranked

6080
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Lipoprotein Metabolism and Lipid Management in Chronic Kidney Disease. <i>Journal of the American Society of Nephrology: JASN</i> , 2007, 18, 1246-1261. | 6.1 | 280 |
| 2 | Endotoxemia is Related to Systemic Inflammation and Atherosclerosis in Peritoneal Dialysis Patients. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2008, 3, 431-436. | 4.5 | 177 |
| 3 | Serum and Urinary Cell-free miR-146a and miR-155 in Patients with Systemic Lupus Erythematosus. <i>Journal of Rheumatology</i> , 2010, 37, 2516-2522. | 2.0 | 174 |
| 4 | Clinical biocompatibility of a neutral peritoneal dialysis solution with minimal glucose-degradation products--A 1-year randomized control trial. <i>Nephrology Dialysis Transplantation</i> , 2006, 22, 552-559. | 0.7 | 152 |
| 5 | Serum and urinary free microRNA level in patients with systemic lupus erythematosus. <i>Lupus</i> , 2011, 20, 493-500. | 1.6 | 142 |
| 6 | Expression of microRNAs in the Urine of Patients With Bladder Cancer. <i>Clinical Genitourinary Cancer</i> , 2012, 10, 106-113. | 1.9 | 134 |
| 7 | Urinary miR-21, miR-29, and miR-93: Novel Biomarkers of Fibrosis. <i>American Journal of Nephrology</i> , 2012, 36, 412-418. | 3.1 | 130 |
| 8 | A Risk Analysis of Continuous Ambulatory Peritoneal Dialysis-Related Peritonitis. <i>Peritoneal Dialysis International</i> , 2005, 25, 374-379. | 2.3 | 127 |
| 9 | Tacrolimus for the treatment of systemic lupus erythematosus with pure class V nephritis. <i>Rheumatology</i> , 2008, 47, 1678-1681. | 1.9 | 115 |
| 10 | Elevated Levels of miR-146a and miR-155 in Kidney Biopsy and Urine from Patients with IgA Nephropathy. <i>Disease Markers</i> , 2011, 30, 171-179. | 1.3 | 109 |
| 11 | Intrarenal Expression of miRNAs in Patients With Hypertensive Nephrosclerosis. <i>American Journal of Hypertension</i> , 2010, 23, 78-84. | 2.0 | 107 |
| 12 | Intrarenal expression of microRNAs in patients with IgA nephropathy. <i>Laboratory Investigation</i> , 2010, 90, 98-103. | 3.7 | 103 |
| 13 | Peritoneal Dialysis Catheter Revision and Replacement by Nephrologist for Peritoneal Dialysis Catheter Malfunction. <i>Nephron</i> , 2018, 138, 214-219. | 1.8 | 103 |
| 14 | Enterobacteriaceae peritonitis complicating peritoneal dialysis: A review of 210 consecutive cases. <i>Kidney International</i> , 2006, 69, 1245-1252. | 5.2 | 102 |
| 15 | Glomerular and tubulointerstitial miR-638, miR-198 and miR-146a expression in lupus nephritis. <i>Nephrology</i> , 2012, 17, 346-351. | 1.6 | 99 |
| 16 | Staphylococcus aureus Peritonitis Complicates Peritoneal Dialysis. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2007, 2, 245-251. | 4.5 | 94 |
| 17 | New-Onset Hyperglycemia in Nondiabetic Chinese Patients Started on Peritoneal Dialysis. <i>American Journal of Kidney Diseases</i> , 2007, 49, 524-532. | 1.9 | 94 |
| 18 | Expression of MicroRNAs in the Urinary Sediment of Patients with IgA Nephropathy. <i>Disease Markers</i> , 2010, 28, 79-86. | 1.3 | 93 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Oral Calcitriol for the Treatment of Persistent Proteinuria in Immunoglobulin A Nephropathy: An Uncontrolled Trial. <i>American Journal of Kidney Diseases</i> , 2008, 51, 724-731. | 1.9 | 87 |
| 20 | Carotid Intima Media Thickness Predicts Cardiovascular Diseases in Chinese Predialysis Patients with Chronic Kidney Disease. <i>Journal of the American Society of Nephrology: JASN</i> , 2007, 18, 1966-1972. | 6.1 | 85 |
| 21 | Elevated levels of miR-146a and miR-155 in kidney biopsy and urine from patients with IgA nephropathy. <i>Disease Markers</i> , 2011, 30, 171-9. | 1.3 | 85 |
| 22 | Hypokalemia in Chinese Peritoneal Dialysis Patients: Prevalence and Prognostic Implication. <i>American Journal of Kidney Diseases</i> , 2005, 46, 128-135. | 1.9 | 84 |
| 23 | The gene expression of type 17 T-helper cell-related cytokines in the urinary sediment of patients with systemic lupus erythematosus. <i>Rheumatology</i> , 2009, 48, 1491-1497. | 1.9 | 79 |
| 24 | Podocyte Loss in Human Hypertensive Nephrosclerosis. <i>American Journal of Hypertension</i> , 2009, 22, 300-306. | 2.0 | 79 |
| 25 | Expression of miR-146a and miR-155 in the urinary sediment of systemic lupus erythematosus. <i>Clinical Rheumatology</i> , 2012, 31, 435-440. | 2.2 | 77 |
| 26 | Expression of microRNAs in the urinary sediment of patients with IgA nephropathy. <i>Disease Markers</i> , 2010, 28, 79-86. | 1.3 | 71 |
| 27 | Change in bacterial aetiology of peritoneal dialysis-related peritonitis over 10 years: experience from a centre in south-east Asia. <i>Clinical Microbiology and Infection</i> , 2005, 11, 837-839. | 6.0 | 70 |
| 28 | Peritoneal Dialysis as the First-line Renal Replacement Therapy in Patients With Autosomal Dominant Polycystic Kidney Disease. <i>American Journal of Kidney Diseases</i> , 2011, 57, 903-907. | 1.9 | 66 |
| 29 | A risk analysis of continuous ambulatory peritoneal dialysis-related peritonitis. <i>Peritoneal Dialysis International</i> , 2005, 25, 374-9. | 2.3 | 66 |
| 30 | Predictors of Residual Renal Function Decline in Patients Undergoing Continuous Ambulatory Peritoneal Dialysis. <i>Peritoneal Dialysis International</i> , 2015, 35, 180-188. | 2.3 | 65 |
| 31 | Cellular and morphological changes during neointimal hyperplasia development in a porcine arteriovenous graft model. <i>Nephrology Dialysis Transplantation</i> , 2007, 22, 3139-3146. | 0.7 | 64 |
| 32 | Recurrent and Relapsing Peritonitis: Causative Organisms and Response to Treatment. <i>American Journal of Kidney Diseases</i> , 2009, 54, 702-710. | 1.9 | 62 |
| 33 | Associations of Body Size with Metabolic Syndrome and Mortality in Moderate Chronic Kidney Disease. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2007, 2, 992-998. | 4.5 | 60 |
| 34 | Bioimpedance Spectroscopy for the Detection of Fluid Overload in Chinese Peritoneal Dialysis Patients. <i>Peritoneal Dialysis International</i> , 2014, 34, 409-416. | 2.3 | 60 |
| 35 | Coagulase Negative Staphylococcal Peritonitis in Peritoneal Dialysis Patients. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2008, 3, 91-97. | 4.5 | 57 |
| 36 | Severe Acute Respiratory Syndrome in Dialysis Patients. <i>Journal of the American Society of Nephrology: JASN</i> , 2004, 15, 1883-1888. | 6.1 | 54 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Gene expression of TWEAK/Fn14 and IP10/CXCR3 in glomerulus and tubulointerstitium of patients with lupus nephritis. <i>Nephrology</i> , 2011, 16, 426-432. | 1.6 | 50 |
| 38 | Urinary sediment miRNA levels in adult nephrotic syndrome. <i>Clinica Chimica Acta</i> , 2013, 418, 5-11. | 1.1 | 49 |
| 39 | Urinary mitochondrial DNA level is an indicator of intra-renal mitochondrial depletion and renal scarring in diabetic nephropathy. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, 784-788. | 0.7 | 49 |
| 40 | Urinary mRNA expression of ACE and ACE2 in human type 2 diabetic nephropathy. <i>Diabetologia</i> , 2008, 51, 1062-1067. | 6.3 | 47 |
| 41 | Asymptomatic isolated microscopic haematuria: long-term follow-up. <i>QJM - Monthly Journal of the Association of Physicians</i> , 2004, 97, 739-745. | 0.5 | 46 |
| 42 | Peritoneal Albumin Excretion is a Strong Predictor of Cardiovascular Events in Peritoneal Dialysis Patients: A Prospective Cohort Study. <i>Peritoneal Dialysis International</i> , 2005, 25, 445-452. | 2.3 | 46 |
| 43 | Predicting 12-Month Mortality for Peritoneal Dialysis Patients Using the "Surprise" Question. <i>Peritoneal Dialysis International</i> , 2013, 33, 60-66. | 2.3 | 45 |
| 44 | Prevalence of silent kidney disease in Hong Kong: The Screening for Hong Kong Asymptomatic Renal Population and Evaluation (SHARE) program. <i>Kidney International</i> , 2005, 67, S36-S40. | 5.2 | 44 |
| 45 | Urinary Expression of Kidney Injury Markers in Renal Transplant Recipients. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2010, 5, 2329-2337. | 4.5 | 44 |
| 46 | Urinary messenger RNA expression of podocyte-associated molecules in patients with diabetic nephropathy treated by angiotensin-converting enzyme inhibitor and angiotensin receptor blocker. <i>European Journal of Endocrinology</i> , 2008, 158, 317-322. | 3.7 | 42 |
| 47 | Geriatric Nutritional Risk Index as a Screening Tool for Malnutrition in Patients on Chronic Peritoneal Dialysis. , 2010, 20, 29-37. | | 41 |
| 48 | Public lacks knowledge on chronic kidney disease: telephone survey. <i>Hong Kong Medical Journal</i> , 2014, 20, 139-44. | 0.1 | 40 |
| 49 | Discrepancy between Intrarenal Messenger RNA and Protein Expression of ACE and ACE2 in Human Diabetic Nephropathy. <i>American Journal of Nephrology</i> , 2009, 29, 524-531. | 3.1 | 39 |
| 50 | Repeat Renal Biopsy in Lupus Nephritis: A Change in Histological Pattern Is Common. <i>American Journal of Nephrology</i> , 2011, 34, 220-225. | 3.1 | 39 |
| 51 | Prognostic Value of Arterial Pulse Wave Velocity in Peritoneal Dialysis Patients. <i>American Journal of Nephrology</i> , 2012, 35, 127-133. | 3.1 | 39 |
| 52 | Urinary sediment ICAM-1 level in lupus nephritis. <i>Lupus</i> , 2012, 21, 1190-1195. | 1.6 | 38 |
| 53 | Asymptomatic fluid overload predicts survival and cardiovascular event in incident Chinese peritoneal dialysis patients. <i>PLoS ONE</i> , 2018, 13, e0202203. | 2.5 | 38 |
| 54 | Cefazolin plus Ceftazidime versus Imipenem / Cilastatin Monotherapy for Treatment of Capd Peritonitis " a Randomized Controlled Trial. <i>Peritoneal Dialysis International</i> , 2004, 24, 440-446. | 2.3 | 37 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Circulating bacterial-derived DNA fragments as a marker * of systemic inflammation in peritoneal dialysis. <i>Nephrology Dialysis Transplantation</i> , 2013, 28, 2139-2145. | 0.7 | 35 |
| 56 | Long-term Outcome of Biopsy-Proven Minimal Change Nephropathy in Chinese Adults. <i>American Journal of Kidney Diseases</i> , 2015, 65, 710-718. | 1.9 | 35 |
| 57 | Bacteria-Derived DNA Fragment in Peritoneal Dialysis Effluent as a Predictor of Relapsing Peritonitis. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2013, 8, 1935-1941. | 4.5 | 31 |
| 58 | Circulating Bacterial-Derived DNA Fragment Level Is a Strong Predictor of Cardiovascular Disease in Peritoneal Dialysis Patients. <i>PLoS ONE</i> , 2015, 10, e0125162. | 2.5 | 31 |
| 59 | Repeat Peritonitis in Peritoneal Dialysis. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2011, 6, 827-833. | 4.5 | 30 |
| 60 | Frailty in Chinese Peritoneal Dialysis Patients: Prevalence and Prognostic Significance. <i>Kidney and Blood Pressure Research</i> , 2016, 41, 736-745. | 2.0 | 30 |
| 61 | Urinary FOXP3 mRNA in patients with lupus nephritis--relation with disease activity and treatment response. <i>Rheumatology</i> , 2009, 48, 755-760. | 1.9 | 29 |
| 62 | Expression of ACE and ACE2 in Patients with Hypertensive Nephrosclerosis. <i>Kidney and Blood Pressure Research</i> , 2011, 34, 141-149. | 2.0 | 28 |
| 63 | Peritoneal albumin excretion is a strong predictor of cardiovascular events in peritoneal dialysis patients: a prospective cohort study. <i>Peritoneal Dialysis International</i> , 2005, 25, 445-52. | 2.3 | 28 |
| 64 | The clinical course of peritoneal dialysis-related peritonitis caused by <i>Corynebacterium</i> species. <i>Nephrology Dialysis Transplantation</i> , 2005, 20, 2793-2796. | 0.7 | 27 |
| 65 | Tenckhoff Catheter Insertion by Nephrologists: Open Dissection Technique. <i>Peritoneal Dialysis International</i> , 2010, 30, 524-527. | 2.3 | 26 |
| 66 | Clinical manifestation of macrolide antibiotic toxicity in CKD and dialysis patients. <i>CKJ: Clinical Kidney Journal</i> , 2014, 7, 507-512. | 2.9 | 26 |
| 67 | Urinary miRNA profile for the diagnosis of IgA nephropathy. <i>BMC Nephrology</i> , 2019, 20, 77. | 1.8 | 26 |
| 68 | mRNA Expression of Target Genes in the Urinary Sediment as a Noninvasive Prognostic Indicator of CKD. <i>American Journal of Kidney Diseases</i> , 2006, 47, 578-586. | 1.9 | 25 |
| 69 | Fatal Pancytopenia in a Hemodialysis Patient After Treatment With Low-Dose Methotrexate. <i>Journal of Clinical Rheumatology</i> , 2009, 15, 177-180. | 0.9 | 25 |
| 70 | Depression and Physical Frailty Have Additive Effect on the Nutritional Status and Clinical Outcome of Chinese Peritoneal Dialysis. <i>Kidney and Blood Pressure Research</i> , 2018, 43, 914-923. | 2.0 | 25 |
| 71 | Treatment of Metabolic Syndrome in Peritoneal Dialysis Patients. <i>Peritoneal Dialysis International</i> , 2009, 29, 149-152. | 2.3 | 24 |
| 72 | Treatment of Early Immunoglobulin A Nephropathy by Angiotensin-converting Enzyme Inhibitor. <i>American Journal of Medicine</i> , 2013, 126, 162-168. | 1.5 | 24 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Randomized controlled study of icodextrin on the treatment of peritoneal dialysis patients during acute peritonitis. <i>Nephrology Dialysis Transplantation</i> , 2014, 29, 1438-1443. | 0.7 | 24 |
| 74 | Effect of Membrane Permeability on Inflammation and Arterial Stiffness. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2010, 5, 652-658. | 4.5 | 23 |
| 75 | Metabolic Syndrome in Peritoneal Dialysis Patients. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2014, 9, 779-787. | 4.5 | 23 |
| 76 | Urinary mitochondrial DNA level in non-diabetic chronic kidney diseases. <i>Clinica Chimica Acta</i> , 2018, 484, 36-39. | 1.1 | 23 |
| 77 | Impact of Early Nephrology Referral on Mortality and Hospitalization in Peritoneal Dialysis Patients. <i>Peritoneal Dialysis International</i> , 2008, 28, 371-376. | 2.3 | 22 |
| 78 | Endotoxemia is Associated with Better Clinical Outcome in Incident Chinese Peritoneal Dialysis Patients: A Prospective Cohort Study. <i>Peritoneal Dialysis International</i> , 2010, 30, 178-186. | 2.3 | 21 |
| 79 | Increased subcutaneous insulin requirements in diabetic patients recently commenced on peritoneal dialysis. <i>Nephrology Dialysis Transplantation</i> , 2007, 22, 1697-1702. | 0.7 | 20 |
| 80 | Peritoneal protein clearance predicts mortality in peritoneal dialysis patients. <i>Clinical and Experimental Nephrology</i> , 2019, 23, 551-560. | 1.6 | 20 |
| 81 | Connective Tissue Growth Factor Is Responsible for Transforming Growth Factor-Beta-Induced Peritoneal Mesothelial Cell Apoptosis. <i>Nephron Experimental Nephrology</i> , 2006, 103, e166-e174. | 2.2 | 19 |
| 82 | CARDIOVASCULAR AND SURVIVAL PARADOXES IN DIALYSIS PATIENTS: A Story Half Untold: Adiposity, Adipokines and Outcomes in Dialysis Population. <i>Seminars in Dialysis</i> , 2007, 20, 493-497. | 1.3 | 19 |
| 83 | Peroxisome Proliferator-Activated Receptor-Gamma Gene Polymorphism and Risk of Cardiovascular Disease in Patients with Diabetic Nephropathy. <i>American Journal of Nephrology</i> , 2008, 28, 715-722. | 3.1 | 19 |
| 84 | Intra-renal and urinary mRNA expression of podocyte-associated molecules for the estimation of glomerular podocyte loss. <i>Renal Failure</i> , 2010, 32, 372-379. | 2.1 | 19 |
| 85 | The Safety and Short-Term Efficacy of Aliskiren in the Treatment of Immunoglobulin A Nephropathy – A Randomized Cross-Over Study. <i>PLoS ONE</i> , 2013, 8, e62736. | 2.5 | 19 |
| 86 | Manifestation of tranexamic acid toxicity in chronic kidney disease and kidney transplant patients: A report of four cases and review of literature. <i>Nephrology</i> , 2017, 22, 316-321. | 1.6 | 19 |
| 87 | Cross sectional survey on the concerns and anxiety of patients waiting for organ transplants. <i>Nephrology</i> , 2012, 17, 514-518. | 1.6 | 18 |
| 88 | Effect of Using Ultrapure Dialysate for Hemodialysis on the Level of Circulating Bacterial Fragment in Renal Failure Patients. <i>Nephron Clinical Practice</i> , 2013, 123, 246-253. | 2.3 | 18 |
| 89 | Urinary mitochondrial DNA level as a biomarker of tissue injury in non-diabetic chronic kidney diseases. <i>BMC Nephrology</i> , 2018, 19, 367. | 1.8 | 18 |
| 90 | Relationship between Plasma Endocan Level and Clinical Outcome of Chinese Peritoneal Dialysis Patients. <i>Kidney and Blood Pressure Research</i> , 2019, 44, 1259-1270. | 2.0 | 18 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | Severe acute respiratory syndrome in a hemodialysis patient. American Journal of Kidney Diseases, 2003, 42, 1069-1074. | 1.9 | 17 |
| 92 | Peritoneal dialysis related peritonitis caused by <i>Gordonia</i> species: Report of four cases and literature review. Nephrology, 2014, 19, 379-383. | 1.6 | 17 |
| 93 | Urinary mRNA levels of ELR-negative CXC chemokine ligand and extracellular matrix in diabetic nephropathy. Diabetes/Metabolism Research and Reviews, 2015, 31, 699-706. | 4.0 | 17 |
| 94 | The Effect of Neutral Peritoneal Dialysis Solution with Low Glucose-Degradation-Product on the Fluid Status and Body Composition – A Randomized Control Trial. PLoS ONE, 2015, 10, e0141425. | 2.5 | 17 |
| 95 | Urinary Mitochondrial DNA Level as a Biomarker of Acute Kidney Injury Severity. Kidney Diseases (Basel), 2017, 1, 1-17. | 2.5 | 17 |
| 96 | Peritoneal dialysis-related peritonitis caused by Pseudomonas species: Insight from a post-millennial case series. PLoS ONE, 2018, 13, e0196499. | 2.5 | 17 |
| 97 | Cefazolin plus ceftazidime versus imipenem/cilastatin monotherapy for treatment of CAPD peritonitis—a randomized controlled trial. Peritoneal Dialysis International, 2004, 24, 440-6. | 2.3 | 17 |
| 98 | Longitudinal Changes of Cardiothoracic Ratio and Vascular Pedicle Width as Predictors of Volume Status during One Year in Chinese Peritoneal Dialysis Patients. Kidney and Blood Pressure Research, 2009, 32, 45-50. | 2.0 | 16 |
| 99 | Relation between MicroRNA Expression in Peritoneal Dialysis Effluent and Peritoneal Transport Characteristics. Disease Markers, 2012, 33, 35-42. | 1.3 | 16 |
| 100 | Extra-high-dose hepatitis B vaccination does not confer longer serological protection in peritoneal dialysis patients: a randomized controlled trial. Nephrology Dialysis Transplantation, 2010, 25, 2303-2309. | 0.7 | 15 |
| 101 | Automated peritoneal dialysis in Hong Kong: There are two distinct groups of patients. Nephrology, 2013, 18, 356-364. | 1.6 | 15 |
| 102 | Peritonitis before Peritoneal Dialysis Training: Analysis of Causative Organisms, Clinical Outcomes, Risk Factors, and Long-Term Consequences. Clinical Journal of the American Society of Nephrology: CJASN, 2016, 11, 1219-1226. | 4.5 | 15 |
| 103 | Treatment of metabolic syndrome in peritoneal dialysis patients. Peritoneal Dialysis International, 2009, 29 Suppl 2, S149-52. | 2.3 | 15 |
| 104 | Adherence to peritoneal dialysis training schedule. Nephrology Dialysis Transplantation, 2006, 22, 545-551. | 0.7 | 14 |
| 105 | Outcome of Hemodialysis Patients Who Had Failed Peritoneal Dialysis. Nephron Clinical Practice, 2010, 116, c300-c306. | 2.3 | 14 |
| 106 | Intrarenal and Urinary Th9 and Th22 Cytokine Gene Expression in Lupus Nephritis. Journal of Rheumatology, 2015, 42, 1150-1155. | 2.0 | 14 |
| 107 | Newer antibiotics for the treatment of peritoneal dialysis-related peritonitis. CKJ: Clinical Kidney Journal, 2016, 9, 616-623. | 2.9 | 14 |
| 108 | Causes of nephrotic syndrome and nephrotic-range proteinuria are different in adult Chinese patients: A single centre study over 33 years. Nephrology, 2018, 23, 565-572. | 1.6 | 14 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Influence of climate on the incidence of thiazide-induced hyponatraemia. <i>International Journal of Clinical Practice</i> , 2006, 61, 449-452. | 1.7 | 13 |
| 110 | Unexplained Exudative Pleural Effusion in Chronic Peritoneal Dialysis Patients. <i>Peritoneal Dialysis International</i> , 2010, 30, 534-540. | 2.3 | 13 |
| 111 | Life expectancy of Chinese patients with chronic kidney disease without dialysis. <i>Nephrology</i> , 2011, 16, 715-719. | 1.6 | 13 |
| 112 | Relationship of Intrarenal Gene Expression and the Histological Class of Lupus Nephritis – A Study on Repeat Renal Biopsy. <i>Journal of Rheumatology</i> , 2012, 39, 1942-1947. | 2.0 | 13 |
| 113 | Monitoring of urinary messenger RNA levels for the prediction of flare in systemic lupus erythematosus. <i>Clinica Chimica Acta</i> , 2012, 413, 448-455. | 1.1 | 13 |
| 114 | Fracture risk after thiazide-associated hyponatraemia. <i>Internal Medicine Journal</i> , 2012, 42, 760-764. | 0.8 | 13 |
| 115 | Association of interleukin-18 promoter polymorphism and atherosclerotic diseases in Chinese patients with diabetic nephropathy. <i>Nephrology</i> , 2009, 14, 606-612. | 1.6 | 12 |
| 116 | Arterial Pulse Wave Velocity and Peritoneal Transport Characteristics Independently Predict Hospitalization in Chinese Peritoneal Dialysis Patients. <i>Peritoneal Dialysis International</i> , 2010, 30, 80-85. | 2.3 | 12 |
| 117 | Persistent Symptomatic Intra-Abdominal Collection after Catheter Removal for Pd-Related Peritonitis. <i>Peritoneal Dialysis International</i> , 2011, 31, 34-38. | 2.3 | 12 |
| 118 | Serial monitoring of nutritional status in Chinese peritoneal dialysis patients by Subjective Global Assessment and comprehensive Malnutrition Inflammation Score. <i>Nephrology</i> , 2009, 14, 143-147. | 1.6 | 11 |
| 119 | Relationship between CRP Polymorphism and Cardiovascular Events in Chinese Peritoneal Dialysis Patients. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2012, 7, 304-309. | 4.5 | 11 |
| 120 | Peritoneal dialysis effluent miR-21 and miR-589 levels correlate with longitudinal change in peritoneal transport characteristics. <i>Clinica Chimica Acta</i> , 2017, 464, 106-112. | 1.1 | 11 |
| 121 | Treatment of Enterococcal Peritonitis in Peritoneal Dialysis Patients by Oral Amoxicillin or Intra-Peritoneal Vancomycin: a Retrospective Study. <i>Kidney and Blood Pressure Research</i> , 2017, 42, 837-843. | 2.0 | 11 |
| 122 | Metabolic syndrome in peritoneal dialysis patients. <i>CKJ: Clinical Kidney Journal</i> , 2008, 1, 206-214. | 2.9 | 10 |
| 123 | Hypokalaemia and cardiac risk in peritoneal dialysis patients. <i>Nature Reviews Nephrology</i> , 2012, 8, 501-503. | 9.6 | 10 |
| 124 | The width of the basement membrane does not influence clinical presentation or outcome of thin glomerular basement membrane disease with persistent hematuria. <i>Kidney International</i> , 2010, 78, 1041-1046. | 5.2 | 9 |
| 125 | Dialysate bacterial endotoxin as a prognostic indicator of peritoneal dialysis related peritonitis. <i>Nephrology</i> , 2016, 21, 1069-1072. | 1.6 | 9 |
| 126 | Plasma Mitochondrial DNA Level is a Prognostic Marker in Peritoneal Dialysis Patients. <i>Kidney and Blood Pressure Research</i> , 2016, 41, 402-412. | 2.0 | 9 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 127 | Depression does not predict clinical outcome of Chinese peritoneal Dialysis patients after adjusting for the degree of frailty. <i>BMC Nephrology</i> , 2020, 21, 329. | 1.8 | 9 |
| 128 | Relation between microRNA expression in peritoneal dialysis effluent and peritoneal transport characteristics. <i>Disease Markers</i> , 2012, 33, 35-42. | 1.3 | 9 |
| 129 | Podocyte mRNA in the urinary sediment of minimal change nephropathy and focal segmental glomerulosclerosis. <i>Clinical Nephrology</i> , 2015, 84 (2015), 198-205. | 0.7 | 9 |
| 130 | Factors Associated with Sudden Death in Peritoneal Dialysis Patients. <i>Peritoneal Dialysis International</i> , 2009, 29, 58-63. | 2.3 | 8 |
| 131 | Effect of cinacalcet treatment on vascular arterial stiffness among peritoneal dialysis patients with secondary hyperparathyroidism. <i>Nephrology</i> , 2014, 19, 339-344. | 1.6 | 8 |
| 132 | Weight change during the first year of peritoneal dialysis: Risk factors and prognostic implications. <i>Hong Kong Journal of Nephrology</i> , 2015, 17, 28-35. | 0.0 | 8 |
| 133 | Longitudinal Changes of NF- κ B Downstream Mediators and Peritoneal Transport Characteristics in Incident Peritoneal Dialysis Patients. <i>Scientific Reports</i> , 2020, 10, 6440. | 3.3 | 8 |
| 134 | Adipose expression of miR-130b and miR-17-5p with wasting, cardiovascular event and mortality in advanced chronic kidney disease patients. <i>Nephrology Dialysis Transplantation</i> , 2022, 37, 1935-1943. | 0.7 | 8 |
| 135 | Measurements on the routine chest radiograph as prognostic markers in Chinese peritoneal dialysis patients. <i>Clinical Nephrology</i> , 2011, 76, 16-22. | 0.7 | 8 |
| 136 | Urinary sediment mRNA level of extracellular matrix molecules in adult nephrotic syndrome. <i>Clinica Chimica Acta</i> , 2016, 456, 157-162. | 1.1 | 7 |
| 137 | The choice of comorbidity scoring system in Chinese peritoneal dialysis patients. <i>Clinical and Experimental Nephrology</i> , 2018, 22, 159-166. | 1.6 | 7 |
| 138 | Haemoglobin variability in Chinese pre-dialysis CKD patients not receiving erythropoietin. <i>Nephrology Dialysis Transplantation</i> , 2011, 26, 2919-2924. | 0.7 | 5 |
| 139 | The relationship between bone morphogenic protein-7 and peritoneal transport characteristics. <i>Nephrology Dialysis Transplantation</i> , 2008, 23, 2989-2994. | 0.7 | 4 |
| 140 | Relationship between Myeloid-Related Protein 8/14 and Survival of Chinese Peritoneal Dialysis Patients. <i>Kidney and Blood Pressure Research</i> , 2012, 35, 489-496. | 2.0 | 4 |
| 141 | Relationship between serum levels of tumour necrosis factor-related apoptosis-inducing ligand and the survival of Chinese peritoneal dialysis patients. <i>Nephrology</i> , 2012, 17, 466-471. | 1.6 | 4 |
| 142 | Urinary mRNA in Systemic Lupus Erythematosus. <i>Advances in Clinical Chemistry</i> , 2013, 62, 197-219. | 3.7 | 4 |
| 143 | Long-term outcome of biopsy-proven minimal-change nephrotic syndrome in Chinese children. <i>Hong Kong Journal of Nephrology</i> , 2013, 15, 22-27. | 0.0 | 4 |
| 144 | Campylobacter Peritonitis Complicating Peritoneal Dialysis: A Review of 12 Consecutive Cases. <i>Peritoneal Dialysis International</i> , 2013, 33, 189-194. | 2.3 | 4 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 145 | Functional and histological improvement after everolimus rescue of chronic allograft dysfunction in renal transplant recipients. <i>Therapeutics and Clinical Risk Management</i> , 2015, 11, 829. | 2.0 | 4 |
| 146 | Peritoneal inflammation and fibrosis in C-reactive protein transgenic mice undergoing peritoneal dialysis solution treatment. <i>Nephrology</i> , 2017, 22, 125-132. | 1.6 | 4 |
| 147 | The efficacy of managing fluid overload in chronic peritoneal dialysis patients by a structured nurse-led intervention protocol. <i>BMC Nephrology</i> , 2019, 20, 454. | 1.8 | 4 |
| 148 | Adipose and serum zinc alpha-2-glycoprotein (ZAG) expressions predict longitudinal change of adiposity, wasting and predict survival in dialysis patients. <i>Scientific Reports</i> , 2022, 12, . | 3.3 | 4 |
| 149 | Predictors and prognostic significance of persistent fluid overload: A longitudinal study in Chinese peritoneal dialysis patients. <i>Peritoneal Dialysis International</i> , 2023, 43, 252-262. | 2.3 | 4 |
| 150 | In Reply To "The Beneficial Effect of Oral Calcitriol Treatment on Proteinuria in IgA Nephropathy: Another Point of View". <i>American Journal of Kidney Diseases</i> , 2008, 52, 805. | 1.9 | 3 |
| 151 | Relationship between HSP70-2 A+1267G Polymorphism and Cardiovascular Events of Chinese Peritoneal Dialysis Patients. <i>Nephron Clinical Practice</i> , 2014, 128, 153-158. | 2.3 | 3 |
| 152 | Is peritoneal dialysis kinder for the heart?. <i>Peritoneal Dialysis International</i> , 2011, 31, 135-7. | 2.3 | 3 |
| 153 | Relationship Between β 1-Adrenergic Receptor Polymorphisms and Cardiovascular Disease in Peritoneal Dialysis Patients. <i>Hong Kong Journal of Nephrology</i> , 2008, 10, 58-63. | 0.0 | 2 |
| 154 | Relationship between β 2-adrenergic receptor polymorphisms and cardiovascular disease in patients with diabetic nephropathy. <i>Nephrology</i> , 2010, 15, 242-247. | 1.6 | 2 |
| 155 | Relationship between glutathione S-transferase M1 polymorphism and clinical outcomes in Chinese peritoneal dialysis patients. <i>Journal of Nephrology</i> , 2012, 25, 310-316. | 2.0 | 2 |
| 156 | Characteristics and clinical outcomes of living renal donors in Hong Kong. <i>Hong Kong Medical Journal</i> , 2018, 24, 11-17. | 0.1 | 2 |
| 157 | Muscle infarction in peritoneal dialysis patients1 1Editor Note: The corresponding author declined to respond.. <i>American Journal of Kidney Diseases</i> , 2003, 42, 1102-1103. | 1.9 | 1 |
| 158 | Clinical practice guidelines for the provision of renal service in Hong Kong: Accreditation of Renal Unit. <i>Nephrology</i> , 2019, 24, 130-132. | 1.6 | 1 |
| 159 | Menstrual loss of renal function: a case of mefenamic acid induced renal cortical necrosis. <i>Clinical Nephrology</i> , 2016, 86, 162-164. | 0.7 | 1 |
| 160 | SUCCESSFUL EMBOLIZATION OF TRANSPLANT KIDNEY ARTERIOVENOUS FISTULA. <i>Nephrology</i> , 2007, 12, 623-624. | 1.6 | 0 |
| 161 | Laparoscopic live donor nephrectomy: Current practice and results of renal transplantation. <i>Surgical Practice</i> , 2012, 16, 17-21. | 0.2 | 0 |
| 162 | Spurious hypernatraemia. <i>Nephrology</i> , 2013, 18, 531-532. | 1.6 | 0 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 163 | Therapeutic drug monitoring of once-daily tacrolimus (Advagraf) in a gastrectomized kidney transplant recipient. <i>Nephrology</i> , 2017, 22, 184-184. | 1.6 | 0 |