Stefano Ciardullo

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/8340669/stefano-ciardullo-publications-by-year.pdf

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

8 15 41 294 h-index g-index citations papers 625 46 5.05 5.3 L-index avg, IF ext. citations ext. papers

#	Paper	IF	Citations
41	Prevalence of Elevated Liver Stiffness Among Potential Candidates for Bariatric Surgery in the United States <i>Obesity Surgery</i> , 2022 , 32, 712	3.7	O
40	Liver Stiffness, Albuminuria and Chronic Kidney Disease in Patients with NAFLD: A Systematic Review and Meta-Analysis <i>Biomolecules</i> , 2022 , 12,	5.9	3
39	Advances in fibrosis biomarkers in nonalcoholic fatty liver disease <i>Advances in Clinical Chemistry</i> , 2022 , 106, 33-65	5.8	O
38	Soluble EKlotho levels, glycemic control and renal function in US adults with type 2 diabetes <i>Acta Diabetologica</i> , 2022 , 1	3.9	O
37	Sex-related association of NAFLD and liver fibrosis with body fat distribution in the general US population <i>American Journal of Clinical Nutrition</i> , 2022 ,	7	2
36	Hepatitis C virus infection and diabetes: a complex bidirectional relationship <i>Diabetes Research and Clinical Practice</i> , 2022 , 109870	7.4	
35	Peripheral artery disease and all-cause and cardiovascular mortality in patients with NAFLD <i>Journal of Endocrinological Investigation</i> , 2022 , 1	5.2	1
34	High Prevalence of Advanced Liver Fibrosis Assessed by Transient Elastography Among U.S. Adults With Type 2 Diabetes. <i>Diabetes Care</i> , 2021 , 44, 519-525	14.6	36
33	Nonalcoholic fatty liver disease and risk of incident hypertension: a systematic review and meta-analysis. <i>European Journal of Gastroenterology and Hepatology</i> , 2021 ,	2.2	6
32	Blood pressure, glycemic status and advanced liver fibrosis assessed by transient elastography in the general United States population. <i>Journal of Hypertension</i> , 2021 , 39, 1621-1627	1.9	4
31	Prevalence of NAFLD, MAFLD and associated advanced fibrosis in the contemporary United States population. <i>Liver International</i> , 2021 , 41, 1290-1293	7.9	39
30	Sodium-glucose transporter 2 inhibitors for renal and cardiovascular protection in US adults with type 2 diabetes: Impact of the 2020 KDIGO clinical practice guidelines. <i>Pharmacological Research</i> , 2021 , 166, 105530	10.2	2
29	NAFLD and Liver Fibrosis Are Not Associated With Reduced Femoral Bone Mineral Density in the General US Population. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021 , 106, e2856-e2865	5.6	4
28	Seasonal variation in estimated cardiovascular risk in patients with type 2 diabetes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021 , 31, 1494-1500	4.5	3
27	Current type 2 diabetes, rather than previous gestational diabetes, is associated with liver disease in U.S. Women. <i>Diabetes Research and Clinical Practice</i> , 2021 , 177, 108879	7.4	3
26	Nonalcoholic Fatty Liver Disease, Liver Fibrosis and Cardiovascular Disease in the Adult US Population. <i>Frontiers in Endocrinology</i> , 2021 , 12, 711484	5.7	1
25	Liver fibrosis assessed by transient elastography is independently associated with albuminuria in the general United States population. <i>Digestive and Liver Disease</i> , 2021 , 53, 866-872	3.3	7

(2020-2021)

24	Prevalence of Liver Steatosis and Fibrosis Detected by Transient Elastography in Adolescents in the 2017-2018 National Health and Nutrition Examination Survey. <i>Clinical Gastroenterology and Hepatology</i> , 2021 , 19, 384-390.e1	6.9	27
23	Impact of diabetes on COVID-19-related in-hospital mortality: a retrospective study from Northern Italy. <i>Journal of Endocrinological Investigation</i> , 2021 , 44, 843-850	5.2	23
22	Hypercortisolism and altered glucose homeostasis in obese patients in the pre-bariatric surgery assessment. <i>Diabetes/Metabolism Research and Reviews</i> , 2021 , 37, e3389	7.5	3
21	Baseline TSH levels and short-term weight loss after different procedures of bariatric surgery. International Journal of Obesity, 2021, 45, 326-330	5.5	1
20	Lack of awareness of liver organ damage in patients with type 2 diabetes. <i>Acta Diabetologica</i> , 2021 , 58, 651-655	3.9	2
19	An unexpected bilateral mass after total thyroidectomy. <i>Endocrine</i> , 2021 , 73, 758-761	4	
18	Metabolic and Psychological Features are Associated with Weight Loss 12 Months After Sleeve Gastrectomy. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021 , 106, e3087-e3097	5.6	1
17	Statin use is associated with lower prevalence of advanced liver fibrosis in patients with type 2 diabetes. <i>Metabolism: Clinical and Experimental</i> , 2021 , 121, 154752	12.7	7
16	Reply to "Liver fibrosis assessed by transient elastography and albuminuria". <i>Digestive and Liver Disease</i> , 2021 , 53, 1056	3.3	
15	Comparing medication persistence among patients with type 2 diabetes using sodium-glucose cotransporter 2 inhibitors or glucagon-like peptide-1 receptor agonists in real-world setting. <i>Diabetes Research and Clinical Practice</i> , 2021 , 180, 109035	7.4	3
14	Metabolic Syndrome, and Not Obesity, Is Associated with Chronic Kidney Disease. <i>American Journal of Nephrology</i> , 2021 , 52, 666-672	4.6	3
13	Comment on "An Observational Data Meta-Analysis on the Differences in Prevalence and Risk Factors Between MAFLD vs NAFLD" <i>Clinical Gastroenterology and Hepatology</i> , 2021 ,	6.9	
12	Resting Energy Expenditure in Obese Women with Primary Hypothyroidism and Appropriate Levothyroxine Replacement Therapy. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020 , 105,	5.6	4
11	Nonalcoholic Fatty Liver Disease and Advanced Fibrosis in US Adults Across Blood Pressure Categories. <i>Hypertension</i> , 2020 , 76, 562-568	8.5	14
10	Resting Whole Body Energy Metabolism in Class 3 Obesity; from Preserved Insulin Sensitivity to Overt Type 2 Diabetes. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy,</i> 2020 , 13, 489-4	9 3 ·4	2
9	Impact of using different biomarkers of liver fibrosis on hepatologic referral of individuals with severe obesity and NAFLD. <i>Journal of Endocrinological Investigation</i> , 2020 , 43, 1019-1026	5.2	5
8	Screening for non-alcoholic fatty liver disease in type 2 diabetes using non-invasive scores and association with diabetic complications. <i>BMJ Open Diabetes Research and Care</i> , 2020 , 8,	4.5	35
7	Is Switching from Oral Antidiabetic Therapy to Insulin Associated with an Increased Fracture Risk?. <i>Clinical Orthopaedics and Related Research</i> , 2020 , 478, 992-1003	2.2	8

6	Reply. Clinical Gastroenterology and Hepatology, 2020 , 18, 3061-3062	6.9	1
5	Screening strategies for nonalcoholic fatty liver disease in type 2 diabetes: Insights from NHANES 2005-2016. <i>Diabetes Research and Clinical Practice</i> , 2020 , 167, 108358	7.4	14
4	Visit-to-visit blood pressure variability in patients with type 2 diabetes with and without previous history of cardiovascular disease. <i>Journal of Hypertension</i> , 2020 , 38, 1737-1744	1.9	6
3	Risk stratification tools for heart failure in the diabetes clinic. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2020 , 30, 1070-1079	4.5	5
2	Screening of SLC2A1 in a large cohort of patients suspected for Glut1 deficiency syndrome: identification of novel variants and associated phenotypes. <i>Journal of Neurology</i> , 2019 , 266, 1439-1448	5.5	11
1	Impact of diabetes on COVID-19-related in-hospital mortality: a retrospective study from Northern Italy	1	3