

Andreas Scorilas

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8343178/publications.pdf>

Version: 2024-02-01

396
papers

13,914
citations

18436

62
h-index

51492

86
g-index

400
all docs

400
docs citations

400
times ranked

13753
citing authors

#	ARTICLE	IF	CITATIONS
1	BCL2 Family of Apoptosis-Related Genes: Functions and Clinical Implications in Cancer. <i>Critical Reviews in Clinical Laboratory Sciences</i> , 2006, 43, 1-67.	2.7	214
2	Overexpression of matrix-metalloproteinase-9 in human breast cancer: a potential favourable indicator in node-negative patients. <i>British Journal of Cancer</i> , 2001, 84, 1488-1496.	2.9	210
3	Human Kallikrein 6 (hK6): A New Potential Serum Biomarker for Diagnosis and Prognosis of Ovarian Carcinoma. <i>Journal of Clinical Oncology</i> , 2003, 21, 1035-1043.	0.8	188
4	Genomic Organization of the Human Kallikrein Gene Family on Chromosome 19q13.3â€“q13.4. <i>Biochemical and Biophysical Research Communications</i> , 2000, 276, 125-133.	1.0	183
5	The Role of BCL2 Family of Apoptosis Regulator Proteins in Acute and Chronic Leukemias. <i>Advances in Hematology</i> , 2012, 2012, 1-15.	0.6	183
6	Evaluation of PD-L1 Expression and Associated Tumor-Infiltrating Lymphocytes in Laryngeal Squamous Cell Carcinoma. <i>Clinical Cancer Research</i> , 2016, 22, 704-713.	3.2	173
7	A new tumor suppressor role for the Notch pathway in bladder cancer. <i>Nature Medicine</i> , 2014, 20, 1199-1205.	15.2	160
8	The miR-17-92 Cluster is Over Expressed in and Has an Oncogenic Effect on Renal Cell Carcinoma. <i>Journal of Urology</i> , 2010, 183, 743-751.	0.2	149
9	Lactate Dehydrogenase A is a potential prognostic marker in clear cell renal cell carcinoma. <i>Molecular Cancer</i> , 2014, 13, 101.	7.9	141
10	Structure and biological properties of the copper(II) complex with the quinolone antibacterial drug N-propyl-norfloxacin and 2,2â€²-bipyridine. <i>Journal of Inorganic Biochemistry</i> , 2007, 101, 64-73.	1.5	137
11	Parallel overexpression of seven kallikrein genes in ovarian cancer. <i>Cancer Research</i> , 2003, 63, 2223-7.	0.4	126
12	A comprehensive nomenclature for serine proteases with homology to tissue kallikreins. <i>Biological Chemistry</i> , 2006, 387, 637-41.	1.2	123
13	The serum concentration of human kallikrein 10 represents a novel biomarker for ovarian cancer diagnosis and prognosis. <i>Cancer Research</i> , 2003, 63, 807-11.	0.4	123
14	Hepsin is Highly Over Expressed in and a New Candidate for a Prognostic Indicator in Prostate Cancer. <i>Journal of Urology</i> , 2004, 171, 187-191.	0.2	117
15	Human kallikrein gene 5 (KLK5) expression is an indicator of poor prognosis in ovarian cancer. <i>British Journal of Cancer</i> , 2001, 84, 643-650.	2.9	116
16	SARS-CoV-2 wastewater surveillance data can predict hospitalizations and ICU admissions. <i>Science of the Total Environment</i> , 2022, 804, 150151.	3.9	116
17	The PRMT1 gene expression pattern in colon cancer. <i>British Journal of Cancer</i> , 2008, 99, 2094-2099.	2.9	114
18	Human kallikrein 5: a potential novel serum biomarker for breast and ovarian cancer. <i>Cancer Research</i> , 2003, 63, 3958-65.	0.4	109

#	ARTICLE	IF	CITATIONS
19	The Clinical Utility of miR-21 as a Diagnostic and Prognostic Marker for Renal Cell Carcinoma. <i>Journal of Molecular Diagnostics</i> , 2012, 14, 385-392.	1.2	106
20	Adverse effects of COVID-19 mRNA vaccines: the spike hypothesis. <i>Trends in Molecular Medicine</i> , 2022, 28, 542-554.	3.5	104
21	The Combination of Human Glandular Kallikrein and Free Prostate-specific Antigen (PSA) Enhances Discrimination Between Prostate Cancer and Benign Prostatic Hyperplasia in Patients with Moderately Increased Total PSA. <i>Clinical Chemistry</i> , 1999, 45, 1960-1966.	1.5	103
22	Molecular Cloning of the Human Kallikrein 15 Gene (KLK15). <i>Journal of Biological Chemistry</i> , 2001, 276, 53-61.	1.6	103
23	Serum Human Glandular Kallikrein-2 Protease Levels Predict the Presence of Prostate Cancer Among Men With Elevated Prostate-Specific Antigen. <i>Journal of Clinical Oncology</i> , 2000, 18, 1036-1036.	0.8	99
24	Decreased concentrations of prostate-specific antigen and human glandular kallikrein 2 in malignant versus nonmalignant prostatic tissue. <i>Urology</i> , 2000, 56, 527-532.	0.5	99
25	<i>Phosphatidylinositol 3-kinase Catalytic Subunit 1</i> Gene Amplification Contributes to the Pathogenesis of Mantle Cell Lymphoma. <i>Clinical Cancer Research</i> , 2009, 15, 5724-5732.	3.2	99
26	The human KLK8 (neuropsin/ovasin) gene: identification of two novel splice variants and its prognostic value in ovarian cancer. <i>Clinical Cancer Research</i> , 2001, 7, 806-11.	3.2	98
27	Higher human kallikrein gene 4 (KLK4) expression indicates poor prognosis of ovarian cancer patients. <i>Clinical Cancer Research</i> , 2001, 7, 2380-6.	3.2	95
28	Human kallikrein 10: a novel tumor marker for ovarian carcinoma?. <i>Clinica Chimica Acta</i> , 2001, 306, 111-118.	0.5	94
29	The lysine-specific methyltransferase <i>KMT2C</i> / <i>MLL3</i> regulates <i>DNA</i> repair components in cancer. <i>EMBO Reports</i> , 2019, 20, .	2.0	93
30	The loss of the tumour-suppressor miR-145 results in the shorter disease-free survival of prostate cancer patients. <i>British Journal of Cancer</i> , 2013, 108, 2573-2581.	2.9	90
31	Non-coding RNAs: the riddle of the transcriptome and their perspectives in cancer. <i>Annals of Translational Medicine</i> , 2018, 6, 241-241.	0.7	90
32	Molecular Cloning, Physical Mapping, and Expression Analysis of a Novel Gene, <i>BCL2L12</i> , Encoding a Proline-Rich Protein with a Highly Conserved BH2 Domain of the Bcl-2 Family. <i>Genomics</i> , 2001, 72, 217-221.	1.3	89
33	Analytical methodologies for the detection of SARS-CoV-2 in wastewater: Protocols and future perspectives. <i>TrAC - Trends in Analytical Chemistry</i> , 2021, 134, 116125.	5.8	88
34	The KLK7 (PRSS6) gene, encoding for the stratum corneum chymotryptic enzyme is a new member of the human kallikrein gene family "genomic characterization, mapping, tissue expression and hormonal regulation. <i>Gene</i> , 2000, 254, 119-128.	1.0	87
35	Apoptosis-related BCL2-family Members: Key Players in Chemotherapy. <i>Anti-Cancer Agents in Medicinal Chemistry</i> , 2014, 14, 353-374.	0.9	85
36	Quantitative analysis of macrophage inhibitory cytokine-1 (MIC-1) gene expression in human prostatic tissues. <i>British Journal of Cancer</i> , 2003, 88, 1101-1104.	2.9	84

#	ARTICLE	IF	CITATIONS
37	Human Kallikrein Gene 5 (KLK5) Expression by Quantitative PCR: An Independent Indicator of Poor Prognosis in Breast Cancer. <i>Clinical Chemistry</i> , 2002, 48, 1241-1250.	1.5	82
38	The emergence of drug resistance to targeted cancer therapies: Clinical evidence. <i>Drug Resistance Updates</i> , 2019, 47, 100646.	6.5	81
39	Kallikrein-related peptidases in prostate, breast, and ovarian cancers: from pathobiology to clinical relevance. <i>Biological Chemistry</i> , 2012, 393, 301-317.	1.2	79
40	Differential Protein Expressions in Renal Cell Carcinoma: New Biomarker Discovery by Mass Spectrometry. <i>Journal of Proteome Research</i> , 2009, 8, 3797-3807.	1.8	78
41	Prognostic value of human kallikrein 10 expression in epithelial ovarian carcinoma. <i>Clinical Cancer Research</i> , 2001, 7, 2372-9.	3.2	78
42	Kallikrein-related peptidase genes as promising biomarkers for prognosis and monitoring of human malignancies. <i>Biological Chemistry</i> , 2010, 391, 505-511.	1.2	75
43	Kallikrein-related peptidases (KLKs): a gene family of novel cancer biomarkers. <i>Clinical Chemistry and Laboratory Medicine</i> , 2012, 50, 1877-1891.	1.4	74
44	RAS/PI3K Crosstalk and Cetuximab Resistance in Head and Neck Squamous Cell Carcinoma. <i>Clinical Cancer Research</i> , 2014, 20, 2933-2946.	3.2	74
45	Human Kallikrein 13 Protein in Ovarian Cancer Cytosols: A New Favorable Prognostic Marker. <i>Journal of Clinical Oncology</i> , 2004, 22, 678-685.	0.8	73
46	Quantitative expression of the human kallikrein gene 9 (KLK9) in ovarian cancer: a new independent and favorable prognostic marker. <i>Cancer Research</i> , 2001, 61, 7811-8.	0.4	72
47	Immunofluorometric quantitation and histochemical localisation of kallikrein 6 protein in ovarian cancer tissue: a new independent unfavourable prognostic biomarker. <i>British Journal of Cancer</i> , 2002, 87, 763-771.	2.9	71
48	Human kallikrein 8, a novel biomarker for ovarian carcinoma. <i>Cancer Research</i> , 2003, 63, 2771-4.	0.4	71
49	Effects of Long-term Androgen Administration on Breast Tissue of Female-to-Male Transsexuals. <i>Journal of Histochemistry and Cytochemistry</i> , 2006, 54, 905-910.	1.3	70
50	Streptavidin-Polyvinylamine Conjugates Labeled with a Europium Chelate: Applications in Immunoassay, Immunohistochemistry, and Microarrays. <i>Clinical Chemistry</i> , 2000, 46, 1450-1455.	1.5	69
51	Prognostic Value of the Human Kallikrein Gene 15 Expression in Ovarian Cancer. <i>Journal of Clinical Oncology</i> , 2003, 21, 3119-3126.	0.8	69
52	A Multiparametric Panel for Ovarian Cancer Diagnosis, Prognosis, and Response to Chemotherapy. <i>Clinical Cancer Research</i> , 2007, 13, 6984-6992.	3.2	69
53	Prognostic value of quantitatively assessed KLK7 expression in ovarian cancer. <i>Clinical Biochemistry</i> , 2003, 36, 135-143.	0.8	68
54	Cathepsin B and cathepsin D expression in the progression of colorectal adenoma to carcinoma. <i>Cancer Letters</i> , 2004, 205, 97-106.	3.2	68

#	ARTICLE	IF	CITATIONS
55	Low Expression of miR-126 Is a Prognostic Marker for Metastatic Clear Cell Renal Cell Carcinoma. <i>American Journal of Pathology</i> , 2015, 185, 693-703.	1.9	68
56	Detection of Human Kallikrein 4 in Healthy and Cancerous Prostatic Tissues by Immunofluorometry and Immunohistochemistry. <i>Clinical Chemistry</i> , 2002, 48, 1232-1240.	1.5	67
57	B7-H4 is over-expressed in early-stage ovarian cancer and is independent of CA125 expression. <i>Gynecologic Oncology</i> , 2007, 106, 334-341.	0.6	67
58	Clinical evaluation of PRMT1 gene expression in breast cancer. <i>Tumor Biology</i> , 2011, 32, 575-582.	0.8	67
59	Uncovering the clinical utility of miR-143, miR-145 and miR-224 for predicting the survival of bladder cancer patients following treatment. <i>Carcinogenesis</i> , 2015, 36, 528-537.	1.3	67
60	Prognostic role and implications of mutation status of tumor suppressor gene ARID1A in cancer: a systematic review and meta-analysis. <i>Oncotarget</i> , 2015, 6, 39088-39097.	0.8	67
61	Third-Generation Sequencing: The Spearhead towards the Radical Transformation of Modern Genomics. <i>Life</i> , 2022, 12, 30.	1.1	67
62	The Chromatin Remodeling Gene ARID1A Is a New Prognostic Marker in Clear Cell Renal Cell Carcinoma. <i>American Journal of Pathology</i> , 2013, 182, 1163-1170.	1.9	66
63	Genomic Organization, Physical Mapping, and Expression Analysis of the Human Protein Arginine Methyltransferase 1 Gene. <i>Biochemical and Biophysical Research Communications</i> , 2000, 278, 349-359.	1.0	65
64	Expression analysis of the human kallikrein 7 (KLK7) in breast tumors: a new potential biomarker for prognosis of breast carcinoma. <i>Thrombosis and Haemostasis</i> , 2004, 91, 180-186.	1.8	65
65	Downregulation and Prognostic Performance of MicroRNA 224 Expression in Prostate Cancer. <i>Clinical Chemistry</i> , 2013, 59, 261-269.	1.5	65
66	Down-regulation of the human kallikrein gene 5 (KLK5) in prostate cancer tissues. <i>Prostate</i> , 2002, 51, 126-132.	1.2	64
67	Genomic Organization, Mapping, Tissue Expression, and Hormonal Regulation of Trypsin-like Serine Protease (TLSP PRSS20), a New Member of the Human Kallikrein Gene Family. <i>Genomics</i> , 2000, 63, 88-96.	1.3	62
68	Steroid Hormone Regulation and Prognostic Value of the Human Kallikrein Gene 14 in Ovarian Cancer. <i>American Journal of Clinical Pathology</i> , 2003, 119, 346-355.	0.4	62
69	Human Kallikrein 8 Protein Is a Favorable Prognostic Marker in Ovarian Cancer. <i>Clinical Cancer Research</i> , 2006, 12, 1487-1493.	3.2	60
70	Prognostic value of the apoptosis related genes BCL2 and BCL2L12 in breast cancer. <i>Cancer Letters</i> , 2007, 247, 48-55.	3.2	60
71	The expression of the CEACAM19 gene, a novel member of the CEA family, is associated with breast cancer progression. <i>International Journal of Oncology</i> , 2013, 42, 1770-1777.	1.4	60
72	Insulin-Like Growth Factor I (IGF-I) and IGF-Binding Protein-3 in Benign Prostatic Hyperplasia and Prostate Cancer. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2001, 86, 694-699.	1.8	59

#	ARTICLE	IF	CITATIONS
73	Differential expression of the human kallikrein gene 14 (KLK14) in normal and cancerous prostatic tissues. <i>Prostate</i> , 2003, 56, 287-292.	1.2	59
74	Human kallikrein gene 13 (KLK13) expression by quantitative RT-PCR: an independent indicator of favourable prognosis in breast cancer. <i>British Journal of Cancer</i> , 2002, 86, 1457-1464.	2.9	58
75	Human Tissue Kallikreins: From Gene Structure to Function and Clinical Applications. <i>Advances in Clinical Chemistry</i> , 2005, 39, 11-79.	1.8	58
76	TNF-alpha expression and apoptosis-regulating proteins in oral lichen planus: a comparative immunohistochemical evaluation. <i>Journal of Oral Pathology and Medicine</i> , 2000, 29, 370-375.	1.4	56
77	Favorable prognostic value of tissue human kallikrein 11 (hK11) in patients with ovarian carcinoma. <i>International Journal of Cancer</i> , 2003, 106, 605-610.	2.3	56
78	The use of kallikrein-related peptidases as adjuvant prognostic markers in colorectal cancer. <i>British Journal of Cancer</i> , 2009, 100, 1659-1665.	2.9	55
79	Prognostic value and biological role of the kallikrein-related peptidases in human malignancies. <i>Future Oncology</i> , 2010, 6, 269-285.	1.1	55
80	miR-210 Is a Prognostic Marker in Clear Cell Renal Cell Carcinoma. <i>Journal of Molecular Diagnostics</i> , 2015, 17, 136-144.	1.2	55
81	Quantitative Analysis of Kallikrein 15 Gene Expression in Prostate Tissue. <i>Journal of Urology</i> , 2003, 169, 361-364.	0.2	53
82	Galectin-1 has potential prognostic significance and is implicated in clear cell renal cell carcinoma progression through the HIF/mTOR signaling axis. <i>British Journal of Cancer</i> , 2014, 110, 1250-1259.	2.9	52
83	Comparative kinetics of SARS-CoV-2 anti-spike protein RBD IgGs and neutralizing antibodies in convalescent and naïve recipients of the BNT162b2 mRNA vaccine versus COVID-19 patients. <i>BMC Medicine</i> , 2021, 19, 208.	2.3	52
84	The role of cordycepin in cancer treatment via induction or inhibition of apoptosis: implication of polyadenylation in a cell type specific manner. <i>Cancer Chemotherapy and Pharmacology</i> , 2007, 61, 251-265.	1.1	50
85	Quantitative expression analysis and prognostic significance of L-DOPA decarboxylase in colorectal adenocarcinoma. <i>British Journal of Cancer</i> , 2010, 102, 1384-1390.	2.9	50
86	Quantitative analysis of the mRNA expression levels of BCL2 and BAX genes in human osteoarthritis and normal articular cartilage: An investigation into their differential expression. <i>Molecular Medicine Reports</i> , 2015, 12, 4514-4521.	1.1	50
87	MicroRNA-194 is a Marker for Good Prognosis in Clear Cell Renal Cell Carcinoma. <i>Cancer Medicine</i> , 2016, 5, 656-664.	1.3	50
88	Differential expression of Kallikrein gene 5 in cancerous and normal testicular tissues. <i>Urology</i> , 2002, 60, 714-718.	0.5	49
89	Decreased concentration of human kallikrein 6 in brain extracts of Alzheimer's disease patients. <i>Clinical Biochemistry</i> , 2002, 35, 225-231.	0.8	49
90	Microvascular density as an independent predictor of clinical outcome in renal cell carcinoma: an automated image analysis study. <i>Laboratory Investigation</i> , 2012, 92, 46-56.	1.7	48

#	ARTICLE	IF	CITATIONS
91	Clinical significance of kallikrein-related peptidase (KLK10) mRNA expression in colorectal cancer. <i>Clinical Biochemistry</i> , 2013, 46, 1453-1461.	0.8	48
92	The oncomiR miR-197 is a novel prognostic indicator for non-small cell lung cancer patients. <i>British Journal of Cancer</i> , 2015, 112, 1527-1535.	2.9	48
93	JQ1 inhibits tumour growth in combination with cisplatin and suppresses JAK/STAT signalling pathway in ovarian cancer. <i>European Journal of Cancer</i> , 2020, 126, 125-135.	1.3	48
94	Human kallikrein 11: an indicator of favorable prognosis in ovarian cancer patients. <i>Clinical Biochemistry</i> , 2004, 37, 823-829.	0.8	47
95	Revisiting Histone Deacetylases in Human Tumorigenesis: The Paradigm of Urothelial Bladder Cancer. <i>International Journal of Molecular Sciences</i> , 2019, 20, 1291.	1.8	47
96	Expression of Gelatinase-A (MMP-2) in Human Colon Cancer and Normal Colon Mucosa. <i>Tumor Biology</i> , 2001, 22, 383-389.	0.8	46
97	The Prognostic Value of the Human Kallikrein Gene 9 (KLK9) in Breast Cancer. <i>Breast Cancer Research and Treatment</i> , 2003, 78, 149-158.	1.1	46
98	Immunofluorometric Quantification of Human Kallikrein 5 Expression in Ovarian Cancer Cytosols and Its Association with Unfavorable Patient Prognosis. <i>Tumor Biology</i> , 2003, 24, 299-309.	0.8	45
99	Polyadenylate polymerase modulations in human epithelioid cervix and breast cancer cell lines, treated with etoposide or cordycepin, follow cell cycle rather than apoptosis induction. <i>Biological Chemistry</i> , 2005, 386, 471-480.	1.2	44
100	Unfavorable Prognostic Value of Human Kallikrein 7 Quantified by ELISA in Ovarian Cancer Cytosols. <i>Clinical Chemistry</i> , 2006, 52, 1879-1886.	1.5	44
101	High miR-96 levels in colorectal adenocarcinoma predict poor prognosis, particularly in patients without distant metastasis at the time of initial diagnosis. <i>Tumor Biology</i> , 2016, 37, 11815-11824.	0.8	44
102	Cisplatin-Induced Apoptosis in HL-60 Human Promyelocytic Leukemia Cells. <i>Annals of the New York Academy of Sciences</i> , 2003, 1010, 153-158.	1.8	43
103	Altered kallikrein 7 and 10 concentrations in cerebrospinal fluid of patients with Alzheimer's disease and frontotemporal dementia. <i>Clinical Biochemistry</i> , 2004, 37, 230-237.	0.8	43
104	Transcriptional upregulation of human tissue kallikrein 6 in ovarian cancer: clinical and mechanistic aspects. <i>British Journal of Cancer</i> , 2007, 96, 362-372.	2.9	43
105	The androgen-regulated gene human kallikrein 15 (KLK15) is an independent and favourable prognostic marker for breast cancer. <i>British Journal of Cancer</i> , 2002, 87, 1294-1300.	2.9	42
106	Determination of Cathepsin B Expression May Offer Additional Prognostic Information for Ovarian Cancer Patients. <i>Biological Chemistry</i> , 2002, 383, 1297-303.	1.2	42
107	Alterations in mRNA Expression of Apoptosis-Related Genes BCL2, BAX, FAS, Caspase-3, and the Novel Member BCL2L12 after Treatment of Human Leukemic Cell Line HL60 with the Antineoplastic Agent Etoposide. <i>Annals of the New York Academy of Sciences</i> , 2006, 1090, 89-97.	1.8	42
108	Computational approaches in cancer multidrug resistance research: Identification of potential biomarkers, drug targets and drug-target interactions. <i>Drug Resistance Updates</i> , 2020, 48, 100662.	6.5	42

#	ARTICLE	IF	CITATIONS
109	Circular RNAs: A New Piece in the Colorectal Cancer Puzzle. <i>Cancers</i> , 2020, 12, 2464.	1.7	42
110	Circular RNAs: Emerging Regulators of the Major Signaling Pathways Involved in Cancer Progression. <i>Cancers</i> , 2021, 13, 2744.	1.7	42
111	The usefulness of serum human kallikrein 11 for discriminating between prostate cancer and benign prostatic hyperplasia. <i>Cancer Research</i> , 2003, 63, 6543-6.	0.4	42
112	Quantitative analysis of hippostasin/KLK11 gene expression in cancerous and noncancerous prostatic tissues. <i>Urology</i> , 2003, 61, 1042-1046.	0.5	41
113	Loss of GAS5 tumour suppressor lncRNA: an independent molecular cancer biomarker for short-term relapse and progression in bladder cancer patients. <i>British Journal of Cancer</i> , 2018, 119, 1477-1486.	2.9	41
114	miRNA and long non-coding RNA: molecular function and clinical value in breast and ovarian cancers. <i>Expert Review of Molecular Diagnostics</i> , 2018, 18, 963-979.	1.5	41
115	Downregulation of the neonatal Fc receptor expression in non-small cell lung cancer tissue is associated with a poor prognosis. <i>Oncotarget</i> , 2016, 7, 54415-54429.	0.8	41
116	Quantitative analysis of human kallikrein gene 14 expression in breast tumours indicates association with poor prognosis. <i>British Journal of Cancer</i> , 2002, 87, 1287-1293.	2.9	40
117	Serum human glandular kallikrein (hK2) and insulin-like growth factor 1 (IGF-1) improve the discrimination between prostate cancer and benign prostatic hyperplasia in combination with total and %free PSA. <i>Prostate</i> , 2003, 54, 220-229.	1.2	40
118	Expression analysis and prognostic significance of human kallikrein 11 in prostate cancer. <i>Clinica Chimica Acta</i> , 2005, 357, 190-195.	0.5	40
119	Quantitative expression analysis and prognostic significance of the novel apoptosis-related gene <i>BCL2L12</i> in colon cancer. <i>Biological Chemistry</i> , 2008, 389, 1467-1475.	1.2	40
120	miR-15a-5p, A Novel Prognostic Biomarker, Predicting Recurrent Colorectal Adenocarcinoma. <i>Molecular Diagnosis and Therapy</i> , 2017, 21, 453-464.	1.6	40
121	Evolution of the Plasma and Tissue Kallikreins, and Their Alternative Splicing Isoforms. <i>PLoS ONE</i> , 2013, 8, e68074.	1.1	40
122	BCL2L12 is a Novel Biomarker for the Prediction of Short-Term Relapse in Nasopharyngeal Carcinoma. <i>Molecular Medicine</i> , 2011, 17, 163-171.	1.9	39
123	The Novel Member of the <i>BCL2</i> Gene Family, <i>BCL2L12</i> , Is Substantially Elevated in Chronic Lymphocytic Leukemia Patients, Supporting Its Value As a Significant Biomarker. <i>Oncologist</i> , 2011, 16, 1280-1291.	1.9	39
124	Enhanced miR-182 transcription is a predictor of poor overall survival in colorectal adenocarcinoma patients. <i>Clinical Chemistry and Laboratory Medicine</i> , 2014, 52, 1217-27.	1.4	39
125	Determination of c-myc amplification and overexpression in breast cancer patients: evaluation of its prognostic value against c-erbB-2, cathepsin-D and clinicopathological characteristics using univariate and multivariate analysis. <i>British Journal of Cancer</i> , 1999, 81, 1385-1391.	2.9	38
126	Breast Cancer Cells Response to the Antineoplastic Agents Cisplatin, Carboplatin, and Doxorubicin at the mRNA Expression Levels of Distinct Apoptosis-Related Genes, Including the New Member, BCL2L12. <i>Annals of the New York Academy of Sciences</i> , 2007, 1095, 35-44.	1.8	38

#	ARTICLE	IF	CITATIONS
127	Emerging clinical importance of the cancer biomarkers kallikrein-related peptidases (KLK) in female and male reproductive organ malignancies. <i>Radiology and Oncology</i> , 2013, 47, 319-329.	0.6	38
128	miR-224 overexpression is a strong and independent prognosticator of short-term relapse and poor overall survival in colorectal adenocarcinoma. <i>International Journal of Oncology</i> , 2015, 46, 849-859.	1.4	38
129	Expression of BCL2L12, a new member of apoptosis-related genes, in breast tumors. <i>Thrombosis and Haemostasis</i> , 2003, 89, 1081-1088.	1.8	37
130	Expression analysis and clinical utility of L-Dopa decarboxylase (DDC) in prostate cancer. <i>Clinical Biochemistry</i> , 2008, 41, 1140-1149.	0.8	37
131	Clinical significance of kallikrein-related peptidase 7 (KLK7) in colorectal cancer. <i>Thrombosis and Haemostasis</i> , 2009, 101, 741-747.	1.8	37
132	Quantitative expression analysis of the apoptosis-related genes BCL2, BAX and BCL2L12 in gastric adenocarcinoma cells following treatment with the anticancer drugs cisplatin, etoposide and taxol. <i>Tumor Biology</i> , 2012, 33, 865-875.	0.8	37
133	Kallikrein-related peptidases (KLKs) in gastrointestinal cancer: Mechanistic and clinical aspects. <i>Thrombosis and Haemostasis</i> , 2013, 110, 450-457.	1.8	37
134	Impact of expression differences of kallikrein-related peptidases and of uPA and PAI-1 between primary tumor and omentum metastasis in advanced ovarian cancer. <i>Annals of Oncology</i> , 2011, 22, 877-883.	0.6	36
135	Kallikrein-related peptidases (KLKs) as emerging therapeutic targets: focus on prostate cancer and skin pathologies. <i>Expert Opinion on Therapeutic Targets</i> , 2016, 20, 801-818.	1.5	36
136	The role of circular RNAs in therapy resistance of patients with solid tumors. <i>Personalized Medicine</i> , 2020, 17, 469-490.	0.8	35
137	Human Glandular Kallikrein in Breast Milk, Amniotic Fluid, and Breast Cyst Fluid. <i>Clinical Chemistry</i> , 1999, 45, 1774-1780.	1.5	34
138	Treatment of MCF-7 cells with taxol and etoposide induces distinct alterations in the expression of apoptosis-related genes BCL2, BCL2L12, BAX, CASPASE-9 and FAS. <i>Biological Chemistry</i> , 2006, 387, 1081-6.	1.2	34
139	Molecular Profile of the <i>BCL2</i> Family of the Apoptosis Related Genes in Breast Cancer Cells after Treatment with Cytotoxic/Cytostatic Drugs. <i>Connective Tissue Research</i> , 2008, 49, 261-264.	1.1	34
140	Molecular analysis and prognostic impact of the novel apoptotic gene BCL2L12 in gastric cancer. <i>Biochemical and Biophysical Research Communications</i> , 2010, 391, 214-218.	1.0	34
141	Kallikrein-related peptidase-6 (KLK6) mRNA expression is an independent prognostic tissue biomarker of poor disease-free and overall survival in colorectal adenocarcinoma. <i>Tumor Biology</i> , 2014, 35, 4673-4685.	0.8	34
142	mRNA expression analysis of a variety of apoptosis-related genes, including the novel gene of the BCL2-family, BCL2L12, in HL-60 leukemia cells after treatment with carboplatin and doxorubicin. <i>Biological Chemistry</i> , 2004, 385, 1099-103.	1.2	33
143	Upregulated miR-16 expression is an independent indicator of relapse and poor overall survival of colorectal adenocarcinoma patients. <i>Clinical Chemistry and Laboratory Medicine</i> , 2017, 55, 737-747.	1.4	33
144	miR-125b predicts childhood acute lymphoblastic leukaemia poor response to BFM chemotherapy treatment. <i>British Journal of Cancer</i> , 2017, 117, 801-812.	2.9	33

#	ARTICLE	IF	CITATIONS
145	Natural Alkaloids Intervening the Insulin Pathway: New Hopes for Anti-Diabetic Agents?. <i>Current Medicinal Chemistry</i> , 2019, 26, 5982-6015.	1.2	33
146	Nature Promises New Anticancer Agents: Interplay with the Apoptosis-related BCL2 Gene Family. <i>Anti-Cancer Agents in Medicinal Chemistry</i> , 2014, 14, 375-399.	0.9	33
147	Human kallikrein gene 5 (KLK5) expression by quantitative PCR: an independent indicator of poor prognosis in breast cancer. <i>Clinical Chemistry</i> , 2002, 48, 1241-50.	1.5	33
148	Polyvinylamine-streptavidin complexes labeled with a europium chelator: a universal detection reagent for solid-phase time resolved fluorometric applications. <i>Clinical Biochemistry</i> , 2000, 33, 345-350.	0.8	32
149	High BAX/BCL2 mRNA ratio predicts favorable prognosis in laryngeal squamous cell carcinoma, particularly in patients with negative lymph nodes at the time of diagnosis. <i>Clinical Biochemistry</i> , 2016, 49, 890-896.	0.8	32
150	MicroRNA-155-5p Overexpression in Peripheral Blood Mononuclear Cells of Chronic Lymphocytic Leukemia Patients Is a Novel, Independent Molecular Biomarker of Poor Prognosis. <i>Disease Markers</i> , 2017, 2017, 1-10.	0.6	32
151	Prostate-Specific Antigen and Human Glandular Kallikrein 2 Are Markedly Elevated in Urine of Patients with Polycystic Ovary Syndrome. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2001, 86, 1558-1561.	1.8	31
152	The effect of the polyadenylation inhibitor cordycepin on human Molt-4 and Daudi leukaemia and lymphoma cell lines. <i>Cancer Chemotherapy and Pharmacology</i> , 2008, 61, 703-711.	1.1	31
153	Elevated expression of miR-24-3p is a potentially adverse prognostic factor in colorectal adenocarcinoma. <i>Clinical Biochemistry</i> , 2017, 50, 285-292.	0.8	31
154	Seroprevalence of Antibodies against SARS-CoV-2 among the Personnel and Students of the National and Kapodistrian University of Athens, Greece: A Preliminary Report. <i>Life</i> , 2020, 10, 214.	1.1	31
155	tRNA-Derived Fragments (tRFs) in Bladder Cancer: Increased 5â€²-tRF-LysCTT Results in Disease Early Progression and Patientsâ€™ Poor Treatment Outcome. <i>Cancers</i> , 2020, 12, 3661.	1.7	31
156	MicroRNAs: Tiny Regulators of Gene Expression with Pivotal Roles in Normal B-Cell Development and B-Cell Chronic Lymphocytic Leukemia. <i>Cancers</i> , 2021, 13, 593.	1.7	31
157	Expression analysis of BCL2L12, a new member of apoptosis-related genes, in colon cancer. <i>Biological Chemistry</i> , 2004, 385, 779-83.	1.2	30
158	Expression analysis and clinical evaluation of kallikrein-related peptidase 10 (KLK10) in colorectal cancer. <i>Tumor Biology</i> , 2011, 32, 737-744.	0.8	30
159	Evaluation and prognostic significance of human tissue kallikrein-related peptidase 6 (KLK6) in colorectal cancer. <i>Pathology Research and Practice</i> , 2012, 208, 104-108.	1.0	30
160	Molecular cloning of novel alternatively spliced variants of BCL2L12, a new member of the BCL2 gene family, and their expression analysis in cancer cells. <i>Gene</i> , 2012, 505, 153-166.	1.0	30
161	Kallikrein-related peptidase 4 (KLK4) mRNA predicts short-term relapse in colorectal adenocarcinoma patients. <i>Cancer Letters</i> , 2013, 330, 106-112.	3.2	30
162	<i>KLK11</i> mRNA expression predicts poor disease-free and overall survival in colorectal adenocarcinoma patients. <i>Biomarkers in Medicine</i> , 2014, 8, 671-685.	0.6	30

#	ARTICLE	IF	CITATIONS
163	The combination of human glandular kallikrein and free prostate-specific antigen (PSA) enhances discrimination between prostate cancer and benign prostatic hyperplasia in patients with moderately increased total PSA. <i>Clinical Chemistry</i> , 1999, 45, 1960-6.	1.5	30
164	Immunohistochemical expression of Bcl2 is an independent predictor of time-to-biochemical failure in patients with clinically localized prostate cancer following radical prostatectomy. <i>Anticancer Research</i> , 2005, 25, 3123-33.	0.5	30
165	The role of human tissue kallikreins 7 and 8 in intracranial malignancies. <i>Biological Chemistry</i> , 2006, 387, 1607-1612.	1.2	29
166	Loss of miR-378 in prostate cancer, a common regulator of <i>KLK2</i> and <i>KLK4</i> , correlates with aggressive disease phenotype and predicts the short-term relapse of the patients. <i>Biological Chemistry</i> , 2014, 395, 1095-1104.	1.2	29
167	Prognostic and predictive biomarkers in prostate cancer. <i>Expert Review of Molecular Diagnostics</i> , 2015, 15, 1567-1576.	1.5	29
168	miR-10b is a prognostic marker in clear cell renal cell carcinoma. <i>Journal of Clinical Pathology</i> , 2017, 70, 854-859.	1.0	29
169	Elevated miR-20b-5p expression in peripheral blood mononuclear cells: A novel, independent molecular biomarker of favorable prognosis in chronic lymphocytic leukemia. <i>Leukemia Research</i> , 2018, 70, 1-7.	0.4	29
170	Primary Tumor Levels of Human Tissue Kallikreins Affect Surgical Success and Survival in Ovarian Cancer Patients. <i>Clinical Cancer Research</i> , 2007, 13, 1742-1748.	3.2	28
171	Kallikrein-related peptidase 4 gene (<i>KLK4</i>) in prostate tumors: Quantitative expression analysis and evaluation of its clinical significance. <i>Prostate</i> , 2011, 71, 1780-1789.	1.2	28
172	Parallel overexpression and clinical significance of kallikrein-related peptidases 7 and 14 (KLK7 & 14) in prostate cancer. <i>Journal of Cellular Biochemistry</i> , 2017, 142, 100-108.	1.8	28
173	High microRNA-28-5p expression in colorectal adenocarcinoma predicts short-term relapse of node-negative patients and poor overall survival of patients with non-metastatic disease. <i>Clinical Chemistry and Laboratory Medicine</i> , 2018, 56, 990-1000.	1.4	28
174	Circulating exosomal miRNAs: clinical significance in human cancers. <i>Expert Review of Molecular Diagnostics</i> , 2019, 19, 979-995.	1.5	28
175	Identification of a novel tRNA-derived RNA fragment exhibiting high prognostic potential in chronic lymphocytic leukemia. <i>Hematological Oncology</i> , 2019, 37, 498-504.	0.8	28
176	Correlation of androgen receptor status, neuroendocrine differentiation and angiogenesis with time-to-biochemical failure after radical prostatectomy in clinically localized prostate cancer. <i>Anticancer Research</i> , 2007, 27, 3651-60.	0.5	28
177	Serum and Urinary Prostate-specific Antigen and Urinary Human Glandular Kallikrein Concentrations Are Significantly Increased after Testosterone Administration in Female-to-Male Transsexuals. <i>Clinical Chemistry</i> , 2000, 46, 859-862.	1.5	27
178	Kallikrein-related peptidase 13 (KLK13) gene expressional status contributes significantly in the prognosis of primary gastric carcinomas. <i>Clinical Biochemistry</i> , 2010, 43, 1205-1211.	0.8	27
179	Assessment of the prognostic significance of endoglin (CD105) in clear cell renal cell carcinoma using automated image analysis. <i>Human Pathology</i> , 2012, 43, 1037-1043.	1.1	27
180	Long Noncoding RNAs in Digestive System Malignancies: A Novel Class of Cancer Biomarkers and Therapeutic Targets?. <i>Gastroenterology Research and Practice</i> , 2015, 2015, 1-18.	0.7	27

#	ARTICLE	IF	CITATIONS
181	KLKB1 mRNA overexpression: A novel molecular biomarker for the diagnosis of chronic lymphocytic leukemia. <i>Clinical Biochemistry</i> , 2015, 48, 849-854.	0.8	27
182	Immunoenzymatically determined pepsinogen C concentration in breast tumor cytosols: an independent favorable prognostic factor in node-positive patients. <i>Clinical Cancer Research</i> , 1999, 5, 1778-85.	3.2	27
183	Enhanced Antileukemic Activity of the Novel Complex 2,5-Dihydroxybenzoate Molybdenum(VI) against 2,5-Dihydroxybenzoate, Polyoxometalate of Mo(VI), and Tetraphenylphosphonium in the Human HL-60 and K562 Leukemic Cell Lines. <i>Journal of Medicinal Chemistry</i> , 2007, 50, 1316-1321.	2.9	26
184	Expression analysis and study of KLK4 in benign and malignant breast tumours. <i>Thrombosis and Haemostasis</i> , 2009, 101, 381-387.	1.8	26
185	Quantitative analysis of BCL2 mRNA expression in nasopharyngeal carcinoma: an unfavorable and independent prognostic factor. <i>Tumor Biology</i> , 2010, 31, 391-399.	0.8	26
186	A Comprehensive Phylogenetic and Structural Analysis of the Carcinoembryonic Antigen (CEA) Gene Family. <i>Genome Biology and Evolution</i> , 2014, 6, 1314-1326.	1.1	26
187	Identification of novel alternative splice variants of the BCL2L12 gene in human cancer cells using next-generation sequencing methodology. <i>Cancer Letters</i> , 2016, 373, 119-129.	3.2	26
188	mRNA overexpression of the hypoxia inducible factor 1 alpha subunit gene (HIF1A): An independent predictor of poor overall survival in chronic lymphocytic leukemia. <i>Leukemia Research</i> , 2017, 53, 65-73.	0.4	26
189	miR-221/222 cluster expression improves clinical stratification of non-muscle invasive bladder cancer (TaT1) patients' risk for short-term relapse and progression. <i>Genes Chromosomes and Cancer</i> , 2018, 57, 150-161.	1.5	26
190	Polyadenylate Polymerase (PAP) and 3' End pre-mRNA Processing: Function, Assays, and Association with Disease. <i>Critical Reviews in Clinical Laboratory Sciences</i> , 2002, 39, 193-224.	2.7	25
191	Kallikreins as Markers of Disseminated Tumour Cells in Ovarian Cancer – A Pilot Study. <i>Tumor Biology</i> , 2006, 27, 104-114.	0.8	25
192	Human tissue kallikrein 7, a novel biomarker for advanced ovarian carcinoma using a novel in situ quantitative method of protein expression. <i>Annals of Oncology</i> , 2008, 19, 1271-1277.	0.6	25
193	Pancreatic duct guidewire placement for biliary cannulation in a single-session therapeutic ERCP. <i>World Journal of Gastroenterology</i> , 2011, 17, 1989.	1.4	25
194	Quantitative expression analysis of the apoptosis-related gene, <i>BCL2L12</i> , in head and neck squamous cell carcinoma. <i>Journal of Oral Pathology and Medicine</i> , 2013, 42, 154-161.	1.4	25
195	Targeting kallikrein-related peptidases in prostate cancer. <i>Expert Opinion on Therapeutic Targets</i> , 2014, 18, 365-383.	1.5	25
196	Profilin-1 expression is associated with high grade and stage and decreased disease-free survival in renal cell carcinoma. <i>Human Pathology</i> , 2015, 46, 673-680.	1.1	25
197	Copper(II) Inverse-[9-Metallacrown-3] Compounds Accommodating Nitrate or Diclofenac Ligands: Structure, Magnetism, and Biological Activity. <i>European Journal of Inorganic Chemistry</i> , 2016, 2016, 219-231.	1.0	25
198	miR-34a overexpression predicts poor prognostic outcome in colorectal adenocarcinoma, independently of clinicopathological factors with established prognostic value. <i>Clinical Biochemistry</i> , 2017, 50, 918-924.	0.8	25

#	ARTICLE	IF	CITATIONS
199	The miR-200 family as prognostic markers in clear cell renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2019, 37, 955-963.	0.8	25
200	Polyadenylate polymerase enzymatic activity in mammary tumor cytosols: A new independent prognostic marker in primary breast cancer. <i>Cancer Research</i> , 2000, 60, 5427-33.	0.4	25
201	tRNAGlyGCC-Derived Internal Fragment (i-tRF-GlyGCC) in Ovarian Cancer Treatment Outcome and Progression. <i>Cancers</i> , 2022, 14, 24.	1.7	25
202	Prognostic value of kallikrein-related peptidase 6 protein expression levels in advanced ovarian cancer evaluated by automated quantitative analysis (AQUA). <i>Cancer Science</i> , 2008, 99, 2224-2229.	1.7	24
203	Trastuzumab plus Paclitaxel or Docetaxel in HER-2 "Negative/HER-2 ECD" Positive Anthracycline- and Taxane-Refractory Advanced Breast Cancer. <i>Oncologist</i> , 2008, 13, 361-369.	1.9	24
204	Diagnostic and prognostic significance of human kallikrein 11 (KLK11) mRNA expression levels in patients with laryngeal cancer. <i>Clinical Biochemistry</i> , 2012, 45, 623-630.	0.8	24
205	Quantitative expression analysis and prognostic significance of the BCL2-associated Xgene in nasopharyngeal carcinoma: a retrospective cohort study. <i>BMC Cancer</i> , 2013, 13, 293.	1.1	24
206	Effect of doxorubicin, oxaliplatin, and methotrexate administration on the transcriptional activity of BCL-2 family gene members in stomach cancer cells. <i>Cancer Biology and Therapy</i> , 2013, 14, 587-596.	1.5	24
207	Kallikrein-related peptidase 6 (KLK6) expression in the progression of colon adenoma to carcinoma. <i>Biological Chemistry</i> , 2014, 395, 1105-1117.	1.2	24
208	Progression of mouse skin carcinogenesis is associated with the orchestrated deregulation of miR-200 family members, miR-205 and their common targets. <i>Molecular Carcinogenesis</i> , 2016, 55, 1229-1242.	1.3	24
209	Identification of a novel, internal tRNA-derived RNA fragment as a new prognostic and screening biomarker in chronic lymphocytic leukemia, using an innovative quantitative real-time PCR assay. <i>Leukemia Research</i> , 2019, 87, 106234.	0.4	24
210	Next generation sequencing targeted gene panel in Greek MODY patients increases diagnostic accuracy. <i>Pediatric Diabetes</i> , 2020, 21, 28-39.	1.2	24
211	Identification of Two Novel Circular RNAs Deriving from BCL2L12 and Investigation of Their Potential Value as a Molecular Signature in Colorectal Cancer. <i>International Journal of Molecular Sciences</i> , 2020, 21, 8867.	1.8	24
212	Predictive value of c-erbB-2 and cathepsin-D for Greek breast cancer patients using univariate and multivariate analysis. <i>Clinical Cancer Research</i> , 1999, 5, 815-21.	3.2	24
213	Molecular profile of breast versus ovarian cancer cells in response to treatment with the anticancer drugs cisplatin, carboplatin, doxorubicin, etoposide and taxol. <i>Biological Chemistry</i> , 2008, 389, 1427-1434.	1.2	23
214	Association between kallikrein-related peptidases (KLKs) and macroscopic indicators of semen analysis: their relation to sperm motility. <i>Biological Chemistry</i> , 2009, 390, 921-929.	1.2	23
215	Cathepsin B protein levels in endometrial cancer: Potential value as a tumour biomarker. <i>Gynecologic Oncology</i> , 2009, 112, 531-536.	0.6	23
216	High clusterin (CLU) mRNA expression levels in tumors of colorectal cancer patients predict a poor prognostic outcome. <i>Clinical Biochemistry</i> , 2020, 75, 62-69.	0.8	23

#	ARTICLE	IF	CITATIONS
217	A novel, mitochondrial, internal tRNA-derived RNA fragment possesses clinical utility as a molecular prognostic biomarker in chronic lymphocytic leukemia. <i>Clinical Biochemistry</i> , 2020, 85, 20-26.	0.8	23
218	Significance of Urokinase-Type Plasminogen Activator and Plasminogen Activator Inhibitor-1 (PAI-1) Expression in Human Colorectal Carcinomas. <i>Tumor Biology</i> , 2002, 23, 170-178.	0.8	22
219	Molecular characterization of a new gene, CEAL1, encoding for a carcinoembryonic antigen-like protein with a highly conserved domain of eukaryotic translation initiation factors. <i>Gene</i> , 2003, 310, 79-89.	1.0	22
220	Down-regulation of kallikrein-related peptidase 5 (KLK5) expression in breast cancer patients: a biomarker for the differential diagnosis of breast lesions. <i>Clinical Proteomics</i> , 2011, 8, 5.	1.1	22
221	The single nucleotide polymorphism g.1548A >G (K469E) of the ICAM-1 gene is associated with worse prognosis in non-small cell lung cancer. <i>Tumor Biology</i> , 2012, 33, 1429-1436.	0.8	22
222	Cloning of a gene (SR-A1), encoding for a new member of the human Ser/Arg-rich family of pre-mRNA splicing factors: overexpression in aggressive ovarian cancer. <i>British Journal of Cancer</i> , 2001, 85, 190-198.	2.9	21
223	Quantitative Analysis of Human Kallikrein 5 (KLK5) Expression in Prostate Needle Biopsies: An Independent Cancer Biomarker. <i>Clinical Chemistry</i> , 2009, 55, 904-913.	1.5	21
224	Human kallikrein-related peptidase 12 (KLK12) splice variants expression in breast cancer and their clinical impact. <i>Tumor Biology</i> , 2012, 33, 1075-1084.	0.8	21
225	Blood-based analysis of type-2 diabetes mellitus susceptibility genes identifies specific transcript variants with deregulated expression and association with disease risk. <i>Scientific Reports</i> , 2019, 9, 1512.	1.6	21
226	Clinical significance of kallikrein-related peptidase 7 (KLK7) in colorectal cancer. <i>Thrombosis and Haemostasis</i> , 2009, 101, 741-7.	1.8	21
227	L-Dopa decarboxylase (<i>DDC</i>) gene expression is related to outcome in patients with prostate cancer. <i>BJU International</i> , 2012, 110, E267-73.	1.3	20
228	Clinical utility of microRNAs in renal cell carcinoma: current evidence and future perspectives. <i>Expert Review of Molecular Diagnostics</i> , 2018, 18, 981-991.	1.5	20
229	Free/Total PSA (F/T ratio) kinetics in patients with clinically localized prostate cancer undergoing radical prostatectomy. <i>Clinica Chimica Acta</i> , 2005, 357, 196-201.	0.5	19
230	Topotecan and methotrexate alter expression of the apoptosis-related genes BCL2, FAS and BCL2L12 in leukemic HL-60 cells. <i>Biological Chemistry</i> , 2006, 387, 1629-33.	1.2	19
231	mRNA expression analysis of human kallikrein 11 (KLK11) may be useful in the discrimination of benign prostatic hyperplasia from prostate cancer after needle prostate biopsy. <i>Biological Chemistry</i> , 2006, 387, 789-793.	1.2	19
232	Treatment of PC3 prostate cancer cells with mitoxantrone, etoposide, doxorubicin and carboplatin induces distinct alterations in the expression of kallikreins 5 and 11. <i>Thrombosis and Haemostasis</i> , 2009, 101, 373-380.	1.8	19
233	Comparative study of balloon and metal olive dilators for endoscopic management of benign anastomotic rectal strictures: clinical and cost-effectiveness outcomes. <i>Surgical Endoscopy and Other Interventional Techniques</i> , 2011, 25, 756-763.	1.3	19
234	L-DOPA decarboxylase mRNA expression is associated with tumor stage and size in head and neck squamous cell carcinoma: a retrospective cohort study. <i>BMC Cancer</i> , 2012, 12, 484.	1.1	19

#	ARTICLE	IF	CITATIONS
235	Cytotoxic activity of sunitinib and everolimus in Caki-1 renal cancer cells is accompanied by modulations in the expression of apoptosis-related microRNA clusters and BCL2 family genes. <i>Biomedicine and Pharmacotherapy</i> , 2015, 70, 33-40.	2.5	19
236	Expression and prognostic significance of kallikrein-related peptidase 8 protein levels in advanced ovarian cancer by using automated quantitative analysis. <i>Thrombosis and Haemostasis</i> , 2009, 101, 541-546.	1.8	19
237	Breast cancer prognostic significance of a single nucleotide polymorphism in the proximal androgen response element of the prostate specific antigen gene promoter. <i>Breast Cancer Research and Treatment</i> , 2000, 61, 111-119.	1.1	18
238	Combined Expression of p53, Bcl-2, and p21WAF-1 Proteins in Lung Cancer and Premalignant Lesions: Association with Clinical Characteristics. <i>Lung</i> , 2001, 179, 265-278.	1.4	18
239	Molecular Response of HL60 Cells to Mitotic Inhibitors Vincristine and Taxol Visualized with Apoptosis-Related Gene Expressions, Including the New Member <i>BCL2L12</i> . <i>Annals of the New York Academy of Sciences</i> , 2009, 1171, 276-283.	1.8	18
240	Expression analysis and study of the <i>KLK15</i> mRNA splice variants in prostate cancer and benign prostatic hyperplasia. <i>Cancer Science</i> , 2010, 101, 693-699.	1.7	18
241	<i>BCL2L12</i> : A promising molecular prognostic biomarker in breast cancer. <i>Clinical Biochemistry</i> , 2014, 47, 257-262.	0.8	18
242	Clinical evaluation of microRNA-145 expression in renal cell carcinoma: a promising molecular marker for discriminating and staging the clear cell histological subtype. <i>Biological Chemistry</i> , 2016, 397, 529-539.	1.2	18
243	MicroRNA-92a-3p overexpression in peripheral blood mononuclear cells is an independent predictor of prolonged overall survival of patients with chronic lymphocytic leukemia. <i>Leukemia and Lymphoma</i> , 2019, 60, 658-667.	0.6	18
244	Discovery of novel transcripts of the human tissue kallikrein (KLK1) and kallikrein-related peptidase 2 (KLK2) in human cancer cells, exploiting Next-Generation Sequencing technology. <i>Genomics</i> , 2019, 111, 642-652.	1.3	18
245	Blood-based analysis of 84 microRNAs identifies molecules deregulated in individuals with type-2 diabetes, risk factors for the disease or metabolic syndrome. <i>Diabetes Research and Clinical Practice</i> , 2020, 164, 108187.	1.1	18
246	The Multifaceted Role and Utility of MicroRNAs in Indolent B-Cell Non-Hodgkin Lymphomas. <i>Biomedicines</i> , 2021, 9, 333.	1.4	18
247	Novel splice variants of prostate-specific antigen and applications in diagnosis of prostate cancer. <i>Clinical Biochemistry</i> , 2008, 41, 591-597.	0.8	17
248	Synthesis, spectroscopic study and anticancer activity of a water-soluble Nb(V) peroxo complex. <i>Journal of Inorganic Biochemistry</i> , 2011, 105, 155-163.	1.5	17
249	Quantified <i>KLK15</i> Gene Expression Levels Discriminate Prostate Cancer From Benign Tumors and Constitute a Novel Independent Predictor of Disease Progression. <i>Prostate</i> , 2013, 73, 1191-1201.	1.2	17
250	Predictions for the future of kallikrein-related peptidases in molecular diagnostics. <i>Expert Review of Molecular Diagnostics</i> , 2014, 14, 713-722.	1.5	17
251	Alpha-enolase is a potential prognostic marker in clear cell renal cell carcinoma. <i>Clinical and Experimental Metastasis</i> , 2015, 32, 531-541.	1.7	17
252	Kallikrein-related peptidase 13: an independent indicator of favorable prognosis for patients with nonsmall cell lung cancer. <i>Tumor Biology</i> , 2015, 36, 4979-4986.	0.8	17

#	ARTICLE	IF	CITATIONS
253	Clinical relevance of the deregulated kallikrein-related peptidase 8 mRNA expression in breast cancer: a novel independent indicator of disease-free survival. <i>Breast Cancer Research and Treatment</i> , 2015, 152, 323-336.	1.1	17
254	Identification and molecular cloning of novel transcripts of the human kallikrein-related peptidase 10 (KLK10) gene using next-generation sequencing. <i>Biochemical and Biophysical Research Communications</i> , 2017, 487, 776-781.	1.0	17
255	The transcriptome of a "sleeping" invader: de novo assembly and annotation of the transcriptome of aestivating <i>Cornu aspersum</i> . <i>BMC Genomics</i> , 2017, 18, 491.	1.2	17
256	Molecular cloning of novel transcripts of human kallikrein-related peptidases 5, 6, 7, 8 and 9 (KLK5-9). <i>Overlock</i> 10	1.6	17
257	Assessing the clinical value of microRNAs in formalin-fixed paraffin-embedded liposarcoma tissues: Overexpressed miR-155 is an indicator of poor prognosis. <i>Oncotarget</i> , 2017, 8, 6896-6913.	0.8	17
258	Clinical utility of miR-143/miR-182 levels in prognosis and risk stratification specificity of BFM-treated childhood acute lymphoblastic leukemia. <i>Annals of Hematology</i> , 2018, 97, 1169-1182.	0.8	17
259	HPV16 E6/E7 expression in circulating tumor cells in oropharyngeal squamous cell cancers: A pilot study. <i>PLoS ONE</i> , 2019, 14, e0215984.	1.1	17
260	Heat shock protein beta 3 (HSPB3) is an unfavorable molecular biomarker in colorectal adenocarcinoma. <i>Molecular Carcinogenesis</i> , 2020, 59, 116-125.	1.3	17
261	Multiple Myeloma Bone Disease: Implication of MicroRNAs in Its Molecular Background. <i>International Journal of Molecular Sciences</i> , 2021, 22, 2375.	1.8	17
262	Alternative Splicing Detection Tool "a novel PERL algorithm for sensitive detection of splicing events, based on next-generation sequencing data analysis. <i>Annals of Translational Medicine</i> , 2018, 6, 244-244.	0.7	17
263	Cathepsin-D and c-erb-B 2 have an additive prognostic value for breast cancer patients. <i>Anticancer Research</i> , 1993, 13, 1895-900.	0.5	17
264	Codon 89 polymorphism in the human 5 α -reductase gene in primary breast cancer. <i>British Journal of Cancer</i> , 2001, 84, 760-767.	2.9	16
265	Treatment of Gastric Cancer Cells with 5-Fluorouracil/Leucovorin and Irinotecan Induces Distinct Alterations in the mRNA Expression of the Apoptosis-Related Genes, Including the Novel Gene <i>BCL2L12</i> . <i>Tumor Biology</i> , 2009, 30, 100-107.	0.8	16
266	KLK5 gene expression is severely upregulated in androgen-independent prostate cancer cells after treatment with the chemotherapeutic agents docetaxel and mitoxantrone. <i>Biological Chemistry</i> , 2010, 391, 467-74.	1.2	16
267	L-DOPA Decarboxylase (DDC) Expression Status as a Novel Molecular Tumor Marker for Diagnostic and Prognostic Purposes in Laryngeal Cancer. <i>Translational Oncology</i> , 2012, 5, 288-296.	1.7	16
268	Expression of Bcl2L12 in chronic lymphocytic leukemia patients: association with clinical and molecular prognostic markers. <i>Medical Oncology</i> , 2013, 30, 405.	1.2	16
269	The Stat3/5 Signaling Biosignature in Hematopoietic Stem/Progenitor Cells Predicts Response and Outcome in Myelodysplastic Syndrome Patients Treated with Azacitidine. <i>Clinical Cancer Research</i> , 2016, 22, 1958-1968.	3.2	16
270	Downregulated KLK13 expression in bladder cancer highlights tumor aggressiveness and unfavorable patients' prognosis. <i>Journal of Cancer Research and Clinical Oncology</i> , 2017, 143, 521-532.	1.2	16

#	ARTICLE	IF	CITATIONS
271	Expression Analysis of miR-29b in Malignant and Benign Breast Tumors: A Promising Prognostic Biomarker for Invasive Ductal Carcinoma With a Possible Histotype-Related Expression Status. <i>Clinical Breast Cancer</i> , 2018, 18, 305-312.e3.	1.1	16
272	Expression of BCL2L12, a new member of apoptosis-related genes, in breast tumors. <i>Thrombosis and Haemostasis</i> , 2003, 89, 1081-8.	1.8	16
273	DRAMATIC SUPPRESSION OF PLASMA AND URINARY PROSTATE SPECIFIC ANTIGEN AND HUMAN GLANDULAR KALLIKREIN BY ANTIANDROGENS IN MALE-TO-FEMALE TRANSSEXUALS. <i>Journal of Urology</i> , 2000, 163, 802-805.	0.2	15
274	Kallikrein-related peptidase 6 (KLK6) gene expression in intracranial tumors. <i>Tumor Biology</i> , 2012, 33, 1375-1383.	0.8	15
275	mRNA overexpression of kallikrein-related peptidase 14 (KLK14) is an independent predictor of poor overall survival in chronic lymphocytic leukemia patients. <i>Clinical Chemistry and Laboratory Medicine</i> , 2016, 54, 315-24.	1.4	15
276	Unravelling a p73-regulated network: The role of a novel p73-dependent target, MIR3158, in cancer cell migration and invasiveness. <i>Cancer Letters</i> , 2017, 388, 96-106.	3.2	15
277	Identification and Characterization of a Novel Human Testis-Specific Kinase Substrate Gene Which Is Downregulated in Testicular Tumors. <i>Biochemical and Biophysical Research Communications</i> , 2001, 285, 400-408.	1.0	14
278	Evaluation and prognostic significance of human tissue kallikrein-related peptidase 10 (KLK10) in colorectal cancer. <i>Tumor Biology</i> , 2012, 33, 1209-1214.	0.8	14
279	The role of transcription factors in laboratory medicine. <i>Clinical Chemistry and Laboratory Medicine</i> , 2013, 51, 1563-1571.	1.4	14
280	Low mRNA expression levels of kallikrein-related peptidase 4 (KLK4) predict short-term relapse in patients with laryngeal squamous cell carcinoma. <i>Biological Chemistry</i> , 2014, 395, 1051-1062.	1.2	14
281	S100A11 is a potential prognostic marker for clear cell renal cell carcinoma. <i>Clinical and Experimental Metastasis</i> , 2016, 33, 63-71.	1.7	14
282	Novel alternative splice variants of the human protein arginine methyltransferase 1 (PRMT1) gene, discovered using next-generation sequencing. <i>Gene</i> , 2019, 699, 135-144.	1.0	14
283	Unraveling UCA1 lncRNA prognostic utility in urothelial bladder cancer. <i>Carcinogenesis</i> , 2019, 40, 965-974.	1.3	14
284	Comparison of the percent free prostate-specific antigen levels in the serum of healthy men and in men with recurrent prostate cancer after radical prostatectomy. <i>Clinica Chimica Acta</i> , 2000, 292, 127-138.	0.5	13
285	Effect of Testosterone Administration on Serum and Urine Kallikrein Concentrations in Female-to-Male Transsexuals. <i>Clinical Chemistry</i> , 2006, 52, 1546-1551.	1.5	13
286	Identification of a STAT5 Target Gene, Dpf3, Provides Novel Insights in Chronic Lymphocytic Leukemia. <i>PLoS ONE</i> , 2013, 8, e76155.	1.1	13
287	A Cancer-Related microRNA Signature Shows Biomarker Utility in Multiple Myeloma. <i>International Journal of Molecular Sciences</i> , 2021, 22, 13144.	1.8	13
288	Enhanced Concentration-Dependent Cytotoxic Effect of the Dinuclear Copper(II) Complex of L-Carnitine [Cu ₂ (L-carnitine) ₂ Cl ₂ (H ₂ O) ₂]Cl ₂ ·2H ₂ O Compared to L-Carnitine or Copper Chloride Dihydrate, in Human Leukemic Cell Lines. <i>Journal of Medicinal Chemistry</i> , 2008, 51, 3713-3719.	0.2	12

#	ARTICLE	IF	CITATIONS
289	Kallikrein-related peptidase 6 (KLK6) expression differentiates tumor subtypes and predicts clinical outcome in breast cancer patients. <i>Clinical and Experimental Medicine</i> , 2018, 18, 203-213.	1.9	12
290	Unraveling novel survivin mRNA transcripts in cancer cells using an in-house developed targeted high-throughput sequencing approach. <i>Genomics</i> , 2021, 113, 573-581.	1.3	12
291	SARS-CoV-2 Infection Is Asymptomatic in Nearly Half of Adults with Robust Anti-Spike Protein Receptor-Binding Domain Antibody Response. <i>Vaccines</i> , 2021, 9, 207.	2.1	12
292	A Molecular Signature of Circulating MicroRNA Can Predict Osteolytic Bone Disease in Multiple Myeloma. <i>Cancers</i> , 2021, 13, 3877.	1.7	12
293	High Expression of a tRNAPro Derivative Associates with Poor Survival and Independently Predicts Colorectal Cancer Recurrence. <i>Biomedicines</i> , 2022, 10, 1120.	1.4	12
294	10-(2-Biotinyloxyethyl)-9-acridone. <i>Journal of Photochemistry and Photobiology A: Chemistry</i> , 2006, 181, 126-131.	2.0	11
295	Comparative HPLC-DAD and UHPLC-ESI(-)-HRMS & MS/MS profiling of Hypericum species and correlation with necrotic cell-death activity in human leukemic cells. <i>Phytochemistry Letters</i> , 2017, 20, 481-490.	0.6	11
296	Human kallikrein-related peptidase 12 (KLK12) splice variants discriminate benign from cancerous breast tumors. <i>Clinical Biochemistry</i> , 2018, 58, 78-85.	0.8	11
297	A 3â€² tRNAâ€derived fragment produced by tRNA LeuAAG and tRNA LeuTAG is associated with poor prognosis in Bâ€cell chronic lymphocytic leukemia, independently of classical prognostic factors. <i>European Journal of Haematology</i> , 2021, 106, 821-830.	1.1	11
298	Novel Nested-Seq Approach for SARS-CoV-2 Real-Time Epidemiology and In-Depth Mutational Profiling in Wastewater. <i>International Journal of Molecular Sciences</i> , 2021, 22, 8498.	1.8	11
299	miRNA-seq and clinical evaluation in multiple myeloma: miR-181a overexpression predicts short-term disease progression and poor post-treatment outcome. <i>British Journal of Cancer</i> , 2022, 126, 79-90.	2.9	11
300	Identification of Novel Circular RNAs of the Human Protein Arginine Methyltransferase 1 (PRMT1) Gene, Expressed in Breast Cancer Cells. <i>Genes</i> , 2022, 13, 1133.	1.0	11
301	Novel biotinylated acridinium derivatives: New reagents for fluorescence immunoassays and proteomics. <i>Clinica Chimica Acta</i> , 2005, 357, 159-167.	0.5	10
302	Quantitative expression analysis and study of the novel human kallikrein-related peptidase 14 gene (KLK14) in malignant and benign breast tissues. <i>Thrombosis and Haemostasis</i> , 2011, 105, 131-137.	1.8	10
303	Overexpression of the novel member of the BCL2 gene family, BCL2L12, is associated with the disease outcome in patients with acute myeloid leukemia. <i>Clinical Biochemistry</i> , 2012, 45, 1362-1367.	0.8	10
304	The kallikrein-related peptidase 13 (KLK13) gene is substantially up-regulated after exposure of gastric cancer cells to antineoplastic agents. <i>Tumor Biology</i> , 2012, 33, 2069-2078.	0.8	10
305	Increased BCL2L12 expression predicts the short-term relapse of patients with TaT1 bladder cancer following transurethral resection of bladder tumors. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014, 32, 39.e29-39.e36.	0.8	10
306	Overexpression of BCL2 and BAX following BFM induction therapy predicts ch-ALL patientsâ€™ poor response to treatment and short-term relapse. <i>Journal of Cancer Research and Clinical Oncology</i> , 2015, 141, 2023-2036.	1.2	10

#	ARTICLE	IF	CITATIONS
307	Cisplatin and Paclitaxel Alter the Expression Pattern of miRâ€143/145 and miRâ€183/96/182 Clusters in T24 Bladder Cancer Cells. <i>Clinical and Translational Science</i> , 2015, 8, 668-675.	1.5	10
308	Gemcitabine impacts differentially on bladder and kidney cancer cells: distinct modulations in the expression patterns of apoptosis-related microRNAs and BCL2 family genes. <i>Tumor Biology</i> , 2015, 36, 3197-3207.	0.8	10
309	BCL2L12: a multiply spliced gene with independent prognostic significance in breast cancer. <i>Clinical Chemistry and Laboratory Medicine</i> , 2018, 57, 276-287.	1.4	10
310	BCL2L12 improves risk stratification and prediction of BFM-chemotherapy response in childhood acute lymphoblastic leukemia. <i>Clinical Chemistry and Laboratory Medicine</i> , 2018, 56, 2104-2118.	1.4	10
311	Novel splice variants of the human kallikrein-related peptidases 11 (<i>KLK11</i>) and 12 (<i>KLK12</i>), unraveled by next-generation sequencing technology. <i>Biological Chemistry</i> , 2018, 399, 1065-1071.	1.2	10
312	Kallikrein-related peptidases and associated microRNAs as promising prognostic biomarkers in gastrointestinal malignancies. <i>Biological Chemistry</i> , 2018, 399, 821-836.	1.2	10
313	Identification of novel alternative splice variants of the human L-DOPA decarboxylase (DDC) gene in human cancer cells, using high-throughput sequencing approaches. <i>Gene</i> , 2019, 719, 144075.	1.0	10
314	miR-203 is an independent molecular predictor of prognosis and treatment outcome in ovarian cancer: a multi-institutional study. <i>Carcinogenesis</i> , 2020, 41, 442-451.	1.3	10
315	Revised Exon Structure of L-DOPA Decarboxylase (DDC) Reveals Novel Splice Variants Associated with Colorectal Cancer Progression. <i>International Journal of Molecular Sciences</i> , 2020, 21, 8568.	1.8	10
316	Next-generation sequencing reveals alternative L-DOPA decarboxylase (DDC) splice variants bearing novel exons, in human hepatocellular and lung cancer cells. <i>Gene</i> , 2021, 768, 145262.	1.0	10
317	Quantitative analysis and study of the mRNA expression levels of apoptotic genes BCL2, BAX and BCL2L12 in the articular cartilage of an animal model of osteoarthritis. <i>Annals of Translational Medicine</i> , 2018, 6, 243-243.	0.7	10
318	Metformin and Anti-Cancer Therapeutics: Hopes for a More Enhanced Armamentarium Against Human Neoplasias?. <i>Current Medicinal Chemistry</i> , 2017, 24, 14-56.	1.2	10
319	Molecular Effects of Treatment of Human Colorectal Cancer Cells with Natural and Classical Chemotherapeutic Drugs: Alterations in the Expression of Apoptosis-related BCL2 Family Members, Including BCL2L12. <i>Current Pharmaceutical Biotechnology</i> , 2019, 19, 1064-1075.	0.9	10
320	Evidence for L-Dopa Decarboxylase Involvement in Cancer Cell Cytotoxicity Induced by Docetaxel and Mitoxantrone. <i>Current Pharmaceutical Biotechnology</i> , 2019, 19, 1087-1096.	0.9	10
321	Relationships between cathepsin-D, pS2 protein and hormonal receptors in breast cancer cytosols: inconsistency with their established prognostic significance. <i>Anticancer Research</i> , 1997, 17, 3665-9.	0.5	10
322	Expression analysis and study of KLK4 in benign and malignant breast tumours. <i>Thrombosis and Haemostasis</i> , 2009, 101, 381-7.	1.8	10
323	Expression and prognostic significance of kallikrein-related peptidase 8 protein levels in advanced ovarian cancer by using automated quantitative analysis. <i>Thrombosis and Haemostasis</i> , 2009, 101, 541-6.	1.8	10
324	SR-A1, a member of the human pre-mRNA splicing factor family, and its expression in colon cancer progression. <i>Biological Chemistry</i> , 2004, 385, 785-90.	1.2	9

#	ARTICLE	IF	CITATIONS
325	Multidisciplinary Therapy of Locally Advanced or Inflammatory Breast Cancer with Fixed Perioperative Sequence of Epirubicin, Vinorelbine, and Fluorouracil Chemotherapy, Surgery, and Radiotherapy: Long-Term Results. <i>Oncologist</i> , 2006, 11, 563-573.	1.9	9
326	BCL2L12 protein overexpression is associated with favorable outcome in diffuse large B-cell lymphoma patients in the rituximab era. <i>Leukemia and Lymphoma</i> , 2016, 57, 2199-2203.	0.6	9
327	Discovery and expression analysis of novel transcripts of the human SR-related CTD-associated factor 1 (SCAF1) gene in human cancer cells using Next-Generation Sequencing. <i>Gene</i> , 2018, 670, 155-165.	1.0	9
328	Molecular cloning of novel transcripts of the adaptor-related protein complex 2 alpha 1 subunit (AP2A1) gene, using Next-Generation Sequencing. <i>Gene</i> , 2018, 678, 55-64.	1.0	9
329	Uncovering the clinical impact of kallikrein-related peptidase 5 (<i>KLK5</i>) mRNA expression in the colorectal adenoma-carcinoma sequence. <i>Clinical Chemistry and Laboratory Medicine</i> , 2019, 57, 1251-1260.	1.4	9
330	c-erbB-2 overexpression may be used as an independent prognostic factor for breast cancer patients. <i>Anticancer Research</i> , 1995, 15, 1543-7.	0.5	9
331	Evaluation of kallikrein-related peptidase 5 expression and its significance for breast cancer patients: association with kallikrein-related peptidase 7 expression. <i>Anticancer Research</i> , 2011, 31, 3093-100.	0.5	9
332	Cloning, physical mapping and structural characterization of the human β -A-adaptin gene. <i>Gene</i> , 2002, 289, 191-199.	1.0	8
333	Expression analysis and prognostic significance of the SRA1 gene, in ovarian cancer. <i>Biochemical and Biophysical Research Communications</i> , 2006, 344, 667-674.	1.0	8
334	L-Dopa decarboxylase (DDC) constitutes an emerging biomarker in predicting patients' survival with stomach adenocarcinomas. <i>Journal of Cancer Research and Clinical Oncology</i> , 2013, 139, 297-306.	1.2	8
335	L-DOPA decarboxylase mRNA levels provide high diagnostic accuracy and discrimination between clear cell and non-clear cell subtypes in renal cell carcinoma. <i>Clinical Biochemistry</i> , 2015, 48, 590-595.	0.8	8
336	A comprehensive clinicopathological evaluation of the differential expression of microRNA-331 in breast tumors and its diagnostic significance. <i>Clinical Biochemistry</i> , 2018, 60, 24-32.	0.8	8
337	Gene-Specific Intron Retention Serves as Molecular Signature that Distinguishes Melanoma from Non-Melanoma Cancer Cells in Greek Patients. <i>International Journal of Molecular Sciences</i> , 2019, 20, 937.	1.8	8
338	Pediatric Ependymoma: A Proteomics Perspective. <i>Cancer Genomics and Proteomics</i> , 2017, 14, 127-136.	1.0	8
339	tRNA Derivatives in Multiple Myeloma: Investigation of the Potential Value of a tRNA-Derived Molecular Signature. <i>Biomedicines</i> , 2021, 9, 1811.	1.4	8
340	A versatile 5' RACE-Seq methodology for the accurate identification of the 5' termini of mRNAs. <i>BMC Genomics</i> , 2022, 23, 163.	1.2	8
341	Expression of the C-terminal domain of novel human SR-A1 protein: Interaction with the CTD domain of RNA polymerase II. <i>Biochemical and Biophysical Research Communications</i> , 2005, 334, 61-68.	1.0	7
342	The Immunohistochemical Expression of Growth Hormone-Releasing Hormone Receptor Splice Variant 1 Is a Favorable Prognostic Marker in Colorectal Cancer. <i>Molecular Medicine</i> , 2009, 15, 242-247.	1.9	7

#	ARTICLE	IF	CITATIONS
343	Immunohistochemical expression of somatostatin receptor subtypes 2 and 5 in colorectal cancer. <i>European Journal of Clinical Investigation</i> , 2012, 42, 777-783.	1.7	7
344	Significant alterations in the expression pattern of kallikrein-related peptidase genes KLK4, KLK5 and KLK14 after treatment of breast cancer cells with the chemotherapeutic agents epirubicin, docetaxel and methotrexate. <i>Tumor Biology</i> , 2013, 34, 369-378.	0.8	7
345	Contribution of miRNAs, tRNAs and tRFs to Aberrant Signaling and Translation Deregulation in Lung Cancer. <i>Cancers</i> , 2020, 12, 3056.	1.7	7
346	miR-181a overexpression predicts the poor treatment response and early progression of serous ovarian cancer patients. <i>International Journal of Cancer</i> , 2020, 147, 3560-3573.	2.3	7
347	Prognostic significance of the expression of SR-A1, encoding a novel SR-related CTD-associated factor, in breast cancer. <i>Biological Chemistry</i> , 2006, 387, 1613-8.	1.2	6
348	Total and free PSA kinetics in patients without prostate cancer undergoing radical cystoprostatectomy. <i>Prostate</i> , 2008, 68, 759-765.	1.2	6
349	Kallikrein-related peptidases (KLKs) as novel potential biomarkers in gastric cancer: An open yet challenging road lies ahead. <i>Journal of Surgical Oncology</i> , 2012, 105, 223-224.	0.8	6
350	Nanopore Sequencing Unveils Diverse Transcript Variants of the Epithelial Cell-Specific Transcription Factor Elf-3 in Human Malignancies. <i>Genes</i> , 2021, 12, 839.	1.0	6
351	Prostate-Specific Antigen and Human Glandular Kallikrein 2 Are Markedly Elevated in Urine of Patients with Polycystic Ovary Syndrome. , 0, .		6
352	Cathepsin D may help in discriminating node-negative breast cancer patients at risk for local-regional recurrence. <i>Anticancer Research</i> , 1998, 18, 2885-90.	0.5	6
353	TA repeat polymorphism of the 5alpha-reductase gene and breast cancer. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2000, 9, 387-93.	1.1	6
354	Treatment of PC3 prostate cancer cells with mitoxantrone, etoposide, doxorubicin and carboplatin induces distinct alterations in the expression of kallikreins 5 and 11. <i>Thrombosis and Haemostasis</i> , 2009, 101, 373-80.	1.8	6
355	Safety and Efficacy of Trastuzumab Every 3 Weeks Combined with Cytotoxic Chemotherapy in Patients with HER2-Positive Recurrent Breast Cancer: Findings from a Case Series. <i>Oncology Research and Treatment</i> , 2005, 28, 558-564.	0.8	5
356	Serum and Urine Tissue Kallikrein Concentrations in Male-to-Female Transsexuals Treated with Antiandrogens and Estrogens. <i>Clinical Chemistry</i> , 2006, 52, 1356-1365.	1.5	5
357	Cathepsin D concentration in tumor cytosols improves the accuracy of prognostic evaluation of primary breast cancer. <i>Anticancer Research</i> , 1997, 17, 1405-9.	0.5	5
358	Overexpression of the GR Riborepressor LncRNA GAS5 Results in Poor Treatment Response and Early Relapse in Childhood B-ALL. <i>Cancers</i> , 2021, 13, 6064.	1.7	5
359	<sc>Epi-miRNAs</sc>: Modern mediators of methylation status in human cancers. <i>Wiley Interdisciplinary Reviews RNA</i> , 2023, 14, e1735.	3.2	5
360	The kallikreins: old proteases with new clinical potentials. <i>Thrombosis and Haemostasis</i> , 2013, 110, 396-398.	1.8	4

#	ARTICLE	IF	CITATIONS
361	Human L-DOPA decarboxylase mRNA is a target of miR-145: A prediction to validation workflow. <i>Gene</i> , 2015, 554, 174-180.	1.0	4
362	Expressional profiling and clinical relevance of RNase H ¹ in prostate cancer: a novel indicator of favorable progression-free survival. <i>Journal of Cancer Research and Clinical Oncology</i> , 2018, 144, 2049-2057.	1.2	4
363	¹²⁵ I-Np63 transcript loss in bladder cancer constitutes an independent molecular predictor of TaT1 patients post-treatment relapse and progression. <i>Journal of Cancer Research and Clinical Oncology</i> , 2019, 145, 3075-3087.	1.2	4
364	Complex transcriptional regulation of the BCL2L12 gene: Novel, active promoter in K562 cells. <i>Gene</i> , 2020, 750, 144723.	1.0	4
365	Molecular Biomarkers of Laryngeal Cancer Laryngeal squamous cell carcinoma (LSCC) Tumor markers. <i>Biomarkers in Disease</i> , 2015, , 891-919.	0.0	4
366	mRNA quantification and clinical evaluation of telomerase reverse transcriptase subunit (hTERT) in intracranial tumours of patients in the island of Crete. <i>British Journal of Cancer</i> , 2005, 93, 152-158.	2.9	3
367	DDC (dopa decarboxylase (aromatic L-amino acid decarboxylase)). <i>Atlas of Genetics and Cytogenetics in Oncology and Haematology</i> , 2012, , .	0.1	3
368	Editorial (Thematic Issue: The Effects of Anticancer Agents on Cell Apoptosis and on the Expression of) <i>Tj ETQq0 0 0 rgBT /Overlock 10 T</i>	0.9	3
369	Multianalyte quantitative competitive PCR on optically encoded microspheres for an eight-gene panel related to prostate cancer. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 971-980.	1.9	3
370	Identification of novel alternative transcripts of the human Ribonuclease H ¹ (RNASEK) gene using 3 ^{â€²} RACE and high-throughput sequencing approaches. <i>Genomics</i> , 2020, 112, 943-951.	1.3	3
371	Identification and expression analysis of novel splice variants of the human carcinoembryonic antigen-related cell adhesion molecule 19 (CEACAM19) gene using a high-throughput sequencing approach. <i>Genomics</i> , 2020, 112, 4268-4276.	1.3	3
372	Jagged Ends of Cell-Free DNA: Rebranding Fragmentomics in Modern Liquid Biopsy Diagnostics. <i>Clinical Chemistry</i> , 2021, 67, 576-578.	1.5	3
373	Pharmacoepigenomics circuits induced by a novel retinoid-polyamine conjugate in human immortalized keratinocytes. <i>Pharmacogenomics Journal</i> , 2021, 21, 638-648.	0.9	3
374	Identification and expression analysis of ten novel small non-coding RNAs (sncRNAs) in cancer cells using a high-throughput sequencing approach. <i>Gene</i> , 2022, 809, 146025.	1.0	3
375	Poly(A)polymerase activity levels in breast tumour cytosols. <i>Journal of Experimental and Clinical Cancer Research</i> , 1998, 17, 511-8.	0.4	3
376	10 Kallikrein-related Peptidases as Biomarkers in Personalized Cancer Medicine. , 2012, , 201-218.		2
377	Quantification and study of the L-DOPA decarboxylase expression in gastric adenocarcinoma cells treated with chemotherapeutic substances. <i>Anti-Cancer Drugs</i> , 2013, 24, 291-299.	0.7	2
378	Kallikreins as Biomarkers in Human Malignancies. <i>Biomarkers in Disease</i> , 2015, , 135-165.	0.0	2

#	ARTICLE	IF	CITATIONS
379	Quantitative Analysis of Kallikrein 15 Gene Expression in Prostate Tissue. <i>Journal of Urology</i> , 2003, , 361-364.	0.2	2
380	Targeted Long-Read Sequencing Decodes the Transcriptional Atlas of the Founding RAS Gene Family Members. <i>International Journal of Molecular Sciences</i> , 2021, 22, 13298.	1.8	2
381	Effect of bleomycin and cisplatin on the expression profile of <i>SRA1</i> , a novel member of pre-mRNA splicing factors, in HL-60 human promyelocytic leukemia cells. <i>Biological Chemistry</i> , 2007, 388, 773-778.	1.2	1
382	Highlight: Kinin 2012 in Paris. <i>Biological Chemistry</i> , 2013, 393, 299-300.	1.2	1
383	THE tRNA-DERIVED RNA FRAGMENTS (tRFs) BEARING THE GLYCINE ANTICODONS GCC AND CCC AS EMERGING MOLECULAR BIOMARKERS OF UNFAVORABLE PROGNOSIS IN CHRONIC LYMPHOBLASTIC LEUKEMIA. <i>Hematological Oncology</i> , 2019, 37, 375-376.	0.8	1
384	A comprehensive nanopore sequencing methodology deciphers the complete transcriptional landscape of cyclin dependent kinase 4 (CDK4) in human malignancies. <i>FEBS Journal</i> , 2021, , .	2.2	1
385	Molecular Biomarkers of Laryngeal Cancer. , 2014, , 1-24.		1
386	A Molecular Signature of Three tRNA-Derived RNA Fragments May Discriminate Smoldering from Symptomatic Multiple Myeloma Patients. <i>Blood</i> , 2019, 134, 5528-5528.	0.6	1
387	Evaluation of the clinical utility of kallikrein-related peptidase 6 gene (KLK6) downregulation in breast cancer.. <i>Journal of Clinical Oncology</i> , 2012, 30, 10606-10606.	0.8	1
388	Kallikreins as Biomarkers in Human Malignancies. , 2014, , 1-25.		1
389	BCL2L12 (BCL2-like 12 (proline-rich)). <i>Atlas of Genetics and Cytogenetics in Oncology and Haematology</i> , 2011, , .	0.1	0
390	SCAF1 (SR-related CTD-associated factor 1). <i>Atlas of Genetics and Cytogenetics in Oncology and Haematology</i> , 2011, , .	0.1	0
391	Manfred Schmitt (1947-2018). <i>Biological Chemistry</i> , 2018, 399, 923-924.	1.2	0
392	The interplay between miR-1245a and BRCA2 in colorectal cancer. <i>Annals of Translational Medicine</i> , 2020, 8, 1043-1043.	0.7	0
393	Identification of six novel alternative transcripts of the human kallikrein-related peptidase 15 (KLK15), using 5'RACE and high-throughput sequencing. <i>Gene</i> , 2020, 749, 144708.	1.0	0
394	Association of BCL2L12 overexpression with prolonged disease-free survival in breast cancer.. <i>Journal of Clinical Oncology</i> , 2014, 32, e22202-e22202.	0.8	0
395	Steroid Hormone Regulation and Prognostic Value of the Human Kallikrein Gene 14 in Ovarian Cancer. <i>American Journal of Clinical Pathology</i> , 2003, 119, 0-0.	0.4	0
396	Elevated levels of both microRNA 378 (miR-378) and kallikrein-related peptidase 4 (KLK4) mRNA are associated with an unfavorable prognosis in triple-negative breast cancer. <i>American Journal of Translational Research (discontinued)</i> , 2021, 13, 1594-1606.	0.0	0