

Andreas Scorilas

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8343178/andreas-scorilas-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

386
papers

11,018
citations

55
h-index

76
g-index

399
ext. papers

12,428
ext. citations

5.1
avg. IF

6.37
L-index

#	Paper	IF	Citations
386	Identification and expression analysis of ten novel small non-coding RNAs (sncRNAs) in cancer cells using a high-throughput sequencing approach. <i>Gene</i> , 2022 , 809, 146025	3.8	0
385	SARS-CoV-2 wastewater surveillance data can predict hospitalizations and ICU admissions. <i>Science of the Total Environment</i> , 2022 , 804, 150151	10.2	13
384	A versatile 5ORACE-Seq methodology for the accurate identification of the 5' termini of mRNAs.. <i>BMC Genomics</i> , 2022 , 23, 163	4.5	2
383	Adverse effects of COVID-19 mRNA vaccines: the spike hypothesis.. <i>Trends in Molecular Medicine</i> , 2022 ,	11.5	14
382	High Expression of a tRNA ^{Pro} Derivative Associates with Poor Survival and Independently Predicts Colorectal Cancer Recurrence. <i>Biomedicines</i> , 2022 , 10, 1120	4.8	0
381	Epi-miRNAs: Modern mediators of methylation status in human cancers.. <i>Wiley Interdisciplinary Reviews RNA</i> , 2022 , e1735	9.3	1
380	Elevated levels of both microRNA 378 (miR-378) and kallikrein-related peptidase 4 (KLK4) mRNA are associated with an unfavorable prognosis in triple-negative breast cancer. <i>American Journal of Translational Research (discontinued)</i> , 2021 , 13, 1594-1606	3	
379	tRNA Derivatives in Multiple Myeloma: Investigation of the Potential Value of a tRNA-Derived Molecular Signature.. <i>Biomedicines</i> , 2021 , 9,	4.8	2
378	miRNA-seq and clinical evaluation in multiple myeloma: miR-181a overexpression predicts short-term disease progression and poor post-treatment outcome. <i>British Journal of Cancer</i> , 2021 ,	8.7	4
377	SARS-CoV-2 Infection Is Asymptomatic in Nearly Half of Adults with Robust Anti-Spike Protein Receptor-Binding Domain Antibody Response. <i>Vaccines</i> , 2021 , 9,	5.3	6
376	The Multifaceted Role and Utility of MicroRNAs in Indolent B-Cell Non-Hodgkin Lymphomas. <i>Biomedicines</i> , 2021 , 9,	4.8	11
375	Nanopore Sequencing Unveils Diverse Transcript Variants of the Epithelial Cell-Specific Transcription Factor Elf-3 in Human Malignancies. <i>Genes</i> , 2021 , 12,	4.2	2
374	Circular RNAs: Emerging Regulators of the Major Signaling Pathways Involved in Cancer Progression. <i>Cancers</i> , 2021 , 13,	6.6	16
373	Pharmacoepiggenomics circuits induced by a novel retinoid-polyamine conjugate in human immortalized keratinocytes. <i>Pharmacogenomics Journal</i> , 2021 , 21, 638-648	3.5	0
372	A Molecular Signature of Circulating MicroRNA Can Predict Osteolytic Bone Disease in Multiple Myeloma. <i>Cancers</i> , 2021 , 13,	6.6	5
371	Next-generation sequencing reveals alternative L-DOPA decarboxylase (DDC) splice variants bearing novel exons, in human hepatocellular and lung cancer cells. <i>Gene</i> , 2021 , 768, 145262	3.8	3
370	Unraveling novel survivin mRNA transcripts in cancer cells using an in-house developed targeted high-throughput sequencing approach. <i>Genomics</i> , 2021 , 113, 573-581	4.3	7

369	Analytical methodologies for the detection of SARS-CoV-2 in wastewater: Protocols and future perspectives. <i>TrAC - Trends in Analytical Chemistry</i> , 2021 , 134, 116125	14.6	40
368	MicroRNAs: Tiny Regulators of Gene Expression with Pivotal Roles in Normal B-Cell Development and B-Cell Chronic Lymphocytic Leukemia. <i>Cancers</i> , 2021 , 13,	6.6	15
367	Multiple Myeloma Bone Disease: Implication of MicroRNAs in Its Molecular Background. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	12
366	A 3'UTR-derived fragment produced by tRNA and tRNA is associated with poor prognosis in B-cell chronic lymphocytic leukemia, independently of classical prognostic factors. <i>European Journal of Haematology</i> , 2021 , 106, 821-830	3.8	5
365	Novel Nested-Seq Approach for SARS-CoV-2 Real-Time Epidemiology and In-Depth Mutational Profiling in Wastewater. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	3
364	Comparative kinetics of SARS-CoV-2 anti-spike protein RBD IgGs and neutralizing antibodies in convalescent and naïve recipients of the BNT162b2 mRNA vaccine versus COVID-19 patients. <i>BMC Medicine</i> , 2021 , 19, 208	11.4	19
363	tRNA-Derived Internal Fragment (i-tRF-GlyGCC) in Ovarian Cancer Treatment Outcome and Progression.. <i>Cancers</i> , 2021 , 14,	6.6	2
362	Third-Generation Sequencing: The Spearhead towards the Radical Transformation of Modern Genomics.. <i>Life</i> , 2021 , 12,	3	5
361	A Cancer-Related microRNA Signature Shows Biomarker Utility in Multiple Myeloma. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	1
360	Targeted Long-Read Sequencing Decodes the Transcriptional Atlas of the Founding RAS Gene Family Members.. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	1
359	tRNA-Derived Fragments (tRFs) in Bladder Cancer: Increased 5'tRF-LysCTT Results in Disease Early Progression and Patients' Poor Treatment Outcome. <i>Cancers</i> , 2020 , 12,	6.6	8
358	Complex transcriptional regulation of the BCL2L12 gene: Novel, active promoter in K562 cells. <i>Gene</i> , 2020 , 750, 144723	3.8	4
357	miR-181a overexpression predicts the poor treatment response and early-progression of serous ovarian cancer patients. <i>International Journal of Cancer</i> , 2020 , 147, 3560-3573	7.5	4
356	JQ1 inhibits tumour growth in combination with cisplatin and suppresses JAK/STAT signalling pathway in ovarian cancer. <i>European Journal of Cancer</i> , 2020 , 126, 125-135	7.5	22
355	Blood-based analysis of 84 microRNAs identifies molecules deregulated in individuals with type-2 diabetes, risk factors for the disease or metabolic syndrome. <i>Diabetes Research and Clinical Practice</i> , 2020 , 164, 108187	7.4	9
354	Identification of six novel alternative transcripts of the human kallikrein-related peptidase 15 (KLK15), using 3'ACE and high-throughput sequencing. <i>Gene</i> , 2020 , 749, 144708	3.8	
353	High clusterin (CLU) mRNA expression levels in tumors of colorectal cancer patients predict a poor prognostic outcome. <i>Clinical Biochemistry</i> , 2020 , 75, 62-69	3.5	15
352	Heat shock protein beta 3 (HSPB3) is an unfavorable molecular biomarker in colorectal adenocarcinoma. <i>Molecular Carcinogenesis</i> , 2020 , 59, 116-125	5	8

351	Computational approaches in cancer multidrug resistance research: Identification of potential biomarkers, drug targets and drug-target interactions. <i>Drug Resistance Updates</i> , 2020 , 48, 100662	23.2	22
350	Next generation sequencing targeted gene panel in Greek MODY patients increases diagnostic accuracy. <i>Pediatric Diabetes</i> , 2020 , 21, 28-39	3.6	12
349	Seroprevalence of Antibodies against SARS-CoV-2 among the Personnel and Students of the National and Kapodistrian University of Athens, Greece: A Preliminary Report. <i>Life</i> , 2020 , 10,	3	20
348	Revised Exon Structure of L-DOPA Decarboxylase () Reveals Novel Splice Variants Associated with Colorectal Cancer Progression. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	1
347	Identification of Two Novel Circular RNAs Deriving from and Investigation of Their Potential Value as a Molecular Signature in Colorectal Cancer. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	9
346	Identification and expression analysis of novel splice variants of the human carcinoembryonic antigen-related cell adhesion molecule 19 (CEACAM19) gene using a high-throughput sequencing approach. <i>Genomics</i> , 2020 , 112, 4268-4276	4.3	3
345	A novel, mitochondrial, internal tRNA-derived RNA fragment possesses clinical utility as a molecular prognostic biomarker in chronic lymphocytic leukemia. <i>Clinical Biochemistry</i> , 2020 , 85, 20-26	3.5	14
344	The role of circular RNAs in therapy resistance of patients with solid tumors. <i>Personalized Medicine</i> , 2020 , 17, 469-490	2.2	17
343	Contribution of miRNAs, tRNAs and tRFs to Aberrant Signaling and Translation Deregulation in Lung Cancer. <i>Cancers</i> , 2020 , 12,	6.6	3
342	Circular RNAs: A New Piece in the Colorectal Cancer Puzzle. <i>Cancers</i> , 2020 , 12,	6.6	25
341	miR-203 is an independent molecular predictor of prognosis and treatment outcome in ovarian cancer: a multi-institutional study. <i>Carcinogenesis</i> , 2020 , 41, 442-451	4.6	4
340	Identification of novel alternative transcripts of the human Ribonuclease II (RNASEK) gene using 30 RACE and high-throughput sequencing approaches. <i>Genomics</i> , 2020 , 112, 943-951	4.3	1
339	Identification of novel alternative splice variants of the human L-DOPA decarboxylase (DDC) gene in human cancer cells, using high-throughput sequencing approaches. <i>Gene</i> , 2019 , 719, 144075	3.8	5
338	The lysine-specific methyltransferase KMT2C/MLL3 regulates DNA repair components in cancer. <i>EMBO Reports</i> , 2019 , 20,	6.5	46
337	THE tRNA-DERIVED RNA FRAGMENTS (tRFs) BEARING THE GLYCINE ANTICODONS GCC AND CCC AS EMERGING MOLECULAR BIOMARKERS OF UNFAVORABLE PROGNOSIS IN CHRONIC LYMPHOCYTIC LEUKEMIA. <i>Hematological Oncology</i> , 2019 , 37, 375-376	1.3	
336	HPV16 E6/E7 expression in circulating tumor cells in oropharyngeal squamous cell cancers: A pilot study. <i>PLoS ONE</i> , 2019 , 14, e0215984	3.7	10
335	Novel alternative splice variants of the human protein arginine methyltransferase 1 (PRMT1) gene, discovered using next-generation sequencing. <i>Gene</i> , 2019 , 699, 135-144	3.8	9
334	Unraveling UCA1 lncRNA prognostic utility in urothelial bladder cancer. <i>Carcinogenesis</i> , 2019 , 40, 965-974.6	4.6	10

333	Gene-Specific Intron Retention Serves as Molecular Signature that Distinguishes Melanoma from Non-Melanoma Cancer Cells in Greek Patients. <i>International Journal of Molecular Sciences</i> , 2019 , 20,	6.3	5
332	Revisiting Histone Deacetylases in Human Tumorigenesis: The Paradigm of Urothelial Bladder Cancer. <i>International Journal of Molecular Sciences</i> , 2019 , 20,	6.3	27
331	Identification of a novel tRNA-derived RNA fragment exhibiting high prognostic potential in chronic lymphocytic leukemia. <i>Hematological Oncology</i> , 2019 , 37, 498-504	1.3	21
330	Blood-based analysis of type-2 diabetes mellitus susceptibility genes identifies specific transcript variants with deregulated expression and association with disease risk. <i>Scientific Reports</i> , 2019 , 9, 1512	4.9	11
329	Uncovering the clinical impact of kallikrein-related peptidase 5 (KLK5) mRNA expression in the colorectal adenoma-carcinoma sequence. <i>Clinical Chemistry and Laboratory Medicine</i> , 2019 , 57, 1251-1260	5.9	4
328	MicroRNA-92a-3p overexpression in peripheral blood mononuclear cells is an independent predictor of prolonged overall survival of patients with chronic lymphocytic leukemia. <i>Leukemia and Lymphoma</i> , 2019 , 60, 658-667	1.9	13
327	Identification of a novel, internal tRNA-derived RNA fragment as a new prognostic and screening biomarker in chronic lymphocytic leukemia, using an innovative quantitative real-time PCR assay. <i>Leukemia Research</i> , 2019 , 87, 106234	2.7	17
326	The miR-200 family as prognostic markers in clear cell renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2019 , 37, 955-963	2.8	17
325	p63 transcript loss in bladder cancer constitutes an independent molecular predictor of TaT1 patients post-treatment relapse and progression. <i>Journal of Cancer Research and Clinical Oncology</i> , 2019 , 145, 3075-3087	4.9	2
324	Circulating exosomal miRNAs: clinical significance in human cancers. <i>Expert Review of Molecular Diagnostics</i> , 2019 , 19, 979-995	3.8	13
323	Molecular characterization, genomic structure and expression analysis of a gene (CATL1/CPT1C) encoding a third member of the human carnitine acyltransferase family. <i>Genomics</i> , 2019 ,	4.3	1
322	The emergence of drug resistance to targeted cancer therapies: Clinical evidence. <i>Drug Resistance Updates</i> , 2019 , 47, 100646	23.2	48
321	A Molecular Signature of Three tRNA-Derived RNA Fragments May Discriminate Smoldering from Symptomatic Multiple Myeloma Patients. <i>Blood</i> , 2019 , 134, 5528-5528	2.2	1
320	Natural Alkaloids Intervening the Insulin Pathway: New Hopes for Anti-Diabetic Agents?. <i>Current Medicinal Chemistry</i> , 2019 , 26, 5982-6015	4.3	21
319	Discovery of novel transcripts of the human tissue kallikrein (KLK1) and kallikrein-related peptidase 2 (KLK2) in human cancer cells, exploiting Next-Generation Sequencing technology. <i>Genomics</i> , 2019 , 111, 642-652	4.3	12
318	High microRNA-28-5p expression in colorectal adenocarcinoma predicts short-term relapse of node-negative patients and poor overall survival of patients with non-metastatic disease. <i>Clinical Chemistry and Laboratory Medicine</i> , 2018 , 56, 990-1000	5.9	20
317	Manfred Schmitt (1947-2018). <i>Biological Chemistry</i> , 2018 , 399, 923-924	4.5	
316	Kallikrein-related peptidase 6 (KLK6) expression differentiates tumor subtypes and predicts clinical outcome in breast cancer patients. <i>Clinical and Experimental Medicine</i> , 2018 , 18, 203-213	4.9	9

315	Elevated miR-20b-5p expression in peripheral blood mononuclear cells: A novel, independent molecular biomarker of favorable prognosis in chronic lymphocytic leukemia. <i>Leukemia Research</i> , 2018 , 70, 1-7	2.7	21
314	Clinical utility of miR-143/miR-182 levels in prognosis and risk stratification specificity of BFM-treated childhood acute lymphoblastic leukemia. <i>Annals of Hematology</i> , 2018 , 97, 1169-1182	3	14
313	Multianalyte quantitative competitive PCR on optically encoded microspheres for an eight-gene panel related to prostate cancer. <i>Analytical and Bioanalytical Chemistry</i> , 2018 , 410, 971-980	4.4	3
312	A comprehensive clinicopathological evaluation of the differential expression of microRNA-331 in breast tumors and its diagnostic significance. <i>Clinical Biochemistry</i> , 2018 , 60, 24-32	3.5	5
311	Molecular cloning of novel transcripts of the adaptor-related protein complex 2 alpha 1 subunit (AP2A1) gene, using Next-Generation Sequencing. <i>Gene</i> , 2018 , 678, 55-64	3.8	7
310	Expressional profiling and clinical relevance of RNase H in prostate cancer: a novel indicator of favorable progression-free survival. <i>Journal of Cancer Research and Clinical Oncology</i> , 2018 , 144, 2049-2057	4.9	3
309	BCL2L12 improves risk stratification and prediction of BFM-chemotherapy response in childhood acute lymphoblastic leukemia. <i>Clinical Chemistry and Laboratory Medicine</i> , 2018 , 56, 2104-2118	5.9	7
308	Novel splice variants of the human kallikrein-related peptidases 11 (KLK11) and 12 (KLK12), unraveled by next-generation sequencing technology. <i>Biological Chemistry</i> , 2018 , 399, 1065-1071	4.5	10
307	Kallikrein-related peptidases and associated microRNAs as promising prognostic biomarkers in gastrointestinal malignancies. <i>Biological Chemistry</i> , 2018 , 399, 821-836	4.5	9
306	Quantitative analysis and study of the mRNA expression levels of apoptotic genes, and in the articular cartilage of an animal model of osteoarthritis. <i>Annals of Translational Medicine</i> , 2018 , 6, 243	3.2	7
305	Non-coding RNAs: the riddle of the transcriptome and their perspectives in cancer. <i>Annals of Translational Medicine</i> , 2018 , 6, 241	3.2	63
304	Alternative Splicing Detection Tool-a novel PERL algorithm for sensitive detection of splicing events, based on next-generation sequencing data analysis. <i>Annals of Translational Medicine</i> , 2018 , 6, 244	3.2	7
303	Molecular Effects of Treatment of Human Colorectal Cancer Cells with Natural and Classical Chemotherapeutic Drugs: Alterations in the Expression of Apoptosis-related BCL2 Family Members, Including BCL2L12. <i>Current Pharmaceutical Biotechnology</i> , 2018 , 19, 1064-1075	2.6	8
302	Evidence for L-Dopa Decarboxylase Involvement in Cancer Cell Cytotoxicity Induced by Docetaxel and Mitoxantrone. <i>Current Pharmaceutical Biotechnology</i> , 2018 , 19, 1087-1096	2.6	6
301	miR-221/222 cluster expression improves clinical stratification of non-muscle invasive bladder cancer (TaT1) patients: Risk for short-term relapse and progression. <i>Genes Chromosomes and Cancer</i> , 2018 , 57, 150-161	5	18
300	Expression Analysis of miR-29b in Malignant and Benign Breast Tumors: A Promising Prognostic Biomarker for Invasive Ductal Carcinoma With a Possible Histotype-Related Expression Status. <i>Clinical Breast Cancer</i> , 2018 , 18, 305-312.e3	3	5
299	Clinical utility of microRNAs in renal cell carcinoma: current evidence and future perspectives. <i>Expert Review of Molecular Diagnostics</i> , 2018 , 18, 981-991	3.8	13
298	BCL2L12: a multiply spliced gene with independent prognostic significance in breast cancer. <i>Clinical Chemistry and Laboratory Medicine</i> , 2018 , 57, 276-287	5.9	2

297	Loss of GAS5 tumour suppressor lncRNA: an independent molecular cancer biomarker for short-term relapse and progression in bladder cancer patients. <i>British Journal of Cancer</i> , 2018 , 119, 1477-1486	8.7	29
296	miRNA and long non-coding RNA: molecular function and clinical value in breast and ovarian cancers. <i>Expert Review of Molecular Diagnostics</i> , 2018 , 18, 963-979	3.8	30
295	Discovery and expression analysis of novel transcripts of the human SR-related CTD-associated factor 1 (SCAF1) gene in human cancer cells using Next-Generation Sequencing. <i>Gene</i> , 2018 , 670, 155-165	3.8	8
294	Human kallikrein-related peptidase 12 (KLK12) splice variants discriminate benign from cancerous breast tumors. <i>Clinical Biochemistry</i> , 2018 , 58, 78-85	3.5	8
293	miR-15a-5p, A Novel Prognostic Biomarker, Predicting Recurrent Colorectal Adenocarcinoma. <i>Molecular Diagnosis and Therapy</i> , 2017 , 21, 453-464	4.5	33
292	Identification and molecular cloning of novel transcripts of the human kallikrein-related peptidase 10 (KLK10) gene using next-generation sequencing. <i>Biochemical and Biophysical Research Communications</i> , 2017 , 487, 776-781	3.4	14
291	mRNA overexpression of the hypoxia inducible factor 1 alpha subunit gene (HIF1A): An independent predictor of poor overall survival in chronic lymphocytic leukemia. <i>Leukemia Research</i> , 2017 , 53, 65-73	2.7	20
290	Unravelling a p73-regulated network: The role of a novel p73-dependent target, MIR3158, in cancer cell migration and invasiveness. <i>Cancer Letters</i> , 2017 , 388, 96-106	9.9	13
289	miR-34a overexpression predicts poor prognostic outcome in colorectal adenocarcinoma, independently of clinicopathological factors with established prognostic value. <i>Clinical Biochemistry</i> , 2017 , 50, 918-924	3.5	18
288	Comparative HPLC-DAD and UHPLC-ESI(-)-HRMS & MS/MS profiling of Hypericum species and correlation with necrotic cell-death activity in human leukemic cells. <i>Phytochemistry Letters</i> , 2017 , 20, 481-490	1.9	7
287	miR-10b is a prognostic marker in clear cell renal cell carcinoma. <i>Journal of Clinical Pathology</i> , 2017 , 70, 854-859	3.9	23
286	Elevated expression of miR-24-3p is a potentially adverse prognostic factor in colorectal adenocarcinoma. <i>Clinical Biochemistry</i> , 2017 , 50, 285-292	3.5	27
285	Upregulated miR-16 expression is an independent indicator of relapse and poor overall survival of colorectal adenocarcinoma patients. <i>Clinical Chemistry and Laboratory Medicine</i> , 2017 , 55, 737-747	5.9	25
284	MicroRNA-155-5p Overexpression in Peripheral Blood Mononuclear Cells of Chronic Lymphocytic Leukemia Patients Is a Novel, Independent Molecular Biomarker of Poor Prognosis. <i>Disease Markers</i> , 2017 , 2017, 2046545	3.2	22
283	Molecular cloning of novel transcripts of human kallikrein-related peptidases 5, 6, 7, 8 and 9 (KLK5 - KLK9), using Next-generation sequencing. <i>Scientific Reports</i> , 2017 , 7, 17299	4.9	15
282	Assessing the clinical value of microRNAs in formalin-fixed paraffin-embedded liposarcoma tissues: Overexpressed miR-155 is an indicator of poor prognosis. <i>Oncotarget</i> , 2017 , 8, 6896-6913	3.3	13
281	miR-125b predicts childhood acute lymphoblastic leukaemia poor response to BFM chemotherapy treatment. <i>British Journal of Cancer</i> , 2017 , 117, 801-812	8.7	21
280	The transcriptome of a "sleeping" invader: de novo assembly and annotation of the transcriptome of aestivating <i>Cornu aspersum</i> . <i>BMC Genomics</i> , 2017 , 18, 491	4.5	14

279	Downregulated KLK13 expression in bladder cancer highlights tumor aggressiveness and unfavorable patients prognosis. <i>Journal of Cancer Research and Clinical Oncology</i> , 2017 , 143, 521-532	4.9	14
278	Metformin and Anti-Cancer Therapeutics: Hopes for a More Enhanced Armamentarium Against Human Neoplasias?. <i>Current Medicinal Chemistry</i> , 2017 , 24, 14-56	4.3	5
277	Pediatric Ependymoma: A Proteomics Perspective. <i>Cancer Genomics and Proteomics</i> , 2017 , 14, 127-136	3.3	6
276	S100A11 is a potential prognostic marker for clear cell renal cell carcinoma. <i>Clinical and Experimental Metastasis</i> , 2016 , 33, 63-71	4.7	12
275	BCL2L12 protein overexpression is associated with favorable outcome in diffuse large B-cell lymphoma patients in the rituximab era. <i>Leukemia and Lymphoma</i> , 2016 , 57, 2199-203	1.9	9
274	Copper(II) Inverse-[9-Metallacrown-3] Compounds Accommodating [Nitrate or Diclofenac Ligands: Structure, Magnetism, and Biological Activity. <i>European Journal of Inorganic Chemistry</i> , 2016 , 2016, 219-231	2.3	23
273	Identification of novel alternative splice variants of the BCL2L12 gene in human cancer cells using next-generation sequencing methodology. <i>Cancer Letters</i> , 2016 , 373, 119-129	9.9	24
272	MicroRNA-194 is a Marker for Good Prognosis in Clear Cell Renal Cell Carcinoma. <i>Cancer Medicine</i> , 2016 , 5, 656-64	4.8	37
271	Clinical evaluation of microRNA-145 expression in renal cell carcinoma: a promising molecular marker for discriminating and staging the clear cell histological subtype. <i>Biological Chemistry</i> , 2016 , 397, 529-39	4.5	16
270	Kallikrein-related peptidases (KLKs) as emerging therapeutic targets: focus on prostate cancer and skin pathologies. <i>Expert Opinion on Therapeutic Targets</i> , 2016 , 20, 801-18	6.4	23
269	The Stat3/5 Signaling Biosignature in Hematopoietic Stem/Progenitor Cells Predicts Response and Outcome in Myelodysplastic Syndrome Patients Treated with Azacitidine. <i>Clinical Cancer Research</i> , 2016 , 22, 1958-68	12.9	16
268	mRNA overexpression of kallikrein-related peptidase 14 (KLK14) is an independent predictor of poor overall survival in chronic lymphocytic leukemia patients. <i>Clinical Chemistry and Laboratory Medicine</i> , 2016 , 54, 315-24	5.9	11
267	Evaluation of PD-L1 Expression and Associated Tumor-Infiltrating Lymphocytes in Laryngeal Squamous Cell Carcinoma. <i>Clinical Cancer Research</i> , 2016 , 22, 704-13	12.9	138
266	Downregulation of the neonatal Fc receptor expression in non-small cell lung cancer tissue is associated with a poor prognosis. <i>Oncotarget</i> , 2016 , 7, 54415-54429	3.3	20
265	Progression of mouse skin carcinogenesis is associated with the orchestrated deregulation of mir-200 family members, mir-205 and their common targets. <i>Molecular Carcinogenesis</i> , 2016 , 55, 1229-42 ⁵		18
264	High miR-96 levels in colorectal adenocarcinoma predict poor prognosis, particularly in patients without distant metastasis at the time of initial diagnosis. <i>Tumor Biology</i> , 2016 , 37, 11815-11824	2.9	36
263	High BAX/BCL2 mRNA ratio predicts favorable prognosis in laryngeal squamous cell carcinoma, particularly in patients with negative lymph nodes at the time of diagnosis. <i>Clinical Biochemistry</i> , 2016 , 49, 890-6	3.5	25
262	L-DOPA decarboxylase mRNA levels provide high diagnostic accuracy and discrimination between clear cell and non-clear cell subtypes in renal cell carcinoma. <i>Clinical Biochemistry</i> , 2015 , 48, 590-5	3.5	7

261	Profilin-1 expression is associated with high grade and stage and decreased disease-free survival in renal cell carcinoma. <i>Human Pathology</i> , 2015 , 46, 673-80	3.7	19
260	Kallikreins as Biomarkers in Human Malignancies. <i>Biomarkers in Disease</i> , 2015 , 135-165		2
259	Clinical relevance of the deregulated kallikrein-related peptidase 8 mRNA expression in breast cancer: a novel independent indicator of disease-free survival. <i>Breast Cancer Research and Treatment</i> , 2015 , 152, 323-36	4.4	11
258	The oncomiR miR-197 is a novel prognostic indicator for non-small cell lung cancer patients. <i>British Journal of Cancer</i> , 2015 , 112, 1527-35	8.7	39
257	KLKB1 mRNA overexpression: A novel molecular biomarker for the diagnosis of chronic lymphocytic leukemia. <i>Clinical Biochemistry</i> , 2015 , 48, 849-54	3.5	23
256	Uncovering the clinical utility of miR-143, miR-145 and miR-224 for predicting the survival of bladder cancer patients following treatment. <i>Carcinogenesis</i> , 2015 , 36, 528-37	4.6	56
255	Gemcitabine impacts differentially on bladder and kidney cancer cells: distinct modulations in the expression patterns of apoptosis-related microRNAs and BCL2 family genes. <i>Tumor Biology</i> , 2015 , 36, 3197-207	2.9	9
254	Prognostic and predictive biomarkers in prostate cancer. <i>Expert Review of Molecular Diagnostics</i> , 2015 , 15, 1567-76	3.8	23
253	Human L-DOPA decarboxylase mRNA is a target of miR-145: A prediction to validation workflow. <i>Gene</i> , 2015 , 554, 174-80	3.8	4
252	Quantitative analysis of the mRNA expression levels of BCL2 and BAX genes in human osteoarthritis and normal articular cartilage: An investigation into their differential expression. <i>Molecular Medicine Reports</i> , 2015 , 12, 4514-4521	2.9	31
251	Long Noncoding RNAs in Digestive System Malignancies: A Novel Class of Cancer Biomarkers and Therapeutic Targets?. <i>Gastroenterology Research and Practice</i> , 2015 , 2015, 319861	2	22
250	Alpha-enolase is a potential prognostic marker in clear cell renal cell carcinoma. <i>Clinical and Experimental Metastasis</i> , 2015 , 32, 531-41	4.7	14
249	Overexpression of BCL2 and BAX following BFM induction therapy predicts ch-ALL patients poor response to treatment and short-term relapse. <i>Journal of Cancer Research and Clinical Oncology</i> , 2015 , 141, 2023-36	4.9	9
248	Cisplatin and Paclitaxel Alter the Expression Pattern of miR-143/145 and miR-183/96/182 Clusters in T24 Bladder Cancer Cells. <i>Clinical and Translational Science</i> , 2015 , 8, 668-75	4.9	5
247	miR-224 overexpression is a strong and independent prognosticator of short-term relapse and poor overall survival in colorectal adenocarcinoma. <i>International Journal of Oncology</i> , 2015 , 46, 849-59	4.4	32
246	miR-210 is a prognostic marker in clear cell renal cell carcinoma. <i>Journal of Molecular Diagnostics</i> , 2015 , 17, 136-44	5.1	49
245	Cytotoxic activity of sunitinib and everolimus in Caki-1 renal cancer cells is accompanied by modulations in the expression of apoptosis-related microRNA clusters and BCL2 family genes. <i>Biomedicine and Pharmacotherapy</i> , 2015 , 70, 33-40	7.5	17
244	Kallikrein-related peptidase 13: an independent indicator of favorable prognosis for patients with nonsmall cell lung cancer. <i>Tumor Biology</i> , 2015 , 36, 4979-86	2.9	13

243	Low expression of miR-126 is a prognostic marker for metastatic clear cell renal cell carcinoma. <i>American Journal of Pathology</i> , 2015 , 185, 693-703	5.8	51
242	Prognostic role and implications of mutation status of tumor suppressor gene ARID1A in cancer: a systematic review and meta-analysis. <i>Oncotarget</i> , 2015 , 6, 39088-97	3.3	49
241	Molecular Biomarkers of Laryngeal Cancer. <i>Biomarkers in Disease</i> , 2015 , 891-919		4
240	Targeting kallikrein-related peptidases in prostate cancer. <i>Expert Opinion on Therapeutic Targets</i> , 2014 , 18, 365-83	6.4	21
239	Kallikrein-related peptidase-6 (KLK6) mRNA expression is an independent prognostic tissue biomarker of poor disease-free and overall survival in colorectal adenocarcinoma. <i>Tumor Biology</i> , 2014 , 35, 4673-85	2.9	28
238	Kallikrein-related peptidase 6 (KLK6) expression in the progression of colon adenoma to carcinoma. <i>Biological Chemistry</i> , 2014 , 395, 1105-17	4.5	16
237	Galectin-1 has potential prognostic significance and is implicated in clear cell renal cell carcinoma progression through the HIF/mTOR signaling axis. <i>British Journal of Cancer</i> , 2014 , 110, 1250-9	8.7	43
236	KLK11 mRNA expression predicts poor disease-free and overall survival in colorectal adenocarcinoma patients. <i>Biomarkers in Medicine</i> , 2014 , 8, 671-85	2.3	23
235	Loss of miR-378 in prostate cancer, a common regulator of KLK2 and KLK4, correlates with aggressive disease phenotype and predicts the short-term relapse of the patients. <i>Biological Chemistry</i> , 2014 , 395, 1095-104	4.5	27
234	A new tumor suppressor role for the Notch pathway in bladder cancer. <i>Nature Medicine</i> , 2014 , 20, 1199-2005	30.5	130
233	BCL2L12: a promising molecular prognostic biomarker in breast cancer. <i>Clinical Biochemistry</i> , 2014 , 47, 257-62	3.5	16
232	Low mRNA expression levels of kallikrein-related peptidase 4 (KLK4) predict short-term relapse in patients with laryngeal squamous cell carcinoma. <i>Biological Chemistry</i> , 2014 , 395, 1051-62	4.5	12
231	Lactate dehydrogenase A is a potential prognostic marker in clear cell renal cell carcinoma. <i>Molecular Cancer</i> , 2014 , 13, 101	42.1	101
230	Predictions for the future of kallikrein-related peptidases in molecular diagnostics. <i>Expert Review of Molecular Diagnostics</i> , 2014 , 14, 713-22	3.8	16
229	RAS/PI3K crosstalk and cetuximab resistance in head and neck squamous cell carcinoma. <i>Clinical Cancer Research</i> , 2014 , 20, 2933-46	12.9	55
228	Increased BCL2L12 expression predicts the short-term relapse of patients with TaT1 bladder cancer following transurethral resection of bladder tumors. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014 , 32, 39.e29-36	2.8	8
227	Translation Regulation by microRNAs in Acute Leukemia 2014 , 1-30		
226	Enhanced miR-182 transcription is a predictor of poor overall survival in colorectal adenocarcinoma patients. <i>Clinical Chemistry and Laboratory Medicine</i> , 2014 , 52, 1217-27	5.9	31

225	A comprehensive phylogenetic and structural analysis of the carcinoembryonic antigen (CEA) gene family. <i>Genome Biology and Evolution</i> , 2014 , 6, 1314-26	3.9	20
224	Nature promises new anticancer agents: Interplay with the apoptosis-related BCL2 gene family. <i>Anti-Cancer Agents in Medicinal Chemistry</i> , 2014 , 14, 375-99	2.2	26
223	Apoptosis-related BCL2-family members: Key players in chemotherapy. <i>Anti-Cancer Agents in Medicinal Chemistry</i> , 2014 , 14, 353-74	2.2	63
222	Molecular Biomarkers of Laryngeal Cancer 2014 , 1-24		1
221	Association of BCL2L12 overexpression with prolonged disease-free survival in breast cancer.. <i>Journal of Clinical Oncology</i> , 2014 , 32, e22202-e22202	2.2	
220	Kallikreins as Biomarkers in Human Malignancies 2014 , 1-25		1
219	Quantitative expression analysis of the apoptosis-related gene, BCL2L12, in head and neck squamous cell carcinoma. <i>Journal of Oral Pathology and Medicine</i> , 2013 , 42, 154-61	3.3	21
218	Quantitative expression analysis and prognostic significance of the BCL2-associated X gene in nasopharyngeal carcinoma: a retrospective cohort study. <i>BMC Cancer</i> , 2013 , 13, 293	4.8	21
217	Clinical significance of kallikrein-related peptidase (KLK10) mRNA expression in colorectal cancer. <i>Clinical Biochemistry</i> , 2013 , 46, 1453-61	3.5	37
216	The role of transcription factors in laboratory medicine. <i>Clinical Chemistry and Laboratory Medicine</i> , 2013 , 51, 1563-71	5.9	9
215	The chromatin remodeling gene ARID1A is a new prognostic marker in clear cell renal cell carcinoma. <i>American Journal of Pathology</i> , 2013 , 182, 1163-70	5.8	55
214	Significant alterations in the expression pattern of kallikrein-related peptidase genes KLK4, KLK5 and KLK14 after treatment of breast cancer cells with the chemotherapeutic agents epirubicin, docetaxel and methotrexate. <i>Tumor Biology</i> , 2013 , 34, 369-78	2.9	7
213	Kallikrein-related peptidase 4 (KLK4) mRNA predicts short-term relapse in colorectal adenocarcinoma patients. <i>Cancer Letters</i> , 2013 , 330, 106-12	9.9	28
212	L-Dopa decarboxylase (DDC) constitutes an emerging biomarker in predicting patients' survival with stomach adenocarcinomas. <i>Journal of Cancer Research and Clinical Oncology</i> , 2013 , 139, 297-306	4.9	6
211	Expression of Bcl2L12 in chronic lymphocytic leukemia patients: association with clinical and molecular prognostic markers. <i>Medical Oncology</i> , 2013 , 30, 405	3.7	13
210	Downregulation and prognostic performance of microRNA 224 expression in prostate cancer. <i>Clinical Chemistry</i> , 2013 , 59, 261-9	5.5	55
209	The expression of the CEACAM19 gene, a novel member of the CEA family, is associated with breast cancer progression. <i>International Journal of Oncology</i> , 2013 , 42, 1770-7	4.4	35
208	The loss of the tumour-suppressor miR-145 results in the shorter disease-free survival of prostate cancer patients. <i>British Journal of Cancer</i> , 2013 , 108, 2573-81	8.7	74

207	Quantified KLK15 gene expression levels discriminate prostate cancer from benign tumors and constitute a novel independent predictor of disease progression. <i>Prostate</i> , 2013 , 73, 1191-201	4.2	14
206	Emerging clinical importance of the cancer biomarkers kallikrein-related peptidases (KLK) in female and male reproductive organ malignancies. <i>Radiology and Oncology</i> , 2013 , 47, 319-29	3.8	33
205	Effect of doxorubicin, oxaliplatin, and methotrexate administration on the transcriptional activity of BCL-2 family gene members in stomach cancer cells. <i>Cancer Biology and Therapy</i> , 2013 , 14, 587-96	4.6	20
204	Kallikrein-related peptidases (KLKs) in gastrointestinal cancer: mechanistic and clinical aspects. <i>Thrombosis and Haemostasis</i> , 2013 , 110, 450-7	7	30
203	Quantification and study of the L-DOPA decarboxylase expression in gastric adenocarcinoma cells treated with chemotherapeutic substances. <i>Anti-Cancer Drugs</i> , 2013 , 24, 291-9	2.4	2
202	Parallel overexpression and clinical significance of kallikrein-related peptidases 7 and 14 (KLK7KLK14) in colon cancer. <i>Thrombosis and Haemostasis</i> , 2013 , 109, 716-25	7	22
201	Evolution of the plasma and tissue kallikreins, and their alternative splicing isoforms. <i>PLoS ONE</i> , 2013 , 8, e68074	3.7	31
200	Identification of a STAT5 target gene, Dpf3, provides novel insights in chronic lymphocytic leukemia. <i>PLoS ONE</i> , 2013 , 8, e76155	3.7	11
199	Immunohistochemical expression of somatostatin receptor subtypes 2 and 5 in colorectal cancer. <i>European Journal of Clinical Investigation</i> , 2012 , 42, 777-83	4.6	6
198	L-dopa decarboxylase (DDC) gene expression is related to outcome in patients with prostate cancer. <i>BJU International</i> , 2012 , 110, E267-73	5.6	16
197	Kallikrein-related peptidases (KLKs): a gene family of novel cancer biomarkers. <i>Clinical Chemistry and Laboratory Medicine</i> , 2012 , 50, 1877-91	5.9	60
196	Diagnostic and prognostic significance of human kallikrein 11 (KLK11) mRNA expression levels in patients with laryngeal cancer. <i>Clinical Biochemistry</i> , 2012 , 45, 623-30	3.5	17
195	Overexpression of the novel member of the BCL2 gene family, BCL2L12, is associated with the disease outcome in patients with acute myeloid leukemia. <i>Clinical Biochemistry</i> , 2012 , 45, 1362-7	3.5	9
194	Human kallikrein-related peptidase 12 (KLK12) splice variants expression in breast cancer and their clinical impact. <i>Tumor Biology</i> , 2012 , 33, 1075-84	2.9	20
193	Evaluation and prognostic significance of human tissue kallikrein-related peptidase 10 (KLK10) in colorectal cancer. <i>Tumor Biology</i> , 2012 , 33, 1209-14	2.9	12
192	Kallikrein-related peptidase 6 (KLK6) gene expression in intracranial tumors. <i>Tumor Biology</i> , 2012 , 33, 1375-83	2.9	12
191	The single nucleotide polymorphism g.1548A >G (K469E) of the ICAM-1 gene is associated with worse prognosis in non-small cell lung cancer. <i>Tumor Biology</i> , 2012 , 33, 1429-36	2.9	18
190	The kallikrein-related peptidase 13 (KLK13) gene is substantially up-regulated after exposure of gastric cancer cells to antineoplastic agents. <i>Tumor Biology</i> , 2012 , 33, 2069-78	2.9	9

189	Evaluation and prognostic significance of human tissue kallikrein-related peptidase 6 (KLK6) in colorectal cancer. <i>Pathology Research and Practice</i> , 2012 , 208, 104-8	3.4	22
188	Assessment of the prognostic significance of endoglin (CD105) in clear cell renal cell carcinoma using automated image analysis. <i>Human Pathology</i> , 2012 , 43, 1037-43	3.7	22
187	Molecular cloning of novel alternatively spliced variants of BCL2L12, a new member of the BCL2 gene family, and their expression analysis in cancer cells. <i>Gene</i> , 2012 , 505, 153-66	3.8	28
186	L-DOPA decarboxylase mRNA expression is associated with tumor stage and size in head and neck squamous cell carcinoma: a retrospective cohort study. <i>BMC Cancer</i> , 2012 , 12, 484	4.8	17
185	l-DOPA Decarboxylase (DDC) Expression Status as a Novel Molecular Tumor Marker for Diagnostic and Prognostic Purposes in Laryngeal Cancer. <i>Translational Oncology</i> , 2012 , 5, 288-96	4.9	14
184	The clinical utility of miR-21 as a diagnostic and prognostic marker for renal cell carcinoma. <i>Journal of Molecular Diagnostics</i> , 2012 , 14, 385-92	5.1	88
183	DDC (dopa decarboxylase (aromatic L-amino acid decarboxylase)). <i>Atlas of Genetics and Cytogenetics in Oncology and Haematology</i> , 2012 ,	2.3	1
182	Quantitative expression analysis of the apoptosis-related genes BCL2, BAX and BCL2L12 in gastric adenocarcinoma cells following treatment with the anticancer drugs cisplatin, etoposide and taxol. <i>Tumor Biology</i> , 2012 , 33, 865-75	2.9	27
181	Microvascular density as an independent predictor of clinical outcome in renal cell carcinoma: an automated image analysis study. <i>Laboratory Investigation</i> , 2012 , 92, 46-56	5.9	45
180	Kallikrein-related peptidases in prostate, breast, and ovarian cancers: from pathobiology to clinical relevance. <i>Biological Chemistry</i> , 2012 , 393, 301-17	4.5	62
179	The Role of BCL2 Family of Apoptosis Regulator Proteins in Acute and Chronic Leukemias. <i>Advances in Hematology</i> , 2012 , 2012, 524308	1.5	129
178	Evaluation of the clinical utility of kallikrein-related peptidase 6 gene (KLK6) downregulation in breast cancer.. <i>Journal of Clinical Oncology</i> , 2012 , 30, 10606-10606	2.2	1
177	BCL2L12 is a novel biomarker for the prediction of short-term relapse in nasopharyngeal carcinoma. <i>Molecular Medicine</i> , 2011 , 17, 163-71	6.2	36
176	Quantitative expression analysis and study of the novel human kallikrein-related peptidase 14 gene (KLK14) in malignant and benign breast tissues. <i>Thrombosis and Haemostasis</i> , 2011 , 105, 131-7	7	7
175	Pancreatic duct guidewire placement for biliary cannulation in a single-session therapeutic ERCP. <i>World Journal of Gastroenterology</i> , 2011 , 17, 1989-95	5.6	24
174	Comparative study of balloon and metal olive dilators for endoscopic management of benign anastomotic rectal strictures: clinical and cost-effectiveness outcomes. <i>Surgical Endoscopy and Other Interventional Techniques</i> , 2011 , 25, 756-63	5.2	11
173	Clinical evaluation of PRMT1 gene expression in breast cancer. <i>Tumor Biology</i> , 2011 , 32, 575-82	2.9	57
172	Expression analysis and clinical evaluation of kallikrein-related peptidase 10 (KLK10) in colorectal cancer. <i>Tumor Biology</i> , 2011 , 32, 737-44	2.9	27

171	Down-regulation of kallikrein-related peptidase 5 (KLK5) expression in breast cancer patients: a biomarker for the differential diagnosis of breast lesions. <i>Clinical Proteomics</i> , 2011 , 8, 5	5	19
170	Kallikrein-related peptidase 4 gene (KLK4) in prostate tumors: quantitative expression analysis and evaluation of its clinical significance. <i>Prostate</i> , 2011 , 71, 1780-9	4.2	21
169	Synthesis, spectroscopic study and anticancer activity of a water-soluble Nb(V) peroxy complex. <i>Journal of Inorganic Biochemistry</i> , 2011 , 105, 155-63	4.2	16
168	Impact of expression differences of kallikrein-related peptidases and of uPA and PAI-1 between primary tumor and omentum metastasis in advanced ovarian cancer. <i>Annals of Oncology</i> , 2011 , 22, 877-883	10.3	34
167	The novel member of the BCL2 gene family, BCL2L12, is substantially elevated in chronic lymphocytic leukemia patients, supporting its value as a significant biomarker. <i>Oncologist</i> , 2011 , 16, 1280-91	5.7	34
166	Evaluation of kallikrein-related peptidase 5 expression and its significance for breast cancer patients: association with kallikrein-related peptidase 7 expression. <i>Anticancer Research</i> , 2011 , 31, 3093-100	10	7
165	Expression analysis and study of the KLK15 mRNA splice variants in prostate cancer and benign prostatic hyperplasia. <i>Cancer Science</i> , 2010 , 101, 693-9	6.9	17
164	Quantitative expression analysis and prognostic significance of L-DOPA decarboxylase in colorectal adenocarcinoma. <i>British Journal of Cancer</i> , 2010 , 102, 1384-90	8.7	42
163	KLK5 gene expression is severely upregulated in androgen-independent prostate cancer cells after treatment with the chemotherapeutic agents docetaxel and mitoxantrone. <i>Biological Chemistry</i> , 2010 , 391, 467-74	4.5	12
162	Prognostic value and biological role of the kallikrein-related peptidases in human malignancies. <i>Future Oncology</i> , 2010 , 6, 269-85	3.6	49
161	The miR-17-92 cluster is over expressed in and has an oncogenic effect on renal cell carcinoma. <i>Journal of Urology</i> , 2010 , 183, 743-51	2.5	128
160	Molecular analysis and prognostic impact of the novel apoptotic gene BCL2L12 in gastric cancer. <i>Biochemical and Biophysical Research Communications</i> , 2010 , 391, 214-8	3.4	29
159	Kallikrein-related peptidase genes as promising biomarkers for prognosis and monitoring of human malignancies. <i>Biological Chemistry</i> , 2010 , 391, 505-11	4.5	58
158	Quantitative analysis of BCL2 mRNA expression in nasopharyngeal carcinoma: an unfavorable and independent prognostic factor. <i>Tumor Biology</i> , 2010 , 31, 391-9	2.9	26
157	Kallikrein-related peptidase 13 (KLK13) gene expressional status contributes significantly in the prognosis of primary gastric carcinomas. <i>Clinical Biochemistry</i> , 2010 , 43, 1205-11	3.5	21
156	Expression analysis and study of KLK4 in benign and malignant breast tumours. <i>Thrombosis and Haemostasis</i> , 2009 , 101, 381-387	7	22
155	The immunohistochemical expression of growth hormone-releasing hormone receptor splice variant 1 is a favorable prognostic marker in colorectal cancer. <i>Molecular Medicine</i> , 2009 , 15, 242-7	6.2	5
154	The use of kallikrein-related peptidases as adjuvant prognostic markers in colorectal cancer. <i>British Journal of Cancer</i> , 2009 , 100, 1659-65	8.7	53

153	Quantitative analysis of human kallikrein 5 (KLK5) expression in prostate needle biopsies: an independent cancer biomarker. <i>Clinical Chemistry</i> , 2009 , 55, 904-13	5.5	19
152	Association between kallikrein-related peptidases (KLKs) and macroscopic indicators of semen analysis: their relation to sperm motility. <i>Biological Chemistry</i> , 2009 , 390, 921-9	4.5	17
151	Phosphatidylinositol 3-kinase catalytic subunit alpha gene amplification contributes to the pathogenesis of mantle cell lymphoma. <i>Clinical Cancer Research</i> , 2009 , 15, 5724-32	12.9	83
150	Cathepsin B protein levels in endometrial cancer: Potential value as a tumour biomarker. <i>Gynecologic Oncology</i> , 2009 , 112, 531-6	4.9	20
149	Molecular response of HL-60 cells to mitotic inhibitors vincristine and taxol visualized with apoptosis-related gene expressions, including the new member BCL2L12. <i>Annals of the New York Academy of Sciences</i> , 2009 , 1171, 276-83	6.5	16
148	Differential protein expressions in renal cell carcinoma: new biomarker discovery by mass spectrometry. <i>Journal of Proteome Research</i> , 2009 , 8, 3797-807	5.6	66
147	Treatment of gastric cancer cells with 5-fluorouracil/leucovorin and irinotecan induces distinct alterations in the mRNA expression of the apoptosis-related genes, including the novel gene BCL2L12. <i>Tumor Biology</i> , 2009 , 30, 100-7	2.9	14
146	Clinical significance of kallikrein-related peptidase 7 (KLK7) in colorectal cancer. <i>Thrombosis and Haemostasis</i> , 2009 , 101, 741-747	7	32
145	Treatment of PC3 prostate cancer cells with mitoxantrone, etoposide, doxorubicin and carboplatin induces distinct alterations in the expression of kallikreins 5 and 11. <i>Thrombosis and Haemostasis</i> , 2009 , 101, 373-380	7	17
144	Expression and prognostic significance of kallikrein-related peptidase 8 protein levels in advanced ovarian cancer by using automated quantitative analysis. <i>Thrombosis and Haemostasis</i> , 2009 , 101, 541-546	7	16
143	Treatment of PC3 prostate cancer cells with mitoxantrone, etoposide, doxorubicin and carboplatin induces distinct alterations in the expression of kallikreins 5 and 11. <i>Thrombosis and Haemostasis</i> , 2009 , 101, 373-80	7	6
142	Expression analysis and study of KLK4 in benign and malignant breast tumours. <i>Thrombosis and Haemostasis</i> , 2009 , 101, 381-7	7	7
141	Expression and prognostic significance of kallikrein-related peptidase 8 protein levels in advanced ovarian cancer by using automated quantitative analysis. <i>Thrombosis and Haemostasis</i> , 2009 , 101, 541-6	7	9
140	Clinical significance of kallikrein-related peptidase 7 (KLK7) in colorectal cancer. <i>Thrombosis and Haemostasis</i> , 2009 , 101, 741-7	7	18
139	The PRMT1 gene expression pattern in colon cancer. <i>British Journal of Cancer</i> , 2008 , 99, 2094-9	8.7	93
138	Prognostic value of kallikrein-related peptidase 6 protein expression levels in advanced ovarian cancer evaluated by automated quantitative analysis (AQUA). <i>Cancer Science</i> , 2008 , 99, 2224-9	6.9	20
137	Molecular profile of breast versus ovarian cancer cells in response to treatment with the anticancer drugs cisplatin, carboplatin, doxorubicin, etoposide and taxol. <i>Biological Chemistry</i> , 2008 , 389, 1427-34	4.5	19
136	Molecular profile of the BCL2 family of the apoptosis related genes in breast cancer cells after treatment with cytotoxic/cytostatic drugs. <i>Connective Tissue Research</i> , 2008 , 49, 261-4	3.3	29

135	Human tissue kallikrein 7, a novel biomarker for advanced ovarian carcinoma using a novel in situ quantitative method of protein expression. <i>Annals of Oncology</i> , 2008 , 19, 1271-1277	10.3	23
134	Quantitative expression analysis and prognostic significance of the novel apoptosis-related gene BCL2L12 in colon cancer. <i>Biological Chemistry</i> , 2008 , 389, 1467-75	4.5	36
133	Trastuzumab plus paclitaxel or docetaxel in HER-2-negative/HER-2 ECD-positive anthracycline- and taxane-refractory advanced breast cancer. <i>Oncologist</i> , 2008 , 13, 361-9	5.7	22
132	The role of cordycepin in cancer treatment via induction or inhibition of apoptosis: implication of polyadenylation in a cell type specific manner. <i>Cancer Chemotherapy and Pharmacology</i> , 2008 , 61, 251-65 ³⁻⁵	3.5	39
131	Expression analysis and clinical utility of L-Dopa decarboxylase (DDC) in prostate cancer. <i>Clinical Biochemistry</i> , 2008 , 41, 1140-9	3.5	33
130	The effect of the polyadenylation inhibitor cordycepin on human Molt-4 and Daudi leukaemia and lymphoma cell lines. <i>Cancer Chemotherapy and Pharmacology</i> , 2008 , 61, 703-11	3.5	24
129	Total and free PSA kinetics in patients without prostate cancer undergoing radical cystoprostatectomy. <i>Prostate</i> , 2008 , 68, 759-65	4.2	4
128	Novel splice variants of prostate-specific antigen and applications in diagnosis of prostate cancer. <i>Clinical Biochemistry</i> , 2008 , 41, 591-7	3.5	15
127	Enhanced concentration-dependent cytotoxic effect of the dinuclear copper(II) complex of L-carnitine [Cu ₂ (L-carnitine) ₂ Cl ₂ (H ₂ O) ₂]Cl ₂ , compared to L-carnitine or copper chloride dihydrate, in human leukemic cell lines. <i>Journal of Medicinal Chemistry</i> , 2008 , 51, 3713-9	8.3	12
126	Enhanced antileukemic activity of the novel complex 2,5-dihydroxybenzoate molybdenum(VI) against 2,5-dihydroxybenzoate, polyoxometalate of Mo(VI), and tetraphenylphosphonium in the human HL-60 and K562 leukemic cell lines. <i>Journal of Medicinal Chemistry</i> , 2007 , 50, 1316-21	8.3	24
125	Structure and biological properties of the copper(II) complex with the quinolone antibacterial drug N-propyl-norfloxacin and 2,2'-bipyridine. <i>Journal of Inorganic Biochemistry</i> , 2007 , 101, 64-73	4.2	126
124	Transcriptional upregulation of human tissue kallikrein 6 in ovarian cancer: clinical and mechanistic aspects. <i>British Journal of Cancer</i> , 2007 , 96, 362-72	8.7	38
123	B7-H4 is over-expressed in early-stage ovarian cancer and is independent of CA125 expression. <i>Gynecologic Oncology</i> , 2007 , 106, 334-41	4.9	58
122	Breast cancer cells response to the antineoplastic agents cisplatin, carboplatin, and doxorubicin at the mRNA expression levels of distinct apoptosis-related genes, including the new member, BCL2L12. <i>Annals of the New York Academy of Sciences</i> , 2007 , 1095, 35-44	6.5	27
121	Effect of bleomycin and cisplatin on the expression profile of SRA1, a novel member of pre-mRNA splicing factors, in HL-60 human promyelocytic leukemia cells. <i>Biological Chemistry</i> , 2007 , 388, 773-8	4.5	1
120	A multiparametric panel for ovarian cancer diagnosis, prognosis, and response to chemotherapy. <i>Clinical Cancer Research</i> , 2007 , 13, 6984-92	12.9	57
119	Primary tumor levels of human tissue kallikreins affect surgical success and survival in ovarian cancer patients. <i>Clinical Cancer Research</i> , 2007 , 13, 1742-8	12.9	23
118	Prognostic value of the apoptosis related genes BCL2 and BCL2L12 in breast cancer. <i>Cancer Letters</i> , 2007 , 247, 48-55	9.9	52

117	Correlation of androgen receptor status, neuroendocrine differentiation and angiogenesis with time-to-biochemical failure after radical prostatectomy in clinically localized prostate cancer. <i>Anticancer Research</i> , 2007 , 27, 3651-60	2.3	28
116	Kallikreins as markers of disseminated tumour cells in ovarian cancer-- a pilot study. <i>Tumor Biology</i> , 2006 , 27, 104-14	2.9	23
115	Human kallikrein 8 protein is a favorable prognostic marker in ovarian cancer. <i>Clinical Cancer Research</i> , 2006 , 12, 1487-93	12.9	47
114	Effect of testosterone administration on serum and urine kallikrein concentrations in female-to-male transsexuals. <i>Clinical Chemistry</i> , 2006 , 52, 1546-51	5.5	13
113	Multidisciplinary therapy of locally far-advanced or inflammatory breast cancer with fixed perioperative sequence of epirubicin, vinorelbine, and Fluorouracil chemotherapy, surgery, and radiotherapy: long-term results. <i>Oncologist</i> , 2006 , 11, 563-73	5.7	7
112	The role of human tissue kallikreins 7 and 8 in intracranial malignancies. <i>Biological Chemistry</i> , 2006 , 387, 1607-12	4.5	24
111	Unfavorable prognostic value of human kallikrein 7 quantified by ELISA in ovarian cancer cytosols. <i>Clinical Chemistry</i> , 2006 , 52, 1879-86	5.5	35
110	Prognostic significance of the expression of SR-A1, encoding a novel SR-related CTD-associated factor, in breast cancer. <i>Biological Chemistry</i> , 2006 , 387, 1613-8	4.5	5
109	A comprehensive nomenclature for serine proteases with homology to tissue kallikreins. <i>Biological Chemistry</i> , 2006 , 387, 637-41	4.5	102
108	mRNA expression analysis of human kallikrein 11 (KLK11) may be useful in the discrimination of benign prostatic hyperplasia from prostate cancer after needle prostate biopsy. <i>Biological Chemistry</i> , 2006 , 387, 789-93	4.5	16
107	Effects of long-term androgen administration on breast tissue of female-to-male transsexuals. <i>Journal of Histochemistry and Cytochemistry</i> , 2006 , 54, 905-10	3.4	53
106	Serum and urine tissue kallikrein concentrations in male-to-female transsexuals treated with antiandrogens and estrogens. <i>Clinical Chemistry</i> , 2006 , 52, 1356-65	5.5	5
105	Topotecan and methotrexate alter expression of the apoptosis-related genes BCL2, FAS and BCL2L12 in leukemic HL-60 cells. <i>Biological Chemistry</i> , 2006 , 387, 1629-33	4.5	16
104	Treatment of MCF-7 cells with taxol and etoposide induces distinct alterations in the expression of apoptosis-related genes BCL2, BCL2L12, BAX, CASPASE-9 and FAS. <i>Biological Chemistry</i> , 2006 , 387, 1081-6	4.5	28
103	Expression analysis and prognostic significance of the SRA1 gene, in ovarian cancer. <i>Biochemical and Biophysical Research Communications</i> , 2006 , 344, 667-74	3.4	7
102	10-(2-Biotinyloxyethyl)-9-acridone: A novel fluorescent label for (strept)avidinBiotin based assays. <i>Journal of Photochemistry and Photobiology A: Chemistry</i> , 2006 , 181, 126-131	4.7	10
101	Alterations in mRNA expression of apoptosis-related genes BCL2, BAX, FAS, caspase-3, and the novel member BCL2L12 after treatment of human leukemic cell line HL60 with the antineoplastic agent etoposide. <i>Annals of the New York Academy of Sciences</i> , 2006 , 1090, 89-97	6.5	36
100	BCL2 family of apoptosis-related genes: functions and clinical implications in cancer. <i>Critical Reviews in Clinical Laboratory Sciences</i> , 2006 , 43, 1-67	9.4	179

99	Safety and efficacy of trastuzumab every 3 weeks combined with cytotoxic chemotherapy in patients with HER2-positive recurrent breast cancer: findings from a case series. <i>Oncology Research and Treatment</i> , 2005 , 28, 558-64	2.8	5
98	Human tissue kallikreins: from gene structure to function and clinical applications. <i>Advances in Clinical Chemistry</i> , 2005 , 39, 11-79	5.8	49
97	Polyadenylate polymerase modulations in human epithelioid cervix and breast cancer cell lines, treated with etoposide or cordycepin, follow cell cycle rather than apoptosis induction. <i>Biological Chemistry</i> , 2005 , 386, 471-80	4.5	38
96	Expression of the C-terminal domain of novel human SR-A1 protein: interaction with the CTD domain of RNA polymerase II. <i>Biochemical and Biophysical Research Communications</i> , 2005 , 334, 61-8	3.4	6
95	Novel biotinylated acridinium derivatives: new reagents for fluorescence immunoassays and proteomics. <i>Clinica Chimica Acta</i> , 2005 , 357, 159-67	6.2	9
94	Expression analysis and prognostic significance of human kallikrein 11 in prostate cancer. <i>Clinica Chimica Acta</i> , 2005 , 357, 190-5	6.2	34
93	Free/Total PSA (F/T ratio) kinetics in patients with clinically localized prostate cancer undergoing radical prostatectomy. <i>Clinica Chimica Acta</i> , 2005 , 357, 196-201	6.2	14
92	mRNA quantification and clinical evaluation of telomerase reverse transcriptase subunit (hTERT) in intracranial tumours of patients in the island of Crete. <i>British Journal of Cancer</i> , 2005 , 93, 152-8	8.7	1
91	Immunohistochemical expression of Bcl2 is an independent predictor of time-to-biochemical failure in patients with clinically localized prostate cancer following radical prostatectomy. <i>Anticancer Research</i> , 2005 , 25, 3123-33	2.3	29
90	Expression analysis of the human kallikrein 7 (KLK7) in breast tumors: a new potential biomarker for prognosis of breast carcinoma. <i>Thrombosis and Haemostasis</i> , 2004 , 91, 180-6	7	54
89	Human kallikrein 13 protein in ovarian cancer cytosols: a new favorable prognostic marker. <i>Journal of Clinical Oncology</i> , 2004 , 22, 678-85	2.2	60
88	SR-A1, a member of the human pre-mRNA splicing factor family, and its expression in colon cancer progression. <i>Biological Chemistry</i> , 2004 , 385, 785-90	4.5	8
87	Expression analysis of BCL2L12, a new member of apoptosis-related genes, in colon cancer. <i>Biological Chemistry</i> , 2004 , 385, 779-83	4.5	27
86	mRNA expression analysis of a variety of apoptosis-related genes, including the novel gene of the BCL2-family, BCL2L12, in HL-60 leukemia cells after treatment with carboplatin and doxorubicin. <i>Biological Chemistry</i> , 2004 , 385, 1099-103	4.5	28
85	Altered kallikrein 7 and 10 concentrations in cerebrospinal fluid of patients with Alzheimer® disease and frontotemporal dementia. <i>Clinical Biochemistry</i> , 2004 , 37, 230-7	3.5	36
84	Human kallikrein 11: an indicator of favorable prognosis in ovarian cancer patients. <i>Clinical Biochemistry</i> , 2004 , 37, 823-9	3.5	43
83	Hepsin is highly over expressed in and a new candidate for a prognostic indicator in prostate cancer. <i>Journal of Urology</i> , 2004 , 171, 187-91	2.5	105
82	Cathepsin B and cathepsin D expression in the progression of colorectal adenoma to carcinoma. <i>Cancer Letters</i> , 2004 , 205, 97-106	9.9	61

81	Quantitative analysis of macrophage inhibitory cytokine-1 (MIC-1) gene expression in human prostatic tissues. <i>British Journal of Cancer</i> , 2003 , 88, 1101-4	8.7	78
80	Immunofluorometric quantification of human kallikrein 5 expression in ovarian cancer cytosols and its association with unfavorable patient prognosis. <i>Tumor Biology</i> , 2003 , 24, 299-309	2.9	43
79	Expression of BCL2L12, a new member of apoptosis-related genes, in breast tumors. <i>Thrombosis and Haemostasis</i> , 2003 , 89, 1081-1088	7	35
78	Cisplatin-induced apoptosis in HL-60 human promyelocytic leukemia cells: differential expression of BCL2 and novel apoptosis-related gene BCL2L12. <i>Annals of the New York Academy of Sciences</i> , 2003 , 1010, 153-8	6.5	39
77	The prognostic value of the human kallikrein gene 9 (KLK9) in breast cancer. <i>Breast Cancer Research and Treatment</i> , 2003 , 78, 149-58	4.4	41
76	Favorable prognostic value of tissue human kallikrein 11 (hK11) in patients with ovarian carcinoma. <i>International Journal of Cancer</i> , 2003 , 106, 605-610	7.5	51
75	Prognostic value of quantitatively assessed KLK7 expression in ovarian cancer. <i>Clinical Biochemistry</i> , 2003 , 36, 135-43	3.5	60
74	Serum human glandular kallikrein (hK2) and insulin-like growth factor 1 (IGF-1) improve the discrimination between prostate cancer and benign prostatic hyperplasia in combination with total and %free PSA. <i>Prostate</i> , 2003 , 54, 220-9	4.2	37
73	Differential expression of the human kallikrein gene 14 (KLK14) in normal and cancerous prostatic tissues. <i>Prostate</i> , 2003 , 56, 287-92	4.2	55
72	Quantitative Analysis of Kallikrein 15 Gene Expression in Prostate Tissue. <i>Journal of Urology</i> , 2003 , 169, 361-364	2.5	47
71	Quantitative analysis of hippostasin/KLK11 gene expression in cancerous and noncancerous prostatic tissues. <i>Urology</i> , 2003 , 61, 1042-6	1.6	37
70	Molecular characterization of a new gene, CEAL1, encoding for a carcinoembryonic antigen-like protein with a highly conserved domain of eukaryotic translation initiation factors. <i>Gene</i> , 2003 , 310, 79-89 ⁸	3.8	22
69	Human kallikrein 6 (hK6): a new potential serum biomarker for diagnosis and prognosis of ovarian carcinoma. <i>Journal of Clinical Oncology</i> , 2003 , 21, 1035-43	2.2	176
68	Steroid hormone regulation and prognostic value of the human kallikrein gene 14 in ovarian cancer. <i>American Journal of Clinical Pathology</i> , 2003 , 119, 346-55	1.9	55
67	Prognostic value of the human kallikrein gene 15 expression in ovarian cancer. <i>Journal of Clinical Oncology</i> , 2003 , 21, 3119-26	2.2	63
66	Steroid Hormone Regulation and Prognostic Value of the Human Kallikrein Gene 14 in Ovarian Cancer. <i>American Journal of Clinical Pathology</i> , 2003 , 119, 0-0	1.9	
65	Quantitative Analysis of Kallikrein 15 Gene Expression in Prostate Tissue. <i>Journal of Urology</i> , 2003 , 361-364	3.6	1
64	The serum concentration of human kallikrein 10 represents a novel biomarker for ovarian cancer diagnosis and prognosis. <i>Cancer Research</i> , 2003 , 63, 807-11	10.1	107

63	Parallel overexpression of seven kallikrein genes in ovarian cancer. <i>Cancer Research</i> , 2003 , 63, 2223-7	10.1	110
62	Human kallikrein 8, a novel biomarker for ovarian carcinoma. <i>Cancer Research</i> , 2003 , 63, 2771-4	10.1	56
61	Expression of BCL2L12, a new member of apoptosis-related genes, in breast tumors. <i>Thrombosis and Haemostasis</i> , 2003 , 89, 1081-8	7	10
60	Human kallikrein 5: a potential novel serum biomarker for breast and ovarian cancer. <i>Cancer Research</i> , 2003 , 63, 3958-65	10.1	93
59	The usefulness of serum human kallikrein 11 for discriminating between prostate cancer and benign prostatic hyperplasia. <i>Cancer Research</i> , 2003 , 63, 6543-6	10.1	38
58	Down-regulation of the human kallikrein gene 5 (KLK5) in prostate cancer tissues. <i>Prostate</i> , 2002 , 51, 126-32	4.2	57
57	Decreased concentration of human kallikrein 6 in brain extracts of Alzheimer® disease patients. <i>Clinical Biochemistry</i> , 2002 , 35, 225-31	3.5	47
56	Quantitative analysis of human kallikrein gene 14 expression in breast tumours indicates association with poor prognosis. <i>British Journal of Cancer</i> , 2002 , 87, 1287-93	8.7	38
55	Human kallikrein gene 13 (KLK13) expression by quantitative RT-PCR: an independent indicator of favourable prognosis in breast cancer. <i>British Journal of Cancer</i> , 2002 , 86, 1457-64	8.7	53
54	The androgen-regulated gene human kallikrein 15 (KLK15) is an independent and favourable prognostic marker for breast cancer. <i>British Journal of Cancer</i> , 2002 , 87, 1294-300	8.7	40
53	Immunofluorometric quantitation and histochemical localisation of kallikrein 6 protein in ovarian cancer tissue: a new independent unfavourable prognostic biomarker. <i>British Journal of Cancer</i> , 2002 , 87, 763-71	8.7	66
52	Polyadenylate polymerase (PAP) and 3' end pre-mRNA processing: function, assays, and association with disease. <i>Critical Reviews in Clinical Laboratory Sciences</i> , 2002 , 39, 193-224	9.4	24
51	Determination of cathepsin B expression may offer additional prognostic information for ovarian cancer patients. <i>Biological Chemistry</i> , 2002 , 383, 1297-303	4.5	36
50	Significance of urokinase-type plasminogen activator and plasminogen activator inhibitor-1 (PAI-1) expression in human colorectal carcinomas. <i>Tumor Biology</i> , 2002 , 23, 170-8	2.9	20
49	Differential expression of Kallikrein gene 5 in cancerous and normal testicular tissues. <i>Urology</i> , 2002 , 60, 714-8	1.6	41
48	Cloning, physical mapping and structural characterization of the human alpha(A)-adaplin gene. <i>Gene</i> , 2002 , 289, 191-9	3.8	6
47	Human Kallikrein Gene 5 (KLK5) Expression by Quantitative PCR: An Independent Indicator of Poor Prognosis in Breast Cancer. <i>Clinical Chemistry</i> , 2002 , 48, 1241-1250	5.5	72
46	Detection of Human Kallikrein 4 in Healthy and Cancerous Prostatic Tissues by Immunofluorometry and Immunohistochemistry. <i>Clinical Chemistry</i> , 2002 , 48, 1232-1240	5.5	60

45	Human kallikrein gene 5 (KLK5) expression by quantitative PCR: an independent indicator of poor prognosis in breast cancer. <i>Clinical Chemistry</i> , 2002 , 48, 1241-50	5.5	19
44	Expression of gelatinase-A (MMP-2) in human colon cancer and normal colon mucosa. <i>Tumor Biology</i> , 2001 , 22, 383-9	2.9	39
43	Combined expression of p53, Bcl-2, and p21WAF-1 proteins in lung cancer and premalignant lesions: association with clinical characteristics. <i>Lung</i> , 2001 , 179, 265-78	2.9	16
42	Human kallikrein gene 5 (KLK5) expression is an indicator of poor prognosis in ovarian cancer. <i>British Journal of Cancer</i> , 2001 , 84, 643-50	8.7	103
41	Codon 89 polymorphism in the human 5 alpha-reductase gene in primary breast cancer. <i>British Journal of Cancer</i> , 2001 , 84, 760-7	8.7	14
40	Overexpression of matrix-metalloproteinase-9 in human breast cancer: a potential favourable indicator in node-negative patients. <i>British Journal of Cancer</i> , 2001 , 84, 1488-96	8.7	182
39	Cloning of a gene (SR-A1), encoding for a new member of the human Ser/Arg-rich family of pre-mRNA splicing factors: overexpression in aggressive ovarian cancer. <i>British Journal of Cancer</i> , 2001 , 85, 190-8	8.7	18
38	Molecular cloning of the human kallikrein 15 gene (KLK15). Up-regulation in prostate cancer. <i>Journal of Biological Chemistry</i> , 2001 , 276, 53-61	5.4	89
37	Insulin-like growth factor I (IGF-I) and IGF-binding protein-3 in benign prostatic hyperplasia and prostate cancer. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2001 , 86, 694-9	5.6	54
36	Prostate-specific antigen and human glandular kallikrein 2 are markedly elevated in urine of patients with polycystic ovary syndrome. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2001 , 86, 1558-61	5.6	26
35	Identification and characterization of a novel human testis-specific kinase substrate gene which is downregulated in testicular tumors. <i>Biochemical and Biophysical Research Communications</i> , 2001 , 285, 400-8	3.4	13
34	Molecular cloning, physical mapping, and expression analysis of a novel gene, BCL2L12, encoding a proline-rich protein with a highly conserved BH2 domain of the Bcl-2 family. <i>Genomics</i> , 2001 , 72, 217-21	4.3	81
33	Human kallikrein 10: a novel tumor marker for ovarian carcinoma?. <i>Clinica Chimica Acta</i> , 2001 , 306, 111-86.2		89
32	The human KLK8 (neuropsin/ovasin) gene: identification of two novel splice variants and its prognostic value in ovarian cancer. <i>Clinical Cancer Research</i> , 2001 , 7, 806-11	12.9	86
31	Prognostic value of human kallikrein 10 expression in epithelial ovarian carcinoma. <i>Clinical Cancer Research</i> , 2001 , 7, 2372-9	12.9	69
30	Higher human kallikrein gene 4 (KLK4) expression indicates poor prognosis of ovarian cancer patients. <i>Clinical Cancer Research</i> , 2001 , 7, 2380-6	12.9	80
29	Quantitative expression of the human kallikrein gene 9 (KLK9) in ovarian cancer: a new independent and favorable prognostic marker. <i>Cancer Research</i> , 2001 , 61, 7811-8	10.1	67
28	Polyvinylamine-streptavidin complexes labeled with a europium chelator: a universal detection reagent for solid-phase time resolved fluorometric applications. <i>Clinical Biochemistry</i> , 2000 , 33, 345-50	3.5	31

27	TNF-alpha expression and apoptosis-regulating proteins in oral lichen planus: a comparative immunohistochemical evaluation. <i>Journal of Oral Pathology and Medicine</i> , 2000 , 29, 370-5	3.3	49
26	Breast cancer prognostic significance of a single nucleotide polymorphism in the proximal androgen response element of the prostate specific antigen gene promoter. <i>Breast Cancer Research and Treatment</i> , 2000 , 61, 111-9	4.4	15
25	Serum and Urinary Prostate-specific Antigen and Urinary Human Glandular Kallikrein Concentrations Are Significantly Increased after Testosterone Administration in Female-to-Male Transsexuals. <i>Clinical Chemistry</i> , 2000 , 46, 859-862	5.5	25
24	Serum human glandular kallikrein-2 protease levels predict the presence of prostate cancer among men with elevated prostate-specific antigen. <i>Journal of Clinical Oncology</i> , 2000 , 18, 1036-42	2.2	92
23	Streptavidin-Polyvinylamine Conjugates Labeled with a Europium Chelate: Applications in Immunoassay, Immunohistochemistry, and Microarrays. <i>Clinical Chemistry</i> , 2000 , 46, 1450-1455	5.5	62
22	Genomic organization of the human kallikrein gene family on chromosome 19q13.3-q13.4. <i>Biochemical and Biophysical Research Communications</i> , 2000 , 276, 125-33	3.4	169
21	Genomic organization, physical mapping, and expression analysis of the human protein arginine methyltransferase 1 gene. <i>Biochemical and Biophysical Research Communications</i> , 2000 , 278, 349-59	3.4	57
20	Genomic organization, mapping, tissue expression, and hormonal regulation of trypsin-like serine protease (TLSP PRSS20), a new member of the human kallikrein gene family. <i>Genomics</i> , 2000 , 63, 88-96	4.3	60
19	The KLK7 (PRSS6) gene, encoding for the stratum corneum chymotryptic enzyme is a new member of the human kallikrein gene family - genomic characterization, mapping, tissue expression and hormonal regulation. <i>Gene</i> , 2000 , 254, 119-28	3.8	72
18	Comparison of the percent free prostate-specific antigen levels in the serum of healthy men and in men with recurrent prostate cancer after radical prostatectomy. <i>Clinica Chimica Acta</i> , 2000 , 292, 127-38	6.2	12
17	Decreased concentrations of prostate-specific antigen and human glandular kallikrein 2 in malignant versus nonmalignant prostatic tissue. <i>Urology</i> , 2000 , 56, 527-32	1.6	89
16	DRAMATIC SUPPRESSION OF PLASMA AND URINARY PROSTATE SPECIFIC ANTIGEN AND HUMAN GLANDULAR KALLIKREIN BY ANTIANDROGENS IN MALE-TO-FEMALE TRANSSEXUALS. <i>Journal of Urology</i> , 2000 , 163, 802-805	2.5	12
15	TA repeat polymorphism of the 5alpha-reductase gene and breast cancer. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2000 , 9, 387-93	4	6
14	Polyadenylate polymerase enzymatic activity in mammary tumor cytosols: A new independent prognostic marker in primary breast cancer. <i>Cancer Research</i> , 2000 , 60, 5427-33	10.1	22
13	Human Glandular Kallikrein in Breast Milk, Amniotic Fluid, and Breast Cyst Fluid. <i>Clinical Chemistry</i> , 1999 , 45, 1774-1780	5.5	31
12	The Combination of Human Glandular Kallikrein and Free Prostate-specific Antigen (PSA) Enhances Discrimination Between Prostate Cancer and Benign Prostatic Hyperplasia in Patients with Moderately Increased Total PSA. <i>Clinical Chemistry</i> , 1999 , 45, 1960-1966	5.5	95
11	Determination of c-myc amplification and overexpression in breast cancer patients: evaluation of its prognostic value against c-erbB-2, cathepsin-D and clinicopathological characteristics using univariate and multivariate analysis. <i>British Journal of Cancer</i> , 1999 , 81, 1385-91	8.7	36
10	Predictive value of c-erbB-2 and cathepsin-D for Greek breast cancer patients using univariate and multivariate analysis. <i>Clinical Cancer Research</i> , 1999 , 5, 815-21	12.9	20

9	Immunoenzymatically determined pepsinogen C concentration in breast tumor cytosols: an independent favorable prognostic factor in node-positive patients. <i>Clinical Cancer Research</i> , 1999 , 5, 1778-85	12.9	27
8	The combination of human glandular kallikrein and free prostate-specific antigen (PSA) enhances discrimination between prostate cancer and benign prostatic hyperplasia in patients with moderately increased total PSA. <i>Clinical Chemistry</i> , 1999 , 45, 1960-6	5.5	24
7	Cathepsin D may help in discriminating node-negative breast cancer patients at risk for local-regional recurrence. <i>Anticancer Research</i> , 1998 , 18, 2885-90	2.3	5
6	Poly(A)polymerase activity levels in breast tumour cytosols. <i>Journal of Experimental and Clinical Cancer Research</i> , 1998 , 17, 511-8		3
5	Cathepsin D concentration in tumor cytosols improves the accuracy of prognostic evaluation of primary breast cancer. <i>Anticancer Research</i> , 1997 , 17, 1405-9	2.3	5
4	Relationships between cathepsin-D, pS2 protein and hormonal receptors in breast cancer cytosols: inconsistency with their established prognostic significance. <i>Anticancer Research</i> , 1997 , 17, 3665-9	2.3	10
3	c-erbB-2 overexpression may be used as an independent prognostic factor for breast cancer patients. <i>Anticancer Research</i> , 1995 , 15, 1543-7	2.3	8
2	Cathepsin-D and c-erb-B 2 have an additive prognostic value for breast cancer patients. <i>Anticancer Research</i> , 1993 , 13, 1895-900	2.3	14
1	Prostate-Specific Antigen and Human Glandular Kallikrein 2 Are Markedly Elevated in Urine of Patients with Polycystic Ovary Syndrome		6