Takashi Takeda

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/8343514/takashi-takeda-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

44 454 13 20 g-index

47 595 2.8 3.62 ext. papers ext. citations avg, IF L-index

#	Paper	IF	Citations
44	Prevalence of Premenstrual Syndrome and Premenstrual Dysphoric Disorder among Mongolian College Students. <i>Kitakanto Medical Journal</i> , 2022 , 72, 43-48	О	
43	Characteristics of the gut microbiota in women with premenstrual symptoms: A cross-sectional study. <i>PLoS ONE</i> , 2022 , 17, e0268466	3.7	0
42	Association Between Serious Psychological Distress and Loneliness During the COVID-19 Pandemic: A Cross-Sectional Study with Pregnant Japanese Women. <i>International Journal of Womenis Health</i> , 2021 , 13, 1087-1093	2.8	O
41	Internet addiction belief, but not Internet use time, is independently associated with menstrual pain severity and interference to social life among adolescents: a cross-sectional study. <i>British Journal of Pain</i> , 2021 , 15, 302-311	2.1	1
40	Psychometric Testing of the Japanese Version of the Daily Record of Severity of Problems Among Japanese Women. <i>International Journal of Womenis Health</i> , 2021 , 13, 361-367	2.8	1
39	The delivery of a placenta/fetus with high gonadal steroid production contributes to postpartum depressive symptoms. <i>Depression and Anxiety</i> , 2021 , 38, 422-430	8.4	3
38	Association between Premenstrual Symptoms and Posttraumatic Stress Symptoms by COVID-19: A Cross-Sectional Study with Japanese High School Students. <i>Tohoku Journal of Experimental Medicine</i> , 2021 , 255, 71-77	2.4	1
37	A Multicenter, Randomized, Double-Blind, Placebo-Controlled Trial to Investigate the Effects of Kamishoyosan, a Traditional Japanese Medicine, on Menopausal Symptoms: The KOSMOS Study. <i>Evidence-based Complementary and Alternative Medicine</i> , 2021 , 2021, 8856149	2.3	2
36	When and how do adolescent girls in Japan become aware of premenstrual symptoms from menarche? A cross-sectional study among senior high school students. <i>BMJ Open</i> , 2021 , 11, e045215	3	O
35	Psychometric Testing of the Premenstrual Symptoms Questionnaire and the Association Between Perceived Injustice and Premenstrual Symptoms: A Cross-Sectional Study Among Japanese High School Students. <i>International Journal of Womens Health</i> , 2020 , 12, 755-763	2.8	5
34	Associations between sleep habits and interference of premenstrual symptoms in athletic performance in Japanese adolescent athletes: a cohort study over a 2-year period. <i>Gynecological Endocrinology</i> , 2020 , 36, 885-889	2.4	2
33	Pain medications during pregnancy: data from the Japan environment and children study. <i>Journal of Anesthesia</i> , 2020 , 34, 202-210	2.2	О
32	Effects of Kamishoyosan, a Traditional Japanese Medicine, on Menopausal Symptoms: A Randomized, Placebo-Controlled, Double-Blind Clinical Trial. <i>Evidence-based Complementary and Alternative Medicine</i> , 2020 , 2020, 9285317	2.3	3
31	Lifestyle Factors Associated with Premenstrual Syndrome: AlCross-sectional Study of Japanese High School Students. <i>Journal of Pediatric and Adolescent Gynecology</i> , 2019 , 32, 590-595	2	10
30	Developing a Japanese version of the Injustice Experience Questionnaire-chronic and the contribution of perceived injustice to severity of menstrual pain: a web-based cross-sectional study. <i>BioPsychoSocial Medicine</i> , 2019 , 13, 17	2.8	5
29	Antidepressive Effects of Kamishoyosan through 5-HT1AReceptor and PKA-CREB-BDNF Signaling in the Hippocampus in Postmenopausal Depression-Model Mice. <i>Evidence-based Complementary and Alternative Medicine</i> , 2019 , 2019, 9475384	2.3	10
28	Preconception dysmenorrhea as a risk factor for psychological distress in pregnancy: The Japan Environment and Children's Study. <i>Journal of Affective Disorders</i> , 2019 , 245, 475-483	6.6	4

27	Premenstrual symptoms interference and equol production status in Japanese collegiate athletes: A cross-sectional study. <i>Journal of Obstetrics and Gynaecology Research</i> , 2018 , 44, 488-494	1.9	3
26	Royal Jelly Supplementation Improves Menopausal Symptoms Such as Backache, Low Back Pain, and Anxiety in Postmenopausal Japanese Women. <i>Evidence-based Complementary and Alternative Medicine</i> , 2018 , 2018, 4868412	2.3	8
25	Effectiveness of natural S-equol supplement for premenstrual symptoms: protocol of a randomised, double-blind, placebo-controlled trial. <i>BMJ Open</i> , 2018 , 8, e023314	3	7
24	Low Proportion of Dietary Plant Protein among Athletes with Premenstrual Syndrome-Related Performance Impairment. <i>Tohoku Journal of Experimental Medicine</i> , 2018 , 244, 119-122	2.4	1
23	Effect of an educational program on adolescent premenstrual syndrome: lessons from the Great East Japan Earthquake. <i>Adolescent Health, Medicine and Therapeutics</i> , 2018 , 9, 95-101	2.7	2
22	Relation between premenstrual syndrome and equol-production status. <i>Journal of Obstetrics and Gynaecology Research</i> , 2016 , 42, 1575-1580	1.9	6
21	Stress fracture and premenstrual syndrome in Japanese adolescent athletes: a cross-sectional study. <i>BMJ Open</i> , 2016 , 6, e013103	3	12
20	Psychological distress during pregnancy in Miyagi after the Great East Japan Earthquake: The Japan Environment and Children Study. <i>Journal of Affective Disorders</i> , 2016 , 190, 341-348	6.6	39
19	Dysmenorrhea and PTSD 2016 , 1315-1327		
18	Fish Consumption and Premenstrual Syndrome and Dysphoric Disorder in Japanese Collegiate Athletes. <i>Journal of Pediatric and Adolescent Gynecology</i> , 2016 , 29, 386-389	2	9
18		2.8	9
	Athletes. Journal of Pediatric and Adolescent Gynecology, 2016, 29, 386-389 The prevalence and risk factors of school absenteeism due to premenstrual disorders in Japanese		
17	Athletes. <i>Journal of Pediatric and Adolescent Gynecology</i> , 2016 , 29, 386-389 The prevalence and risk factors of school absenteeism due to premenstrual disorders in Japanese high school students-a school-based cross-sectional study. <i>BioPsychoSocial Medicine</i> , 2016 , 10, 13 Additional data to Relation between premenstrual syndrome and equol-production status <i>Journal</i>	2.8	
17 16	Athletes. Journal of Pediatric and Adolescent Gynecology, 2016, 29, 386-389 The prevalence and risk factors of school absenteeism due to premenstrual disorders in Japanese high school students-a school-based cross-sectional study. BioPsychoSocial Medicine, 2016, 10, 13 Additional data to Relation between premenstrual syndrome and equol-production status V Journal of Obstetrics and Gynaecology Research, 2016, 42, 1631 The anti-diabetic drug metformin inhibits vascular endothelial growth factor expression via the mammalian target of rapamycin complex 1/hypoxia-inducible factor-1ßignaling pathway in ELT-3	2.8	13
17 16 15	The prevalence and risk factors of school absenteeism due to premenstrual disorders in Japanese high school students-a school-based cross-sectional study. <i>BioPsychoSocial Medicine</i> , 2016 , 10, 13 Additional data to Relation between premenstrual syndrome and equol-production status <i>Journal of Obstetrics and Gynaecology Research</i> , 2016 , 42, 1631 The anti-diabetic drug metformin inhibits vascular endothelial growth factor expression via the mammalian target of rapamycin complex 1/hypoxia-inducible factor-1 ignaling pathway in ELT-3 cells. <i>Molecular and Cellular Endocrinology</i> , 2015 , 399, 1-8 Effectiveness of ethinylestradiol/drospirenone for premenstrual symptoms in Japanese patients with dysmenorrhea: Open-label pilot study. <i>Journal of Obstetrics and Gynaecology Research</i> , 2015 ,	2.8 1.9	13
17 16 15	Athletes. Journal of Pediatric and Adolescent Gynecology, 2016, 29, 386-389 The prevalence and risk factors of school absenteeism due to premenstrual disorders in Japanese high school students-a school-based cross-sectional study. BioPsychoSocial Medicine, 2016, 10, 13 Additional data to Relation between premenstrual syndrome and equol-production statusV Journal of Obstetrics and Gynaecology Research, 2016, 42, 1631 The anti-diabetic drug metformin inhibits vascular endothelial growth factor expression via the mammalian target of rapamycin complex 1/hypoxia-inducible factor-1ßignaling pathway in ELT-3 cells. Molecular and Cellular Endocrinology, 2015, 399, 1-8 Effectiveness of ethinylestradiol/drospirenone for premenstrual symptoms in Japanese patients with dysmenorrhea: Open-label pilot study. Journal of Obstetrics and Gynaecology Research, 2015, 41, 1584-90 The Kampo Medicine Yokukansan Decreases MicroRNA-18 Expression and Recovers Glucocorticoid Receptors Protein Expression in the Hypothalamus of Stressed Mice. BioMed Research International,	2.8 1.9 4.4	13 31 3
17 16 15 14	Athletes. Journal of Pediatric and Adolescent Gynecology, 2016, 29, 386-389 The prevalence and risk factors of school absenteeism due to premenstrual disorders in Japanese high school students-a school-based cross-sectional study. BioPsychoSocial Medicine, 2016, 10, 13 Additional data to Relation between premenstrual syndrome and equol-production status Journal of Obstetrics and Gynaecology Research, 2016, 42, 1631 The anti-diabetic drug metformin inhibits vascular endothelial growth factor expression via the mammalian target of rapamycin complex 1/hypoxia-inducible factor-1ßignaling pathway in ELT-3 cells. Molecular and Cellular Endocrinology, 2015, 399, 1-8 Effectiveness of ethinylestradiol/drospirenone for premenstrual symptoms in Japanese patients with dysmenorrhea: Open-label pilot study. Journal of Obstetrics and Gynaecology Research, 2015, 41, 1584-90 The Kampo Medicine Yokukansan Decreases MicroRNA-18 Expression and Recovers Glucocorticoid Receptors Protein Expression in the Hypothalamus of Stressed Mice. BioMed Research International, 2015, 2015, 797280 Premenstrual Syndrome and Premenstrual Dysphoric Disorder in Japanese Collegiate Athletes.	2.8 1.9 4.4 1.9	13 31 3 20

9	Epigallocatechin-3-gallate potentiates curcumin\ ability to suppress uterine leiomyosarcoma cell growth and induce apoptosis. <i>International Journal of Clinical Oncology</i> , 2013 , 18, 380-8	4.2	17
8	Relationship between dysmenorrhea and posttraumatic stress disorder in Japanese high school students 9 months after the Great East Japan Earthquake. <i>Journal of Pediatric and Adolescent Gynecology</i> , 2013 , 26, 355-7	2	20
7	The antidiabetic drug metformin inhibits uterine leiomyoma cell proliferation via an AMP-activated protein kinase signaling pathway. <i>Gynecological Endocrinology</i> , 2013 , 29, 87-90	2.4	12
6	Premenstrual symptoms and posttraumatic stress disorder in Japanese high school students 9 months after the great East-Japan earthquake. <i>Tohoku Journal of Experimental Medicine</i> , 2013 , 230, 151	1 -2 4	25
5	Relationship between premenstrual symptoms and dysmenorrhea in Japanese high school students. <i>Archives of Womenis Mental Health</i> , 2012 , 15, 131-3	5	31
4	Curcumin disrupts uterine leiomyosarcoma cells through AKT-mTOR pathway inhibition. <i>Gynecologic Oncology</i> , 2011 , 122, 141-8	4.9	33
3	Changes of blood flow volume in the superior mesenteric artery and brachial artery with abdominal thermal stimulation. <i>Evidence-based Complementary and Alternative Medicine</i> , 2011 , 2011, 214089	2.3	15
2	Establishment of a novel xenograft model for human uterine leiomyoma in immunodeficient mice. <i>Tohoku Journal of Experimental Medicine</i> , 2010 , 222, 55-61	2.4	17
1	Prevalence of premenstrual syndrome and premenstrual dysphoric disorder in Japanese high school students. <i>Archives of Womens Mental Health</i> , 2010 , 13, 535-7	5	41