
Yoshiteru Takekita

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8346965/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Effects of the Serotonin Type 2A, 3A and 3B Receptor and the Serotonin Transporter Genes on Paroxetine and Fluvoxamine Efficacy and Adverse Drug Reactions in Depressed Japanese Patients. Neuropsychobiology, 2006, 53, 186-195.	0.9	143
2	ABCB1 (MDR1) gene polymorphisms are associated with the clinical response to paroxetine in patients with major depressive disorder. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2008, 32, 398-404.	2.5	126
3	Effect of 5â€HT1A gene polymorphisms on antidepressant response in major depressive disorder. American Journal of Medical Genetics Part B: Neuropsychiatric Genetics, 2009, 150B, 115-123.	1.1	89
4	Differences in quantitative EEG between frontotemporal dementia and Alzheimer's disease as revealed by LORETA. Clinical Neurophysiology, 2011, 122, 1718-1725.	0.7	69
5	The Alpha 2A-Adrenergic Receptor Gene Polymorphism Modifies Antidepressant Responses to Milnacipran. Journal of Clinical Psychopharmacology, 2008, 28, 518-524.	0.7	30
6	Antidepressant response and intolerance to SSRI is not influenced by G-protein β3 subunit gene C825T polymorphism in Japanese major depressive patients. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2008, 32, 1041-1044.	2.5	29
7	Switching to antipsychotic monotherapy vs. staying on antipsychotic polypharmacy in schizophrenia: A systematic review and meta-analysis. Schizophrenia Research, 2019, 209, 50-57.	1.1	27
8	HTR1A Polymorphisms and Clinical Efficacy of Antipsychotic Drug Treatment in Schizophrenia: A Meta-Analysis. International Journal of Neuropsychopharmacology, 2016, 19, pyv125.	1.0	26
9	Effect of basic fibroblast growth factor (FGF2) gene polymorphisms on SSRIs treatment response and side effects. European Neuropsychopharmacology, 2009, 19, 718-725.	0.3	24
10	Remifentanil in electroconvulsive therapy: a systematic review and meta-analysis of randomized controlled trials. European Archives of Psychiatry and Clinical Neuroscience, 2016, 266, 703-717.	1.8	24
11	HTR1A Gene Polymorphisms and 5-HT1A Receptor Partial Agonist Antipsychotics Efficacy in Schizophrenia. Journal of Clinical Psychopharmacology, 2015, 35, 220-227.	0.7	22
12	The Comparative Effects of Risperidone Long-Acting Injection and Paliperidone Palmitate on Social Functioning in Schizophrenia: A 6-Month, Open-Label, Randomized Controlled Pilot Trial. Neuropsychobiology, 2016, 73, 35-42.	0.9	21
13	Genetic variants in combination with early partial improvement as a clinical utility predictor of treatment outcome in major depressive disorder: the result of two pooled RCTs. Translational Psychiatry, 2015, 5, e513-e513.	2.4	20
14	Therapeutic Response to Paroxetine in Major Depressive Disorder Predicted by DNA Methylation. Neuropsychobiology, 2017, 75, 81-88.	0.9	19
15	Non response at week 4 as clinically useful indicator for antidepressant combination in major depressive disorder. A sequential RCT. Journal of Psychiatric Research, 2017, 89, 97-104.	1.5	17
16	Cognitive function and risperidone long-acting injection vs. paliperidone palmitate in schizophrenia: a 6-month, open-label, randomized, pilot trial. BMC Psychiatry, 2016, 16, 172.	1.1	16
17	A 12-week randomized, open-label study of perospirone versus aripiprazole in the treatment of Japanese schizophrenia patients. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2013, 40, 110-114.	2.5	13
18	5-HTTLPR rs25531A > G Differentially Influence Paroxetine and Fluvoxamine Antidepressant Efficacy. Journal of Clinical Psychopharmacology, 2013, 33, 131-132.	0.7	12

#	Article	IF	CITATIONS
19	Polymorphism of rs3813034 in Serotonin Transporter Gene SLC6A4 Is Associated With the Selective Serotonin and Serotonin-Norepinephrine Reuptake Inhibitor Response in Depressive Disorder. Journal of Clinical Psychopharmacology, 2016, 36, 27-31.	0.7	11
20	Syndrome of inappropriate secretion of antiâ€diuretic hormone in an elderly depressive patient receiving paroxetine: a case report. International Journal of Geriatric Psychiatry, 2010, 25, 433-434.	1.3	9
21	Delirium Associated with Paroxetine in an Elderly Depressive Patient: A Case Report. Pharmacopsychiatry, 2007, 40, 199-200.	1.7	8
22	Serotonin 7 Receptor Variants Are Not Associated with Response to Second-Generation Antipsychotics in Japanese Schizophrenia Patients. Neuropsychobiology, 2015, 72, 118-125.	0.9	8
23	Case of adultâ€onset type <scp>II</scp> citrullinemia treated as schizophrenia for a long time. Psychiatry and Clinical Neurosciences, 2015, 69, 306-307.	1.0	8
24	Multiple Pre-Treatment miRNAs Levels in Untreated Major Depressive Disorder Patients Predict Early Response to Antidepressants and Interact with Key Pathways. International Journal of Molecular Sciences, 2022, 23, 3873.	1.8	8
25	Neuropsychological Evaluation and Cerebral Blood Flow Effects of Apolipoprotein E4 in Alzheimer's Disease Patients after One Year of Treatment: An Exploratory Study. Dementia and Geriatric Cognitive Disorders Extra, 2015, 5, 414-423.	0.6	7
26	Pre-treatment plasma cytokine levels as potential predictors of short-term remission of depression. World Journal of Biological Psychiatry, 2022, 23, 785-793.	1.3	6
27	Antagonist and partial agonist at the dopamine D2 receptors in drug-naÃ⁻ve and non-drug-naÃ⁻ve schizophrenia: a randomized, controlled trial. European Archives of Psychiatry and Clinical Neuroscience, 2015, 265, 579-588.	1.8	5
28	Pharmacological treatment algorithms for the acute phase, agitation, and maintenance phase of firstâ€episode schizophrenia: Japanese Society of Clinical Neuropsychopharmacology treatment algorithms. Human Psychopharmacology, 2021, 36, e2804.	0.7	5
29	Brain Volume-Related Polymorphisms of the Glycogen Synthase Kinase-3β Gene and Their Effect on Antidepressant Treatment in Major Depressive Disorder. Neuropsychobiology, 2019, 78, 136-144.	0.9	4
30	Personality as a basis for antidepressant selection for patients with depression: A two-point outcome study at 4 and 8Âweeks. Journal of Affective Disorders, 2022, 314, 27-33.	2.0	4
31	A series of case reports on abnormal sensation on eye movement associated with paroxetine discontinuation. International Clinical Psychopharmacology, 2006, 21, A29-A30.	0.9	3
32	Olfactory reference syndrome treated by blonanserin augmentation. Psychiatry and Clinical Neurosciences, 2011, 65, 203-204.	1.0	3
33	Sevoflurane in electroconvulsive therapy: A systematic review and meta-analysis of randomised trials. Journal of Psychiatric Research, 2021, 141, 16-25.	1.5	3
34	Predictive factors for hyperglycaemic progression in patients with schizophrenia or bipolar disorder. BJPsych Open, 2018, 4, 454-460.	0.3	2
35	Divergence of dose–response with asenapine: a cluster analysis of randomized, double-blind, and placebo control study. CNS Spectrums, 2022, 27, 369-377.	0.7	2
36	5-HTTLPR rs25531A > G Differentially Influence Paroxetine and Fluvoxamine Antidepressant Efficacy. Journal of Clinical Psychopharmacology, 2012, , 1.	0.7	1

#	Article	IF	CITATIONS
37	ls long-acting injectable aripiprazole useful for the treatment of acute exacerbation of schizophrenia?. Evidence-Based Mental Health, 2016, 19, e25-e25.	2.2	0
38	<p>Rehospitalization Risk of Receptor-Affinity Profile in Antipsychotic Drug Treatment: A Propensity Score Matching Analysis Using a Japanese Employment-Based Health Insurance Database. Neuropsychiatric Disease and Treatment, 2020, Volume 16, 2871-2879.</p>	1.0	0
39	Sevoflurane anesthesia in electroconvulsive therapy: a meta-analysis of randomized controlled trials. Proceedings for Annual Meeting of the Japanese Pharmacological Society, 2018, WCP2018, PO3-1-90.	0.0	0