

# Satoshi Nakata

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8359932/publications.pdf>

Version: 2024-02-01

17  
papers

215  
citations

1307594

7  
h-index

996975

15  
g-index

17  
all docs

17  
docs citations

17  
times ranked

463  
citing authors

#	ARTICLE	IF	CITATIONS
1	GLI3s Associated With Neuronal Differentiation in SHH-Activated and WNT-Activated Medulloblastoma. <i>Journal of Neuropathology and Experimental Neurology</i> , 2021, 80, 129-136.	1.7	5
2	Ependymoma-like tumor with mesenchymal differentiation harboring <i>C11orf95</i> or <i>NCOA1</i> or <i>RELA</i> fusion: A hitherto unclassified tumor related to ependymoma. <i>Brain Pathology</i> , 2021, 31, e12943.	4.1	16
3	BCOR Internal Tandem Duplication Expression in Neural Stem Cells Promotes Growth, Invasion, and Expression of PRC2 Targets. <i>International Journal of Molecular Sciences</i> , 2021, 22, 3913.	4.1	0
4	Secondary INI1-deficient rhabdoid tumors of the central nervous system: analysis of four cases and literature review. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2020, 476, 763-772.	2.8	8
5	CNS Low-grade Diffusely Infiltrative Tumors With INI1 Deficiency, Possessing a High Propensity to Progress to Secondary INI1-deficient Rhabdoid Tumors. <i>American Journal of Surgical Pathology</i> , 2020, 44, 1459-1468.	3.7	8
6	Increased Tau Expression Correlates With IDH Mutation in Infiltrating Gliomas and Impairs Cell Migration. <i>Journal of Neuropathology and Experimental Neurology</i> , 2020, 79, 493-499.	1.7	2
7	Oligodendroglioma showing pleomorphic xanthoastrocytoma-like perivascular microlesion: With IDH1, TERT promoter mutation and 1p/19q codeletion detected in both components. <i>Pathology International</i> , 2020, 70, 40-46.	1.3	3
8	Clinicopathological characteristics of circumscribed high-grade astrocytomas with an unusual combination of BRAF V600E, ATRX, and CDKN2A/B alternations. <i>Brain Tumor Pathology</i> , 2019, 36, 103-111.	1.7	5
9	Inhibition of enhancer of zest homologue 2 is a potential therapeutic target for high-grade MYC medulloblastoma. <i>Neuropathology</i> , 2019, 39, 71-77.	1.2	8
10	Cerebellar high-grade astrocytoma with IDH mutations in the elderly: A report of two cases. <i>Neuropathology</i> , 2018, 38, 411-416.	1.2	5
11	A case report of adult cerebellar high-grade glioma with H3.1 K27M mutation: a rare example of an H3 K27M mutant cerebellar tumor. <i>Brain Tumor Pathology</i> , 2018, 35, 29-35.	1.7	5
12	CNS high-grade neuroepithelial tumor with BCOR internal tandem duplication: a comparison with its counterparts in the kidney and soft tissue. <i>Brain Pathology</i> , 2018, 28, 710-720.	4.1	67
13	PATH-46. NEURONAL DIFFERENTIATION IS INDUCED BY Gli3 IN WNT- AND SHH- ACTIVATED MEDULLOBLASTOMA. <i>Neuro-Oncology</i> , 2018, 20, vi168-vi169.	1.2	0
14	Adult-onset atypical teratoid/rhabdoid tumor featuring long spindle cells with nuclear palisading and perivascular pseudorosettes. <i>Neuropathology</i> , 2017, 37, 52-57.	1.2	15
15	A case of high-grade astrocytoma with BRAF and ATRX mutations following a long-standing course over two decades. <i>Neuropathology</i> , 2017, 37, 351-357.	1.2	5
16	Sellar Atypical Teratoid/Rhabdoid Tumor (AT/RT). <i>American Journal of Surgical Pathology</i> , 2017, 41, 932-940.	3.7	38
17	Histone H3 K27M mutations in adult cerebellar high-grade gliomas. <i>Brain Tumor Pathology</i> , 2017, 34, 113-119.	1.7	25