

Yun-fei Yuan

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8362327/yun-fei-yuan-publications-by-year.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

150
papers

6,948
citations

44
h-index

80
g-index

159
ext. papers

8,144
ext. citations

7.4
avg, IF

5.45
L-index

| # | Paper | IF | Citations |
|-----|--|------|-----------|
| 150 | Influence of AFP on surgical outcomes in non-B non-C patients with curative resection for hepatocellular carcinoma.. <i>Clinical and Experimental Medicine</i> , 2022 , 1 | 4.9 | 0 |
| 149 | The nociceptin receptor promotes autophagy through NF-kB signaling and is transcriptionally regulated by E2F1 in HCC.. <i>Cell Death Discovery</i> , 2022 , 8, 165 | 6.9 | 0 |
| 148 | More Liver Metastases Detected Intraoperatively Indicates Worse Prognosis for Colorectal Liver Metastases Patients after Resection Combined with Microwave Ablation.. <i>Journal of Oncology</i> , 2022 , 2022, 3819564 | 4.5 | 0 |
| 147 | Comprehensive Analysis to Identify the Encoded Gens of Sodium Channels as a Prognostic Biomarker in Hepatocellular Carcinoma.. <i>Frontiers in Genetics</i> , 2021 , 12, 802067 | 4.5 | 0 |
| 146 | RALYL increases hepatocellular carcinoma stemness by sustaining the mRNA stability of TGF- β . <i>Nature Communications</i> , 2021 , 12, 1518 | 17.4 | 12 |
| 145 | Elafin promotes tumour metastasis and attenuates the anti-metastatic effects of erlotinib via binding to EGFR in hepatocellular carcinoma. <i>Journal of Experimental and Clinical Cancer Research</i> , 2021 , 40, 113 | 12.8 | 8 |
| 144 | Conversion to Resectability Using Transarterial Chemoembolization Combined With Hepatic Arterial Infusion Chemotherapy for Initially Unresectable Hepatocellular Carcinoma. <i>Annals of Surgery Open</i> , 2021 , 2, e057 | 1 | 1 |
| 143 | SNRPC promotes hepatocellular carcinoma cell motility by inducing epithelial-mesenchymal transition. <i>FEBS Open Bio</i> , 2021 , 11, 1757-1770 | 2.7 | 2 |
| 142 | Surgical Resection versus Re-Ablation for Intrahepatic Recurrent Hepatocellular Carcinoma after Initial Ablation Therapy. <i>Digestive Surgery</i> , 2021 , 38, 46-57 | 2.5 | 1 |
| 141 | PARP inhibitor Olaparib overcomes Sorafenib resistance through reshaping the pluripotent transcriptome in hepatocellular carcinoma. <i>Molecular Cancer</i> , 2021 , 20, 20 | 42.1 | 12 |
| 140 | PGC7 promotes tumor oncogenic dedifferentiation through remodeling DNA methylation pattern for key developmental transcription factors. <i>Cell Death and Differentiation</i> , 2021 , 28, 1955-1970 | 12.7 | 5 |
| 139 | Long-term outcome for colorectal liver metastases: combining hepatectomy with intraoperative ultrasound guided open microwave ablation versus hepatectomy alone. <i>International Journal of Hyperthermia</i> , 2021 , 38, 372-381 | 3.7 | 0 |
| 138 | Transarterial chemoembolization (TACE) combined with apatinib versus TACE combined with sorafenib in advanced hepatocellular carcinoma patients: a multicenter retrospective study. <i>Annals of Translational Medicine</i> , 2021 , 9, 283 | 3.2 | 4 |
| 137 | Primary tumor immune score fails to predict the prognosis of colorectal cancer liver metastases after hepatectomy in Chinese populations. <i>Annals of Translational Medicine</i> , 2021 , 9, 310 | 3.2 | 2 |
| 136 | A novel prognostic nomogram for colorectal cancer liver metastasis patients with recurrence after hepatectomy. <i>Cancer Medicine</i> , 2021 , 10, 1535-1544 | 4.8 | 2 |
| 135 | Myofibroblast-Specific Msi2 Knockout Inhibits HCC Progression in a Mouse Model. <i>Hepatology</i> , 2021 , 74, 458-473 | 11.2 | 1 |
| 134 | Dynamic monitoring of circulating tumor DNA to predict prognosis and efficacy of adjuvant chemotherapy after resection of colorectal liver metastases. <i>Theranostics</i> , 2021 , 11, 7018-7028 | 12.1 | 6 |

| | | | |
|-----|---|------|----|
| 133 | Primary tumor location affects recurrence-free survival for patients with colorectal liver metastases after hepatectomy: a propensity score matching analysis. <i>World Journal of Surgical Oncology</i> , 2020 , 18, 98 | 3.4 | 1 |
| 132 | Histopathological growth patterns correlate with the immunoscore in colorectal cancer liver metastasis patients after hepatectomy. <i>Cancer Immunology, Immunotherapy</i> , 2020 , 69, 2623-2634 | 7.4 | 5 |
| 131 | PIM2 promotes hepatocellular carcinoma tumorigenesis and progression through activating NF- κ B signaling pathway. <i>Cell Death and Disease</i> , 2020 , 11, 510 | 9.8 | 10 |
| 130 | Author response to Letter to the Editor: Are inflammation-based models feasible tools in predicting the outcome of patients with hepatocellular carcinoma? <i>Liver International</i> , 2020 , 40, 1499-1500 | 7.0 | 7 |
| 129 | A hepatocyte differentiation model reveals two subtypes of liver cancer with different oncofetal properties and therapeutic targets. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020 , 117, 6103-6113 | 11.5 | 17 |
| 128 | Transcatheter arterial chemoembolization alone or combined with ablation for recurrent intermediate-stage hepatocellular carcinoma: a propensity score matching study. <i>Journal of Cancer Research and Clinical Oncology</i> , 2020 , 146, 2669-2680 | 4.9 | 3 |
| 127 | Comparison of the prognostic value of inflammation-based scores in early recurrent hepatocellular carcinoma after hepatectomy. <i>Liver International</i> , 2020 , 40, 229-239 | 7.9 | 26 |
| 126 | PDSS2-Del2, a new variant of PDSS2, promotes tumor cell metastasis and angiogenesis in hepatocellular carcinoma via activating NF- κ B. <i>Molecular Oncology</i> , 2020 , 14, 3184-3197 | 7.9 | 7 |
| 125 | Increased dopamine and its receptor dopamine receptor D1 promote tumor growth in human hepatocellular carcinoma. <i>Cancer Communications</i> , 2020 , 40, 694-710 | 9.4 | 9 |
| 124 | Amide-type local anesthetics may suppress tumor cell proliferation and sensitize Human Hepatocellular Carcinoma Cells to Cisplatin upregulation of expression and demethylation. <i>Journal of Cancer</i> , 2020 , 11, 7312-7319 | 4.5 | 2 |
| 123 | Apatinib versus sorafenib in patients with advanced hepatocellular carcinoma: a preliminary study. <i>Annals of Translational Medicine</i> , 2020 , 8, 1000 | 3.2 | 7 |
| 122 | Dysregulated Sp1/miR-130b-3p/HOXA5 axis contributes to tumor angiogenesis and progression of hepatocellular carcinoma. <i>Theranostics</i> , 2020 , 10, 5209-5224 | 12.1 | 23 |
| 121 | Microwave ablation resection for hepatocellular carcinoma within the Milan criteria: a propensity-score analysis. <i>Therapeutic Advances in Medical Oncology</i> , 2019 , 11, 1758835919874652 | 5.4 | 12 |
| 120 | Lymphoid enhancer-binding factor-1 promotes stemness and poor differentiation of hepatocellular carcinoma by directly activating the NOTCH pathway. <i>Oncogene</i> , 2019 , 38, 4061-4074 | 9.2 | 19 |
| 119 | Lipiodol deposition in portal vein tumour thrombus predicts treatment outcome in HCC patients after transarterial chemoembolisation. <i>European Radiology</i> , 2019 , 29, 5752-5762 | 8 | 15 |
| 118 | LINC01554-Mediated Glucose Metabolism Reprogramming Suppresses Tumorigenicity in Hepatocellular Carcinoma via Downregulating PKM2 Expression and Inhibiting Akt/mTOR Signaling Pathway. <i>Theranostics</i> , 2019 , 9, 796-810 | 12.1 | 71 |
| 117 | The heterogeneity of plasma miRNA profiles in hepatocellular carcinoma patients and the exploration of diagnostic circulating miRNAs for hepatocellular carcinoma. <i>PLoS ONE</i> , 2019 , 14, e02115817 | 3.7 | 10 |
| 116 | Resection vs Ablation for Multifocal Hepatocellular Carcinomas meeting the Barcelona-Clinic Liver Cancer A Classification: A Propensity Score Matching Study. <i>Journal of Cancer</i> , 2019 , 10, 2857-2867 | 4.5 | 4 |

| | | | |
|-----|---|------|-----|
| 115 | The mu-opioid receptor is a molecular marker for poor prognosis in hepatocellular carcinoma and represents a potential therapeutic target. <i>British Journal of Anaesthesia</i> , 2019 , 122, e157-e167 | 5.4 | 29 |
| 114 | Hepatoma cell-secreted exosomal microRNA-103 increases vascular permeability and promotes metastasis by targeting junction proteins. <i>Hepatology</i> , 2018 , 68, 1459-1475 | 11.2 | 184 |
| 113 | Nomogram to Predict Survival of Patients With Recurrence of Hepatocellular Carcinoma After Surgery. <i>Clinical Gastroenterology and Hepatology</i> , 2018 , 16, 756-764.e10 | 6.9 | 28 |
| 112 | Germline Duplication of SNORA18L5 Increases Risk for HBV-related Hepatocellular Carcinoma by Altering Localization of Ribosomal Proteins and Decreasing Levels of p53. <i>Gastroenterology</i> , 2018 , 155, 542-556 | 13.3 | 39 |
| 111 | Hepatocellular Carcinoma Cell-Secreted Exosomal MicroRNA-210 Promotes Angiogenesis In Vitro and In Vivo. <i>Molecular Therapy - Nucleic Acids</i> , 2018 , 11, 243-252 | 10.7 | 118 |
| 110 | Microwave vs radiofrequency ablation for hepatocellular carcinoma within the Milan criteria: a propensity score analysis. <i>Alimentary Pharmacology and Therapeutics</i> , 2018 , 48, 671-681 | 6.1 | 44 |
| 109 | Resection versus Resection with Preoperative Transcatheter Arterial Chemoembolization for Resectable Hepatocellular Carcinoma Recurrence. <i>Journal of Cancer</i> , 2018 , 9, 2778-2785 | 4.5 | 3 |
| 108 | The Immunoscore system predicts prognosis after liver metastasectomy in colorectal cancer liver metastases. <i>Cancer Immunology, Immunotherapy</i> , 2018 , 67, 435-444 | 7.4 | 31 |
| 107 | Overexpression of MUC13, a Poor Prognostic Predictor, Promotes Cell Growth by Activating Wnt Signaling in Hepatocellular Carcinoma. <i>American Journal of Pathology</i> , 2018 , 188, 378-391 | 5.8 | 20 |
| 106 | The efficacy and safety of long- versus short-interval transarterial chemoembolization in unresectable hepatocellular carcinoma. <i>Journal of Cancer</i> , 2018 , 9, 4000-4008 | 4.5 | 6 |
| 105 | Impact of follow-up interval on patients with hepatocellular carcinoma after curative ablation. <i>BMC Cancer</i> , 2018 , 18, 1186 | 4.8 | 7 |
| 104 | Letter: is microwave ablation superior to radiofrequency ablation for early stage hepatocellular carcinoma? AuthorsReply. <i>Alimentary Pharmacology and Therapeutics</i> , 2018 , 48, 1326-1327 | 6.1 | |
| 103 | Long- versus short-interval follow-up after resection of hepatocellular carcinoma: a retrospective cohort study. <i>Cancer Communications</i> , 2018 , 38, 26 | 9.4 | 8 |
| 102 | Letter: microwave vs radiofrequency ablation for hepatocellular carcinoma within the Milan criteria-AuthorsReply. <i>Alimentary Pharmacology and Therapeutics</i> , 2018 , 48, 1027-1028 | 6.1 | |
| 101 | PRMT6 Regulates RAS/RAF Binding and MEK/ERK-Mediated Cancer Stemness Activities in Hepatocellular Carcinoma through CRAF Methylation. <i>Cell Reports</i> , 2018 , 25, 690-701.e8 | 10.6 | 53 |
| 100 | In Reply. <i>Anesthesiology</i> , 2018 , 128, 423 | 4.3 | |
| 99 | Deficiency Induces Hepatocarcinogenesis by Decreasing Mitochondrial Respiration and Reprogramming Glucose Metabolism. <i>Cancer Research</i> , 2018 , 78, 4471-4481 | 10.1 | 16 |
| 98 | CpG Methylation Signature Predicts Recurrence in Early-Stage Hepatocellular Carcinoma: Results From a Multicenter Study. <i>Journal of Clinical Oncology</i> , 2017 , 35, 734-742 | 2.2 | 90 |

| | | | |
|----|--|------|-----|
| 97 | Sorafenib therapy following resection prolongs disease-free survival in patients with advanced hepatocellular carcinoma at a high risk of recurrence. <i>Oncology Letters</i> , 2017 , 13, 984-992 | 2.6 | 13 |
| 96 | Lidocaine Induces Apoptosis and Suppresses Tumor Growth in Human Hepatocellular Carcinoma Cells In Vitro and in a Xenograft Model In Vivo. <i>Anesthesiology</i> , 2017 , 126, 868-881 | 4.3 | 70 |
| 95 | Calcium-binding protein 39 promotes hepatocellular carcinoma growth and metastasis by activating extracellular signal-regulated kinase signaling pathway. <i>Hepatology</i> , 2017 , 66, 1529-1545 | 11.2 | 35 |
| 94 | Pathologic response after preoperative therapy predicts prognosis of Chinese colorectal cancer patients with liver metastases. <i>Chinese Journal of Cancer</i> , 2017 , 36, 78 | | 7 |
| 93 | The Role of Adjuvant Chemotherapy for Colorectal Liver Metastasectomy after Pre-Operative Chemotherapy: Is the Treatment Worthwhile?. <i>Journal of Cancer</i> , 2017 , 8, 1179-1186 | 4.5 | 11 |
| 92 | The Prognostic Value of Peripheral Benzodiazepine Receptor in Patients with Esophageal Squamous Cell Carcinoma. <i>Journal of Cancer</i> , 2017 , 8, 3343-3355 | 4.5 | 1 |
| 91 | Mutual Regulation of MiR-199a-5p and HIF-1 α Modulates the Warburg Effect in Hepatocellular Carcinoma. <i>Journal of Cancer</i> , 2017 , 8, 940-949 | 4.5 | 30 |
| 90 | Reply to A. Braillon. <i>Journal of Clinical Oncology</i> , 2017 , 35, 2098-2099 | 2.2 | |
| 89 | Interacts with Integrin $\alpha 5$ to Suppress HCC Angiogenesis and Metastasis by Inhibiting JAK2/STAT3 Signaling. <i>Cancer Research</i> , 2017 , 77, 5831-5845 | 10.1 | 51 |
| 88 | TP53INP1 Downregulation Activates a p73-Dependent DUSP10/ERK Signaling Pathway to Promote Metastasis of Hepatocellular Carcinoma. <i>Cancer Research</i> , 2017 , 77, 4602-4612 | 10.1 | 27 |
| 87 | Long non-coding RNA UICLM promotes colorectal cancer liver metastasis by acting as a ceRNA for microRNA-215 to regulate ZEB2 expression. <i>Theranostics</i> , 2017 , 7, 4836-4849 | 12.1 | 206 |
| 86 | Preoperative lymphocyte-to-monocyte ratio represents a superior predictor compared with neutrophil-to-lymphocyte and platelet-to-lymphocyte ratios for colorectal liver-only metastases survival. <i>OncoTargets and Therapy</i> , 2017 , 10, 3789-3799 | 4.4 | 23 |
| 85 | Response to Ablation and resection for hepatocellular carcinoma within the Milan criteria and high alpha-fetoprotein levels. <i>Liver International</i> , 2016 , 36, 1878 | 7.9 | |
| 84 | Octamer 4/microRNA-1246 signaling axis drives Wnt/ β -catenin activation in liver cancer stem cells. <i>Hepatology</i> , 2016 , 64, 2062-2076 | 11.2 | 122 |
| 83 | Bevacizumab with preoperative chemotherapy versus preoperative chemotherapy alone for colorectal cancer liver metastases: a retrospective cohort study. <i>Medicine (United States)</i> , 2016 , 95, e4767 ^{1.8} | | 3 |
| 82 | Safety and efficacy of sorafenib therapy in patients with hepatocellular carcinoma: final outcome from the Chinese patient subset of the GIDEON study. <i>Oncotarget</i> , 2016 , 7, 6639-48 | 3.3 | 10 |
| 81 | Nomograms for Pre- and Postoperative Prediction of Long-term Survival for Patients Who Underwent Hepatectomy for Multiple Hepatocellular Carcinomas. <i>Annals of Surgery</i> , 2016 , 263, 778-86 | 7.8 | 46 |
| 80 | Preoperative mean corpuscular hemoglobin affecting long-term outcomes of hepatectomized patients with hepatocellular carcinoma. <i>Molecular and Clinical Oncology</i> , 2016 , 4, 229-236 | 1.6 | 4 |

| | | | |
|----|--|------|-----|
| 79 | CHD1L promotes lineage reversion of hepatocellular carcinoma through opening chromatin for key developmental transcription factors. <i>Hepatology</i> , 2016 , 63, 1544-59 | 11.2 | 26 |
| 78 | Resection vs. ablation for alpha-fetoprotein positive hepatocellular carcinoma within the Milan criteria: a propensity score analysis. <i>Liver International</i> , 2016 , 36, 1677-1687 | 7.9 | 15 |
| 77 | MicroRNAs miR-125b and miR-100 suppress metastasis of hepatocellular carcinoma by disrupting the formation of vessels that encapsulate tumour clusters. <i>Journal of Pathology</i> , 2016 , 240, 450-460 | 9.4 | 47 |
| 76 | High preoperative serum CA19-9 level is predictive of poor prognosis for patients with colorectal liver oligometastases undergoing hepatic resection. <i>Medical Oncology</i> , 2016 , 33, 121 | 3.7 | 18 |
| 75 | A serum microRNA classifier for early detection of hepatocellular carcinoma: a multicentre, retrospective, longitudinal biomarker identification study with a nested case-control study. <i>Lancet Oncology</i> , 2015 , 16, 804-15 | 21.7 | 194 |
| 74 | Systemic delivery of microRNA-101 potentially inhibits hepatocellular carcinoma in vivo by repressing multiple targets. <i>PLoS Genetics</i> , 2015 , 11, e1004873 | 6 | 76 |
| 73 | Loss of ATOH8 Increases Stem Cell Features of Hepatocellular Carcinoma Cells. <i>Gastroenterology</i> , 2015 , 149, 1068-81.e5 | 13.3 | 40 |
| 72 | ANXA3/JNK Signaling Promotes Self-Renewal and Tumor Growth, and Its Blockade Provides a Therapeutic Target for Hepatocellular Carcinoma. <i>Stem Cell Reports</i> , 2015 , 5, 45-59 | 8 | 58 |
| 71 | The role of clinically significant portal hypertension in hepatic resection for hepatocellular carcinoma patients: a propensity score matching analysis. <i>BMC Cancer</i> , 2015 , 15, 263 | 4.8 | 15 |
| 70 | HBP21, a chaperone of heat shock protein 70, functions as a tumor suppressor in hepatocellular carcinoma. <i>Carcinogenesis</i> , 2015 , 36, 1111-20 | 4.6 | 10 |
| 69 | Resection versus ablation in hepatitis B virus-related hepatocellular carcinoma patients with portal hypertension: A propensity score matching study. <i>Surgery</i> , 2015 , 158, 1235-43 | 3.6 | 10 |
| 68 | Hepatectomy Versus Hepatectomy With Lymphadenectomy in Hepatocellular Carcinoma: A Prospective, Randomized Controlled Clinical Trial. <i>Journal of Clinical Gastroenterology</i> , 2015 , 49, 520-8 | 3 | 8 |
| 67 | A novel vascular pattern promotes metastasis of hepatocellular carcinoma in an epithelial-mesenchymal transition-independent manner. <i>Hepatology</i> , 2015 , 62, 452-65 | 11.2 | 88 |
| 66 | Impact of oral anti-hepatitis B therapy on the survival of patients with hepatocellular carcinoma initially treated with chemoembolization. <i>Chinese Journal of Cancer</i> , 2015 , 34, 205-16 | | 16 |
| 65 | Overexpression of N-terminal kinase like gene promotes tumorigenicity of hepatocellular carcinoma by regulating cell cycle progression and cell motility. <i>Oncotarget</i> , 2015 , 6, 1618-30 | 3.3 | 9 |
| 64 | Checkpoint kinase 1 is negatively regulated by miR-497 in hepatocellular carcinoma. <i>Medical Oncology</i> , 2014 , 31, 844 | 3.7 | 53 |
| 63 | MicroRNA-130a is down-regulated in hepatocellular carcinoma and associates with poor prognosis. <i>Medical Oncology</i> , 2014 , 31, 230 | 3.7 | 41 |
| 62 | Allele-specific imbalance of oxidative stress-induced growth inhibitor 1 associates with progression of hepatocellular carcinoma. <i>Gastroenterology</i> , 2014 , 146, 1084-96 | 13.3 | 26 |

| | | | |
|----|---|------|-----|
| 61 | Upregulation of microRNA-106b is associated with poor prognosis in hepatocellular carcinoma. <i>Diagnostic Pathology</i> , 2014 , 9, 226 | 3 | 19 |
| 60 | Long-term survival after resection of hepatocellular carcinoma: a potential risk associated with the choice of postoperative analgesia. <i>Anesthesia and Analgesia</i> , 2014 , 118, 1309-16 | 3.9 | 12 |
| 59 | Phase I trial of hepatic arterial infusion (HAI) of floxuridine with modified oxaliplatin, 5-fluorouracil and leucovorin (m-FOLFOX6) in Chinese patients with unresectable liver metastases from colorectal cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2014 , 74, 1079-87 | 3.5 | 12 |
| 58 | Partial hepatectomy for liver metastases from nasopharyngeal carcinoma: a comparative study and review of the literature. <i>BMC Cancer</i> , 2014 , 14, 818 | 4.8 | 17 |
| 57 | A disrupted RNA editing balance mediated by ADARs (Adenosine DeAminases that act on RNA) in human hepatocellular carcinoma. <i>Gut</i> , 2014 , 63, 832-43 | 19.2 | 136 |
| 56 | Expression of variant isoforms of the tyrosine kinase SYK determines the prognosis of hepatocellular carcinoma. <i>Cancer Research</i> , 2014 , 74, 1845-56 | 10.1 | 28 |
| 55 | Maelstrom promotes hepatocellular carcinoma metastasis by inducing epithelial-mesenchymal transition by way of Akt/GSK-3 β /Snail signaling. <i>Hepatology</i> , 2014 , 59, 531-43 | 11.2 | 98 |
| 54 | Regulatory role of miR-142-3p on the functional hepatic cancer stem cell marker CD133. <i>Oncotarget</i> , 2014 , 5, 5725-35 | 3.3 | 61 |
| 53 | Identification of medium-sized genomic deletions with low coverage, mate-paired restricted tags. <i>BMC Genomics</i> , 2013 , 14, 51 | 4.5 | 1 |
| 52 | Hypermethylation and prognostic implication of Syk gene in human colorectal cancer. <i>Medical Oncology</i> , 2013 , 30, 586 | 3.7 | 17 |
| 51 | MicroRNA-195 suppresses angiogenesis and metastasis of hepatocellular carcinoma by inhibiting the expression of VEGF, VAV2, and CDC42. <i>Hepatology</i> , 2013 , 58, 642-53 | 11.2 | 168 |
| 50 | Genome-wide mutational signatures of aristolochic acid and its application as a screening tool. <i>Science Translational Medicine</i> , 2013 , 5, 197ra101 | 17.5 | 194 |
| 49 | Characterization of the oncogenic function of centromere protein F in hepatocellular carcinoma. <i>Biochemical and Biophysical Research Communications</i> , 2013 , 436, 711-8 | 3.4 | 50 |
| 48 | SPOCK1 is regulated by CHD1L and blocks apoptosis and promotes HCC cell invasiveness and metastasis in mice. <i>Gastroenterology</i> , 2013 , 144, 179-191.e4 | 13.3 | 75 |
| 47 | Recoding RNA editing of AZIN1 predisposes to hepatocellular carcinoma. <i>Nature Medicine</i> , 2013 , 19, 209-16 | 50.5 | 313 |
| 46 | Clinical significance and prognostic value of microRNA expression signatures in hepatocellular carcinoma. <i>Clinical Cancer Research</i> , 2013 , 19, 4780-91 | 12.9 | 88 |
| 45 | MicroRNA-125b promotes apoptosis by regulating the expression of Mcl-1, Bcl-w and IL-6R. <i>Oncogene</i> , 2013 , 32, 3071-9 | 9.2 | 154 |
| 44 | Effects of antiviral therapy on hepatitis B virus reactivation and liver function after resection or chemoembolization for hepatocellular carcinoma. <i>Liver International</i> , 2013 , 33, 595-604 | 7.9 | 63 |

| | | | |
|----|---|------|-----|
| 43 | Efficacy and safety of thermal ablation in patients with liver metastases. <i>European Journal of Gastroenterology and Hepatology</i> , 2013 , 25, 442-6 | 2.2 | 31 |
| 42 | MACC1 as a prognostic biomarker for early-stage and AFP-normal hepatocellular carcinoma. <i>PLoS ONE</i> , 2013 , 8, e64235 | 3.7 | 28 |
| 41 | A modified radiofrequency ablation approach for treating distant lymph node metastasis in two patients with late-stage cancer. <i>Chinese Journal of Cancer</i> , 2013 , 32, 567-70 | | 2 |
| 40 | Translational control of tumor protein induces mitotic defects and chromosome missegregation in hepatocellular carcinoma development. <i>Hepatology</i> , 2012 , 55, 491-505 | 11.2 | 62 |
| 39 | TGF- β 1-miR-34a-CCL22 signaling-induced Treg cell recruitment promotes venous metastases of HBV-positive hepatocellular carcinoma. <i>Cancer Cell</i> , 2012 , 22, 291-303 | 24.3 | 372 |
| 38 | Liver hypertrophy and accelerated growth of implanted tumors in nonembolized liver of rabbit after left portal vein embolization. <i>Journal of Surgical Research</i> , 2012 , 178, 255-63 | 2.5 | 5 |
| 37 | Clinical features and outcome of multiple primary malignancies involving hepatocellular carcinoma: a long-term follow-up study. <i>BMC Cancer</i> , 2012 , 12, 148 | 4.8 | 11 |
| 36 | The beta2-adrenergic receptor is a potential prognostic biomarker for human hepatocellular carcinoma after curative resection. <i>Annals of Surgical Oncology</i> , 2012 , 19, 3556-65 | 3.1 | 32 |
| 35 | Interleukin 23 promotes hepatocellular carcinoma metastasis via NF-kappa B induced matrix metalloproteinase 9 expression. <i>PLoS ONE</i> , 2012 , 7, e46264 | 3.7 | 58 |
| 34 | Serum and glucocorticoid kinase 3 at 8q13.1 promotes cell proliferation and survival in hepatocellular carcinoma. <i>Hepatology</i> , 2012 , 55, 1754-65 | 11.2 | 31 |
| 33 | Expression and prognostic significance of CIP2A mRNA in hepatocellular carcinoma and nontumoral liver tissues. <i>Biomarkers</i> , 2012 , 17, 422-9 | 2.6 | 14 |
| 32 | Hepatectomy for hepatocellular carcinoma patients with macronodular cirrhosis. <i>European Journal of Gastroenterology and Hepatology</i> , 2012 , 24, 575-82 | 2.2 | 8 |
| 31 | CHK1 targets spleen tyrosine kinase (L) for proteolysis in hepatocellular carcinoma. <i>Journal of Clinical Investigation</i> , 2012 , 122, 2165-75 | 15.9 | 84 |
| 30 | Upregulator of cell proliferation predicts poor prognosis in hepatocellular carcinoma and contributes to hepatocarcinogenesis by downregulating FOXO3a. <i>PLoS ONE</i> , 2012 , 7, e40607 | 3.7 | 33 |
| 29 | Allele loss and down-regulation of heparanase gene are associated with the progression and poor prognosis of hepatocellular carcinoma. <i>PLoS ONE</i> , 2012 , 7, e44061 | 3.7 | 10 |
| 28 | LIN28 expression and prognostic value in hepatocellular carcinoma patients who meet the Milan criteria and undergo hepatectomy. <i>Chinese Journal of Cancer</i> , 2012 , 31, 223-32 | | 10 |
| 27 | Novel therapeutic potential in targeting microtubules by nanoparticle albumin-bound paclitaxel in hepatocellular carcinoma. <i>International Journal of Oncology</i> , 2011 , 38, 721-31 | 4.4 | 19 |
| 26 | Changes in hepatitis B virus DNA levels and liver function after transcatheter arterial chemoembolization of hepatocellular carcinoma. <i>Hepatology Research</i> , 2011 , 41, 553-63 | 5.1 | 35 |

| | | | |
|----|--|------|-----|
| 25 | Interleukin 17A promotes hepatocellular carcinoma metastasis via NF- κ B induced matrix metalloproteinases 2 and 9 expression. <i>PLoS ONE</i> , 2011 , 6, e21816 | 3.7 | 131 |
| 24 | Prognostic value of Wnt inhibitory factor-1 expression in hepatocellular carcinoma that is independent of gene methylation. <i>Tumor Biology</i> , 2011 , 32, 233-40 | 2.9 | 11 |
| 23 | NADPH oxidase DUOX1 and DUOX2 but not NOX4 are independent predictors in hepatocellular carcinoma after hepatectomy. <i>Tumor Biology</i> , 2011 , 32, 1173-82 | 2.9 | 24 |
| 22 | Identification of MACC1 as a novel prognostic marker in hepatocellular carcinoma. <i>Journal of Translational Medicine</i> , 2011 , 9, 166 | 8.5 | 67 |
| 21 | Role of Sox2 and Oct4 in predicting survival of hepatocellular carcinoma patients after hepatectomy. <i>Clinical Biochemistry</i> , 2011 , 44, 582-9 | 3.5 | 62 |
| 20 | Clinical significance of CHD1L in hepatocellular carcinoma and therapeutic potentials of virus-mediated CHD1L depletion. <i>Gut</i> , 2011 , 60, 534-43 | 19.2 | 40 |
| 19 | Genome-wide association study identifies 1p36.22 as a new susceptibility locus for hepatocellular carcinoma in chronic hepatitis B virus carriers. <i>Nature Genetics</i> , 2010 , 42, 755-8 | 36.3 | 288 |
| 18 | CpG island methylator phenotype associated with tumor recurrence in tumor-node-metastasis stage I hepatocellular carcinoma. <i>Annals of Surgical Oncology</i> , 2010 , 17, 1917-26 | 3.1 | 37 |
| 17 | Application of tumor-node-metastasis staging 2002 version in locally advanced hepatocellular carcinoma: is it predictive of surgical outcome?. <i>BMC Cancer</i> , 2010 , 10, 535 | 4.8 | 7 |
| 16 | Liver cancer: EphrinA2 promotes tumorigenicity through Rac1/Akt/NF-kappaB signaling pathway. <i>Hepatology</i> , 2010 , 51, 535-44 | 11.2 | 32 |
| 15 | A novel GSK-3 beta-C/EBP alpha-miR-122-insulin-like growth factor 1 receptor regulatory circuitry in human hepatocellular carcinoma. <i>Hepatology</i> , 2010 , 52, 1702-12 | 11.2 | 121 |
| 14 | CHD1L promotes hepatocellular carcinoma progression and metastasis in mice and is associated with these processes in human patients. <i>Journal of Clinical Investigation</i> , 2010 , 120, 1178-91 | 15.9 | 114 |
| 13 | LOH analysis of genes around D4S2964 identifies ARD1B as a prognostic predictor of hepatocellular carcinoma. <i>World Journal of Gastroenterology</i> , 2010 , 16, 2046-54 | 5.6 | 12 |
| 12 | Transarterial chemoembolization as initial treatment for unresectable hepatocellular carcinoma in southern China. <i>World Journal of Gastroenterology</i> , 2010 , 16, 264-9 | 5.6 | 43 |
| 11 | Long-term outcomes and prognostic factors of elderly patients with hepatocellular carcinoma undergoing hepatectomy. <i>Journal of Gastrointestinal Surgery</i> , 2009 , 13, 1627-35 | 3.3 | 97 |
| 10 | MicroRNA-101, down-regulated in hepatocellular carcinoma, promotes apoptosis and suppresses tumorigenicity. <i>Cancer Research</i> , 2009 , 69, 1135-42 | 10.1 | 537 |
| 9 | Large-scale analysis of the genetic and epigenetic alterations in hepatocellular carcinoma from Southeast China. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2008 , 641, 27-35 | 3.3 | 39 |
| 8 | A novel functional polymorphism in the Cdc6 promoter is associated with the risk for hepatocellular carcinoma. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2008 , 643, 70-4 | 3.3 | 12 |

| | | | |
|---|--|------|-----|
| 7 | Clinicopathologic features and long-term outcomes of Chinese patients with hepatocellular carcinoma in non-cirrhotic liver. <i>Digestive Surgery</i> , 2008 , 25, 376-82 | 2.5 | 27 |
| 6 | A functional polymorphism in the miR-146a gene is associated with the risk for hepatocellular carcinoma. <i>Carcinogenesis</i> , 2008 , 29, 2126-31 | 4.6 | 297 |
| 5 | Frequent epigenetic inactivation of spleen tyrosine kinase gene in human hepatocellular carcinoma. <i>Clinical Cancer Research</i> , 2006 , 12, 6687-95 | 12.9 | 59 |
| 4 | Reactivation of SYK expression by inhibition of DNA methylation suppresses breast cancer cell invasiveness. <i>International Journal of Cancer</i> , 2005 , 113, 654-9 | 7.5 | 29 |
| 3 | High-dose iodized oil transcatheter arterial chemoembolization for patients with large hepatocellular carcinoma. <i>World Journal of Gastroenterology</i> , 2002 , 8, 74-8 | 5.6 | 79 |
| 2 | Randomized study of chemoembolization as an adjuvant therapy for primary liver carcinoma after hepatectomy. <i>Journal of Cancer Research and Clinical Oncology</i> , 1995 , 121, 364-6 | 4.9 | 47 |
| 1 | Results of hepatectomy for huge primary liver cancer. <i>Chinese Journal of Cancer Research: Official Journal of China Anti-Cancer Association, Beijing Institute for Cancer Research</i> , 1994 , 6, 91-94 | 3.8 | |