

Margareta Norberg

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8366328/publications.pdf>

Version: 2024-02-01

72
papers

2,165
citations

218677

26
h-index

243625

44
g-index

73
all docs

73
docs citations

73
times ranked

3587
citing authors

#	ARTICLE	IF	CITATIONS
1	Time to initiation of lipid-lowering drugs for subclinical atherosclerosis: sub-study of VIPVIZA randomized controlled trial, with single-arm cross-over. European Heart Journal Open, 2022, 2, .	2.3	2
2	Association of education and feedback on hypertension management with risk for stroke and cardiovascular disease. Blood Pressure, 2022, 31, 31-39.	1.5	2
3	Long-term exposure to particulate air pollution and presence and progression of carotid artery plaques - A northern Sweden VIPVIZA cohort study. Environmental Research, 2022, 211, 113061.	7.5	5
4	No association between moist oral snuff (snus) use and oral cancer: pooled analysis of nine prospective observational studies. Scandinavian Journal of Public Health, 2021, 49, 833-840.	2.3	7
5	Increased knowledge makes a difference! â€“ general practitionersâ€™ experiences of pictorial information about subclinical atherosclerosis for primary prevention: an interview study from the VIPVIZA trial. Scandinavian Journal of Primary Health Care, 2021, 39, 77-84.	1.5	10
6	Quality of life of type 2 diabetes mellitus patients in Ramallah and al-Bireh Governorateâ€“Palestine: a part of the Palestinian diabetes complications and control study (PDCCS). Quality of Life Research, 2021, 30, 1407-1416.	3.1	6
7	Pictorial information about subclinical atherosclerosis reduces the CVD risk: Results from the VIPVIZA RCT. European Journal of Preventive Cardiology, 2021, 28, .	1.8	0
8	The right pick: Does a self-assessment measurement tool correctly identify health care consumers with inadequate health literacy?. Patient Education and Counseling, 2021, , .	2.2	1
9	Prescription of Lipid-Lowering and Antihypertensive Drugs Following Pictorial Information About Subclinical Atherosclerosis. JAMA Network Open, 2021, 4, e2121683.	5.9	7
10	The beneficial effect over 3 years by pictorial information to patients and their physician about subclinical atherosclerosis and cardiovascular risk: Results from the VIPVIZA randomized clinical trial. American Journal of Preventive Cardiology, 2021, 7, 100199.	3.0	21
11	Using Ecological Momentary Assessment to Study the Development of COVID-19 Worries in Sweden: Longitudinal Study. Journal of Medical Internet Research, 2021, 23, e26743.	4.3	3
12	Reduced progression of carotid intima media thickness by pictorial presentation of subclinical atherosclerosis 3-year follow-up from VIPVIZA a randomized controlled trial. European Heart Journal, 2021, 42, .	2.2	0
13	Health literacy is independently and inversely associated with carotid artery plaques and cardiovascular risk. European Journal of Preventive Cardiology, 2020, 27, 209-215.	1.8	20
14	Association of Physician Education and Feedback on Hypertension Management With Patient Blood Pressure and Hypertension Control. JAMA Network Open, 2020, 3, e1918625.	5.9	16
15	Lifetime risk predictions for cardiovascular diseases: Competing risks analyses on a population-based cohort in Sweden. Atherosclerosis, 2020, 312, 90-98.	0.8	11
16	Vitamin D status was not associated with anxiety, depression, or health-related quality of life in Middle Eastern and African-born immigrants in Sweden. Nutrition Research, 2020, 75, 109-118.	2.9	2
17	Are the resources adoptive for conducting team-based diabetes management clinics? An explorative study at primary health care centers in Muscat, Oman. Primary Health Care Research and Development, 2019, 20, e3.	1.2	6
18	Is intima-media thickness a predictor for cardiovascular risk? â€“ Authors' reply. Lancet, The, 2019, 394, 381.	13.7	1

#	ARTICLE	IF	CITATIONS
19	Sex-specific associations between body mass index and death before life expectancy: a comparative study from the USA and Sweden. <i>Global Health Action</i> , 2019, 12, 1580973.	1.9	4
20	Flawed conclusions on the VÄsterbotten Intervention Program by San Sebastian et .al. <i>BMC Public Health</i> , 2019, 19, 1095.	2.9	0
21	Patient“doctor engagement in cardiovascular prevention “ Authors' reply. <i>Lancet, The</i> , 2019, 394, e27.	13.7	0
22	Association between changes in waist circumferences and disability among older adults: WHO-INDEPTH study on global ageing and adult health (SAGE) in Indonesia. <i>Obesity Research and Clinical Practice</i> , 2019, 13, 462-468.	1.8	6
23	Arterial hypertension and diastolic blood pressure associate with aortic stenosis. <i>Scandinavian Cardiovascular Journal</i> , 2019, 53, 91-97.	1.2	6
24	Waist Circumference and All-Cause Mortality among Older Adults in Rural Indonesia. <i>International Journal of Environmental Research and Public Health</i> , 2019, 16, 116.	2.6	13
25	Visualization of asymptomatic atherosclerotic disease for optimum cardiovascular prevention (VIPVIZA): a pragmatic, open-label, randomised controlled trial. <i>Lancet, The</i> , 2019, 393, 133-142.	13.7	142
26	A cost-effectiveness analysis of a community based CVD program in Sweden based on a retrospective register cohort. <i>BMC Public Health</i> , 2018, 18, 452.	2.9	12
27	Separate and combined associations of obesity and metabolic health with coronary heart disease: a pan-European case-cohort analysis. <i>European Heart Journal</i> , 2018, 39, 397-406.	2.2	209
28	Complications of type 2 diabetes mellitus in Ramallah and al-Bireh: The Palestinian Diabetes Complications and Control Study (PDCCS). <i>Primary Care Diabetes</i> , 2018, 12, 547-557.	1.8	16
29	Vitamin D is associated with lower limb muscle strength and grip strength in Middle Eastern- and African-born immigrants in Sweden. <i>Nutrition Research</i> , 2018, 59, 29-35.	2.9	13
30	Association of Pregnancy Complications and Characteristics With Future Risk of Elevated Blood Pressure. <i>Hypertension</i> , 2017, 69, 475-483.	2.7	51
31	Smokeless tobacco (snus) is associated with an increased risk of type 2 diabetes: results from five pooled cohorts. <i>Journal of Internal Medicine</i> , 2017, 281, 398-406.	6.0	34
32	Impact of weight maintenance and loss on diabetes risk and burden: a population-based study in 33,184 participants. <i>BMC Public Health</i> , 2017, 17, 170.	2.9	40
33	Tobacco use is not associated with groin hernia repair, a population-based study. <i>Hernia: the Journal of Hernias and Abdominal Wall Surgery</i> , 2017, 21, 517-523.	2.0	9
34	Use of moist oral snuff (snus) and pancreatic cancer: Pooled analysis of nine prospective observational studies. <i>International Journal of Cancer</i> , 2017, 141, 687-693.	5.1	22
35	Change in lifestyle behaviors and diabetes risk: evidence from a population-based cohort study with 10Åyear follow-up. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2017, 14, 39.	4.6	40
36	The cost-effectiveness of interventions targeting lifestyle change for the prevention of diabetes in a Swedish primary care and community based prevention program. <i>European Journal of Health Economics</i> , 2017, 18, 905-919.	2.8	29

#	ARTICLE	IF	CITATIONS
37	Traditional Cardiovascular Risk Factors and Their Relation to Future Surgery for Valvular Heart Disease or Ascending Aortic Disease: A Case–Referent Study. Journal of the American Heart Association, 2017, 6, .	3.7	26
38	Smokeless tobacco (snus) use and colorectal cancer incidence and survival: Results from nine pooled cohorts. Scandinavian Journal of Public Health, 2017, 45, 741-748.	2.3	7
39	Screening for type 2 diabetes: do screen-detected cases fare better?. Diabetologia, 2017, 60, 2200-2209.	6.3	30
40	Lipoprotein(a) and the Apolipoprotein B/A1 Ratio Independently Associate With Surgery for Aortic Stenosis Only in Patients With Concomitant Coronary Artery Disease. Journal of the American Heart Association, 2017, 6, .	3.7	23
41	Leptin independently predicts development of sepsis and its outcome. Journal of Inflammation, 2017, 14, 19.	3.4	17
42	Socioeconomic inequality in abdominal obesity among older people in Purworejo District, Central Java, Indonesia – a decomposition analysis approach. International Journal for Equity in Health, 2017, 16, 214.	3.5	13
43	Fish consumption and risk of stroke: a second prospective case-control study from northern Sweden. Nutrition Journal, 2016, 15, 98.	3.4	5
44	Prevalence of vitamin D deficiency and its association with nutrition, travelling and clothing habits in an immigrant population in Northern Sweden. European Journal of Clinical Nutrition, 2016, 70, 373-379.	2.9	31
45	From efficacy in trials to effectiveness in clinical practice: The Swedish Stroke Prevention Study. Blood Pressure, 2016, 25, 206-211.	1.5	3
46	Impact of a combined community and primary care prevention strategy on all-cause and cardiovascular mortality: a cohort analysis based on 1 million person-years of follow-up in Västerbotten County, Sweden, during 1990–2006. BMJ Open, 2015, 5, e009651.	1.9	53
47	Comparisons of weight change, eating habits and physical activity between women in Northern Sweden and Rural New York State- results from a longitudinal study. Nutrition Journal, 2015, 14, 88.	3.4	9
48	Adequate vitamin D levels in a Swedish population living above latitude 63°N: The 2009 Northern Sweden MONICA study. International Journal of Circumpolar Health, 2015, 74, 27963.	1.2	34
49	What kills middle-aged Swedes? A Prospective cohort study on NCD risk factors and mortality. European Journal of Public Health, 2015, 25, .	0.3	0
50	Self-rated health and standard risk factors for myocardial infarction: a cohort study. BMJ Open, 2015, 5, e006589-e006589.	1.9	24
51	Healthy behaviours and 10-year incidence of diabetes: A population cohort study. Preventive Medicine, 2015, 71, 121-127.	3.4	34
52	Use of moist smokeless tobacco (snus) and the risk of development of alcohol dependence: A cohort study in a middle-aged population in Sweden. Drug and Alcohol Dependence, 2015, 149, 151-157.	3.2	10
53	Health-related quality of life for pre-diabetic states and type 2 diabetes mellitus: a cross-sectional study in Västerbotten Sweden. Health and Quality of Life Outcomes, 2014, 12, 150.	2.4	37
54	No association found between cardiovascular mortality, and job demands and decision latitude: Experience from the Västerbotten Intervention Programme in Sweden. Social Science and Medicine, 2014, 117, 58-66.	3.8	15

#	ARTICLE	IF	CITATIONS
55	Temporal Shifts in Cardiovascular Risk Factor Distribution. American Journal of Preventive Medicine, 2014, 46, 112-121.	3.0	15
56	A reversal of decreasing trends in population cholesterol levels in VÄsterbotten County, Sweden. Global Health Action, 2012, 5, 10367.	1.9	26
57	Trends of blood pressure levels and management in VÄsterbotten County, Sweden, during 1990â€“2010. Global Health Action, 2012, 5, 18195.	1.9	37
58	Overweight, cardiovascular diseases and diabetes: Health in Sweden: The National Public Health Report 2012. Chapter 7. Scandinavian Journal of Public Health, 2012, 40, 135-163.	2.3	18
59	Community participation and sustainability â€“ evidence over 25 years in the VÄsterbotten Intervention Programme. Global Health Action, 2012, 5, 19166.	1.9	72
60	How to diagnose and classify diabetes in primary health care: Lessons learned from the Diabetes Register in Northern Sweden (DiabNorth). Scandinavian Journal of Primary Health Care, 2012, 30, 81-87.	1.5	26
61	Self-rated health does not predict 10-year weight change among middle-aged adults in a longitudinal population study. BMC Public Health, 2011, 11, 748.	2.9	14
62	Who is using snus? - Time trends, socioeconomic and geographic characteristics of snus users in the ageing Swedish population. BMC Public Health, 2011, 11, 929.	2.9	36
63	Changing patterns of tobacco use in a middle-aged population â€“ the role of snus, gender, age, and education. Global Health Action, 2011, 4, 5613.	1.9	59
64	Increasing physical activity, but persisting social gaps among middle-aged people: trends in Northern Sweden from 1990 to 2007. Global Health Action, 2011, 4, 6347.	1.9	43
65	The obesity epidemic slows among the middle-aged population in Sweden while the socioeconomic gap widens. Global Health Action, 2010, 3, 5149.	1.9	47
66	The VÄsterbotten Intervention Programme: background, design and implications. Global Health Action, 2010, 3, 4643.	1.9	268
67	Recruiting High-Risk Individuals to a Diabetes Prevention Program: How hard can it be?. Diabetes Care, 2007, 30, e61-e61.	8.6	29
68	Work stress and low emotional support is associated with increased risk of future type 2 diabetes in women. Diabetes Research and Clinical Practice, 2007, 76, 368-377.	2.8	104
69	Components of Metabolic Syndrome Predicting Diabetes: No Role of Inflammation or Dyslipidemia. Obesity, 2007, 15, 1875-1885.	3.0	24
70	A combination of HbA1c, fasting glucose and BMI is effective in screening for individuals at risk of future type 2 diabetes: OGTT is not needed. Journal of Internal Medicine, 2006, 260, 263-271.	6.0	71
71	Contribution of Swedish moist snuff to the metabolic syndrome: A wolf in sheep's clothing?. Scandinavian Journal of Public Health, 2006, 34, 576-583.	2.3	53
72	Low Levels of Tissue Inhibitors of Metalloproteinases With a High Matrix Metalloproteinase-9/Tissue Inhibitor of Metalloproteinase-1 Ratio Are Present in Tracheal Aspirate Fluids of Infants Who Develop Chronic Lung Disease. Pediatrics, 2004, 113, 1709-1714.	2.1	86