

# Yoshihisa Tokumaru

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/8369251/yoshihisa-tokumaru-publications-by-year.pdf>  
**Version:** 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.  
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

57 papers	778 citations	18 h-index	26 g-index
64 ext. papers	1,128 ext. citations	5.2 avg, IF	4.93 L-index

#	Paper	IF	Citations
57	Abstract P5-09-03: Mieap, a p53-downstream gene, is associated with suppression of breast cancer cell proliferation and better survival. <i>Cancer Research</i> , <b>2022</b> , 82, P5-09-03-P5-09-03	10.1	
56	High RAD51 gene expression is associated with aggressive biology and with poor survival in breast cancer.. <i>Breast Cancer Research and Treatment</i> , <b>2022</b> , 193, 49	4.4	1
55	Intratumoral density of regulatory T cells is a predictor of host immune response and chemotherapy response in colorectal cancer.. <i>American Journal of Cancer Research</i> , <b>2022</b> , 12, 490-503	4.4	
54	Intratumoral lymphatic endothelial cell infiltration reflecting lymphangiogenesis is counterbalanced by immune responses and better cancer biology in the breast cancer tumor microenvironment.. <i>American Journal of Cancer Research</i> , <b>2022</b> , 12, 504-520	4.4	
53	APOBEC3F expression in triple-negative breast cancer is associated with tumor microenvironment infiltration and activation of cancer immunity and improved survival.. <i>American Journal of Cancer Research</i> , <b>2022</b> , 12, 744-762	4.4	
52	Breast reconstruction using a tissue expander after enucleation of a giant fibroadenoma: A case report.. <i>International Journal of Surgery Case Reports</i> , <b>2021</b> , 90, 106723	0.8	1
51	Angiogenesis is associated with an attenuated tumor microenvironment, aggressive biology, and worse survival in gastric cancer patients. <i>American Journal of Cancer Research</i> , <b>2021</b> , 11, 1659-1671	4.4	5
50	G2M checkpoint pathway alone is associated with drug response and survival among cell proliferation-related pathways in pancreatic cancer. <i>American Journal of Cancer Research</i> , <b>2021</b> , 11, 3070-3084	4.4	2
49	Low expression of miR-195 is associated with cell proliferation, glycolysis and poor survival in estrogen receptor (ER)-positive but not in triple negative breast cancer. <i>American Journal of Cancer Research</i> , <b>2021</b> , 11, 3320-3334	4.4	1
48	A novel five-gene score to predict complete pathological response to neoadjuvant chemotherapy in ER-positive/HER2-negative breast cancer. <i>American Journal of Cancer Research</i> , <b>2021</b> , 11, 3611-3627	4.4	2
47	Increased apoptosis is associated with robust immune cell infiltration and cytolytic activity in breast cancer. <i>American Journal of Cancer Research</i> , <b>2021</b> , 11, 3674-3687	4.4	2
46	Sphingosine 1-phosphate (S1P) produced by sphingosine kinase 1 (SphK1) and exported via ABCC1 is related to hepatocellular carcinoma (HCC) progression. <i>American Journal of Cancer Research</i> , <b>2021</b> , 11, 4394-4407	4.4	1
45	Increased intratumor heterogeneity, angiogenesis and epithelial to mesenchymal transition pathways in metaplastic breast cancer. <i>American Journal of Cancer Research</i> , <b>2021</b> , 11, 4408-4420	4.4	1
44	MELK expression in breast cancer is associated with infiltration of immune cell and pathological complete response (pCR) after neoadjuvant chemotherapy. <i>American Journal of Cancer Research</i> , <b>2021</b> , 11, 4421-4437	4.4	1
43	Abundance of Microvascular Endothelial Cells Is Associated with Response to Chemotherapy and Prognosis in Colorectal Cancer. <i>Cancers</i> , <b>2021</b> , 13,	6.6	6
42	Annexin A1 Expression Is Associated with Epithelial-Mesenchymal Transition (EMT), Cell Proliferation, Prognosis, and Drug Response in Pancreatic Cancer. <i>Cells</i> , <b>2021</b> , 10,	7.9	7
41	A Novel Three-Gene Score as a Predictive Biomarker for Pathologically Complete Response after Neoadjuvant Chemotherapy in Triple-Negative Breast Cancer. <i>Cancers</i> , <b>2021</b> , 13,	6.6	4

40	Urine as a Source of Liquid Biopsy for Cancer. <i>Cancers</i> , <b>2021</b> , 13,	6.6	11
39	Adipogenesis in triple-negative breast cancer is associated with unfavorable tumor immune microenvironment and with worse survival. <i>Scientific Reports</i> , <b>2021</b> , 11, 12541	4.9	5
38	Low expression of miR-29a is associated with aggressive biology and worse survival in gastric cancer. <i>Scientific Reports</i> , <b>2021</b> , 11, 14134	4.9	4
37	Prevalence and clinical relevance of tumor-associated tissue eosinophilia (TATE) in breast cancer. <i>Surgery</i> , <b>2021</b> , 169, 1234-1239	3.6	11
36	Enhanced DNA Repair Pathway is Associated with Cell Proliferation and Worse Survival in Hepatocellular Carcinoma (HCC). <i>Cancers</i> , <b>2021</b> , 13,	6.6	18
35	Th2 cell infiltrations predict neoadjuvant chemotherapy response of estrogen receptor-positive breast cancer. <i>Gland Surgery</i> , <b>2021</b> , 10, 154-165	2.2	13
34	Organoids Are Limited in Modeling the Colon Adenoma-Carcinoma Sequence. <i>Cells</i> , <b>2021</b> , 10,	7.9	7
33	Breast Cancer in Jamaica: Stage, Grade and Molecular Subtype Distributions Across Age Blocks, the Implications for Screening and Treatment. <i>World Journal of Oncology</i> , <b>2021</b> , 12, 93-103	16.7	3
32	Conflicting roles of expression by subtypes in breast cancer. <i>American Journal of Cancer Research</i> , <b>2021</b> , 11, 5094-5110	4.4	1
31	Low intratumoral genetic neutrophil-to-lymphocyte ratio (NLR) is associated with favorable tumor immune microenvironment and with survival in triple negative breast cancer (TNBC). <i>American Journal of Cancer Research</i> , <b>2021</b> , 11, 5743-5755	4.4	
30	, a p53-downstream gene, is associated with suppression of breast cancer cell proliferation and better survival.. <i>American Journal of Cancer Research</i> , <b>2021</b> , 11, 6060-6073	4.4	
29	Inflammation Is Associated with Worse Outcome in the Whole Cohort but with Better Outcome in Triple-Negative Subtype of Breast Cancer Patients. <i>Journal of Immunology Research</i> , <b>2020</b> , 2020, 5618786	4.5	19
28	A Novel Four-Gene Score to Predict Pathologically Complete (R0) Resection and Survival in Pancreatic Cancer. <i>Cancers</i> , <b>2020</b> , 12,	6.6	11
27	High Expression of miR-34a Associated with Less Aggressive Cancer Biology but Not with Survival in Breast Cancer. <i>International Journal of Molecular Sciences</i> , <b>2020</b> , 21,	6.3	31
26	High Expression of microRNA-143 is Associated with Favorable Tumor Immune Microenvironment and Better Survival in Estrogen Receptor Positive Breast Cancer. <i>International Journal of Molecular Sciences</i> , <b>2020</b> , 21,	6.3	29
25	Rectal neuroendocrine tumor developing lateral lymph node metastasis after curative resection: a case report. <i>World Journal of Surgical Oncology</i> , <b>2020</b> , 18, 74	3.4	2
24	Effects of MIR143 on rat sarcoma signaling networks in solid tumors: A brief overview. <i>Cancer Science</i> , <b>2020</b> , 111, 1076-1083	6.9	10
23	Should we target "intermediate expression" of HER2 in older estrogen receptor positive patients?. <i>Translational Cancer Research</i> , <b>2020</b> , 9, 4056-4059	0.3	1

22	KRAS signaling enriched triple negative breast cancer is associated with favorable tumor immune microenvironment and better survival. <i>American Journal of Cancer Research</i> , <b>2020</b> , 10, 897-907	4.4	50
21	Current status and limitations of immunotherapy for breast cancer. <i>Surgery</i> , <b>2020</b> , 167, 628-630	3.6	30
20	CD8 T Cell Score as a Prognostic Biomarker for Triple Negative Breast Cancer. <i>International Journal of Molecular Sciences</i> , <b>2020</b> , 21,	6.3	56
19	Abundance of Regulatory T Cell (Treg) as a Predictive Biomarker for Neoadjuvant Chemotherapy in Triple-Negative Breast Cancer. <i>Cancers</i> , <b>2020</b> , 12,	6.6	39
18	High G2M Pathway Score Pancreatic Cancer is Associated with Worse Survival, Particularly after Margin-Positive (R1 or R2) Resection. <i>Cancers</i> , <b>2020</b> , 12,	6.6	21
17	M1 Macrophage and M1/M2 ratio defined by transcriptomic signatures resemble only part of their conventional clinical characteristics in breast cancer. <i>Scientific Reports</i> , <b>2020</b> , 10, 16554	4.9	45
16	ITPKC as a Prognostic and Predictive Biomarker of Neoadjuvant Chemotherapy for Triple Negative Breast Cancer. <i>Cancers</i> , <b>2020</b> , 12,	6.6	28
15	The E2F Pathway Score as a Predictive Biomarker of Response to Neoadjuvant Therapy in ER+/HER2- Breast Cancer. <i>Cells</i> , <b>2020</b> , 9,	7.9	40
14	Plasmacytoid Dendritic Cell (pDC) Infiltration Correlate with Tumor Infiltrating Lymphocytes, Cancer Immunity, and Better Survival in Triple Negative Breast Cancer (TNBC) More Strongly than Conventional Dendritic Cell (cDC). <i>Cancers</i> , <b>2020</b> , 12,	6.6	34
13	Degree of Early Estrogen Response Predict Survival after Endocrine Therapy in Primary and Metastatic ER-Positive Breast Cancer. <i>Cancers</i> , <b>2020</b> , 12,	6.6	20
12	Transcriptomic Profile of Lymphovascular Invasion, a Known Risk Factor of Pancreatic Ductal Adenocarcinoma Metastasis. <i>Cancers</i> , <b>2020</b> , 12,	6.6	12
11	Intra-Tumoral Angiogenesis Is Associated with Inflammation, Immune Reaction and Metastatic Recurrence in Breast Cancer. <i>International Journal of Molecular Sciences</i> , <b>2020</b> , 21,	6.3	31
10	Intratumoral Adipocyte-High Breast Cancer Enrich for Metastatic and Inflammation-Related Pathways but Associated with Less Cancer Cell Proliferation. <i>International Journal of Molecular Sciences</i> , <b>2020</b> , 21,	6.3	27
9	High Expression of NRF2 Is Associated with Increased Tumor-Infiltrating Lymphocytes and Cancer Immunity in ER-Positive/HER2-Negative Breast Cancer. <i>Cancers</i> , <b>2020</b> , 12,	6.6	21
8	G2M Cell Cycle Pathway Score as a Prognostic Biomarker of Metastasis in Estrogen Receptor (ER)-Positive Breast Cancer. <i>International Journal of Molecular Sciences</i> , <b>2020</b> , 21,	6.3	60
7	Combination chemotherapy with TAS-102 plus bevacizumab in salvage-line treatment of metastatic colorectal cancer: A single-center, retrospective study examining the prognostic value of the modified Glasgow Prognostic Score in salvage-line therapy of metastatic colorectal cancer. <i>Molecular and Clinical Oncology</i> , <b>2019</b> , 11, 390-396	1.6	3
6	MicroRNA-143/Musashi-2/KRAS cascade contributes positively to carcinogenesis in human bladder cancer. <i>Cancer Science</i> , <b>2019</b> , 110, 2189-2199	6.9	14
5	Efficacy of combination therapy with zoledronic acid and cetuximab for unresectable rectal cancer with bone metastases: A case report. <i>Molecular and Clinical Oncology</i> , <b>2019</b> , 10, 571-574	1.6	3

4	Synthetic miR-143 Inhibits Growth of HER2-Positive Gastric Cancer Cells by Suppressing KRAS Networks Including DDX6 RNA Helicase. <i>International Journal of Molecular Sciences</i> , <b>2019</b> , 20,	6.3	14
3	DEAD-Box Protein RNA-Helicase DDX6 Regulates the Expression of HER2 and FGFR2 at the Post-Transcriptional Step in Gastric Cancer Cells. <i>International Journal of Molecular Sciences</i> , <b>2018</b> , 19,	6.3	13
2	The mode of progressive disease affects the prognosis of patients with metastatic breast cancer. <i>World Journal of Surgical Oncology</i> , <b>2018</b> , 16, 169	3.4	3
1	Preoperative Embolization of the Common Hepatic Artery for Radical Surgery of Extrahepatic Cholangiocarcinoma. <i>American Surgeon</i> , <b>2011</b> , 77, 269-271	0.8	2