Hiroaki Fukumoto

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8373163/publications.pdf

Version: 2024-02-01

1163117 1588992 1,745 8 8 8 citations h-index g-index papers 11 11 11 2357 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	$\hat{l}^2\text{-Secretase}$ Protein and Activity Are Increased in the Neocortex in Alzheimer Disease. Archives of Neurology, 2002, 59, 1381.	4.5	656
2	\hat{l}^2 -Secretase Activity Increases with Aging in Human, Monkey, and Mouse Brain. American Journal of Pathology, 2004, 164, 719-725.	3.8	267
3	Age but Not Diagnosis Is the Main Predictor of Plasma Amyloid \hat{l}^2 -Protein Levels. Archives of Neurology, 2003, 60, 958.	4.5	250
4	Highâ€molecularâ€weight βâ€amyloid oligomers are elevated in cerebrospinal fluid of Alzheimer patients. FASEB Journal, 2010, 24, 2716-2726.	0.5	230
5	Induction of the Cholesterol Transporter ABCA1 in Central Nervous System Cells by Liver X Receptor Agonists Increases Secreted A \hat{I}^2 Levels. Journal of Biological Chemistry, 2002, 277, 48508-48513.	3.4	139
6	Plasma Amyloid \hat{l}^2 -Protein and C-reactive Protein in Relation to the Rate of Progression of Alzheimer Disease. Archives of Neurology, 2008, 65, 776-85.	4.5	84
7	APOE ε3/ε4 heterozygotes have an elevated proportion of apolipoprotein E4 in cerebrospinal fluid relative to plasma, independent of Alzheimer's disease diagnosis. Experimental Neurology, 2003, 183, 249-253.	4.1	63
8	Primary cultures of neuronal and non-neuronal rat brain cells secrete similar proportions of amyloid \hat{l}^2 peptides ending at A \hat{l}^2 40 and A \hat{l}^2 42. NeuroReport, 1999, 10, 2965-2969.	1.2	55