

# Hiroaki Fukumoto

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8373163/publications.pdf>

Version: 2024-02-01

8  
papers

1,745  
citations

1162367

8  
h-index

1588620

8  
g-index

11  
all docs

11  
docs citations

11  
times ranked

2357  
citing authors

#	ARTICLE	IF	CITATIONS
1	High-molecular-weight $\beta$ -amyloid oligomers are elevated in cerebrospinal fluid of Alzheimer patients. <i>FASEB Journal</i> , 2010, 24, 2716-2726.	0.2	230
2	Plasma Amyloid $\beta$ -Protein and C-reactive Protein in Relation to the Rate of Progression of Alzheimer Disease. <i>Archives of Neurology</i> , 2008, 65, 776-85.	4.9	84
3	$\beta$ -Secretase Activity Increases with Aging in Human, Monkey, and Mouse Brain. <i>American Journal of Pathology</i> , 2004, 164, 719-725.	1.9	267
4	APOE $\epsilon$ 3/ $\epsilon$ 4 heterozygotes have an elevated proportion of apolipoprotein E4 in cerebrospinal fluid relative to plasma, independent of Alzheimer's disease diagnosis. <i>Experimental Neurology</i> , 2003, 183, 249-253.	2.0	63
5	Age but Not Diagnosis Is the Main Predictor of Plasma Amyloid $\beta$ -Protein Levels. <i>Archives of Neurology</i> , 2003, 60, 958.	4.9	250
6	Induction of the Cholesterol Transporter ABCA1 in Central Nervous System Cells by Liver X Receptor Agonists Increases Secreted $A\beta$ Levels. <i>Journal of Biological Chemistry</i> , 2002, 277, 48508-48513.	1.6	139
7	$\beta$ -Secretase Protein and Activity Are Increased in the Neocortex in Alzheimer Disease. <i>Archives of Neurology</i> , 2002, 59, 1381.	4.9	656
8	Primary cultures of neuronal and non-neuronal rat brain cells secrete similar proportions of amyloid $\beta$ peptides ending at $A\beta$ 40 and $A\beta$ 42. <i>NeuroReport</i> , 1999, 10, 2965-2969.	0.6	55