

Hiroaki Fukumoto

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8373163/publications.pdf>

Version: 2024-02-01

8

papers

1,745

citations

1163117

8

h-index

1588992

8

g-index

11

all docs

11

docs citations

11

times ranked

2357

citing authors

#	ARTICLE	IF	CITATIONS
1	High-molecular-weight β -amyloid oligomers are elevated in cerebrospinal fluid of Alzheimer patients. FASEB Journal, 2010, 24, 2716-2726.	0.5	230
2	Plasma Amyloid β -Protein and C-reactive Protein in Relation to the Rate of Progression of Alzheimer Disease. Archives of Neurology, 2008, 65, 776-85.	4.5	84
3	β -Secretase Activity Increases with Aging in Human, Monkey, and Mouse Brain. American Journal of Pathology, 2004, 164, 719-725.	3.8	267
4	APOE ϵ 3/ ϵ 4 heterozygotes have an elevated proportion of apolipoprotein E4 in cerebrospinal fluid relative to plasma, independent of Alzheimer's disease diagnosis. Experimental Neurology, 2003, 183, 249-253.	4.1	63
5	Age but Not Diagnosis Is the Main Predictor of Plasma Amyloid β -Protein Levels. Archives of Neurology, 2003, 60, 958.	4.5	250
6	Induction of the Cholesterol Transporter ABCA1 in Central Nervous System Cells by Liver X Receptor Agonists Increases Secreted β Levels. Journal of Biological Chemistry, 2002, 277, 48508-48513.	3.4	139
7	β -Secretase Protein and Activity Are Increased in the Neocortex in Alzheimer Disease. Archives of Neurology, 2002, 59, 1381.	4.5	656
8	Primary cultures of neuronal and non-neuronal rat brain cells secrete similar proportions of amyloid β peptides ending at β 40 and β 42. NeuroReport, 1999, 10, 2965-2969.	1.2	55