

Josã© G Barroso

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8380371/publications.pdf>

Version: 2024-02-01

99
papers

4,076
citations

147726

31
h-index

128225

60
g-index

101
all docs

101
docs citations

101
times ranked

4516
citing authors

#	ARTICLE	IF	CITATIONS
1	Essential Oil Variability of Azorean <i>Cryptomeria japonica</i> Leaves under Different Distillation Methods, Part 1: Color, Yield and Chemical Composition Analysis. <i>Applied Sciences (Switzerland)</i> , 2022, 12, 452.	1.3	5
2	Characterization of cuticular compounds of the cerambycid beetles <i>Monochamus galloprovincialis</i> , <i>Arhopalus syriacus</i> , and <i>Pogonocherus perroudi</i> , potential vectors of pinewood nematode. <i>Entomologia Experimentalis Et Applicata</i> , 2021, 169, 183-194.	0.7	3
3	Chemotypes and terpene synthase genes in <i>Thymus</i> genus: State of the art. <i>Industrial Crops and Products</i> , 2018, 124, 530-547.	2.5	35
4	<i>Pinus halepensis</i> , <i>Pinus pinaster</i> , <i>Pinus pinea</i> and <i>Pinus sylvestris</i> Essential Oils Chemotypes and Monoterpene Hydrocarbon Enantiomers, before and after Inoculation with the Pinewood Nematode <i>Bursaphelenchus xylophilus</i> . <i>Chemistry and Biodiversity</i> , 2017, 14, e1600153.	1.0	35
5	Mineral and volatile composition of Açugua-mel from Portugal. <i>European Food Research and Technology</i> , 2016, 242, 171-178.	1.6	3
6	Medicinal and Aromatic Plants (MAP): How Do They Adapt to the Environment?. <i>Medicinal and Aromatic Plants of the World</i> , 2015, , 87-112.	0.1	2
7	Essential oil from <i>Artemisia herba-alba</i> Asso grown wild in Algeria: Variability assessment and comparison with an updated literature survey. <i>Arabian Journal of Chemistry</i> , 2014, 7, 243-251.	2.3	59
8	Identification and characterization of a second isogene encoding β -terpinene synthase in <i>Thymus caespititius</i> . <i>Journal of Plant Physiology</i> , 2014, 171, 1017-1027.	1.6	24
9	Volatile Oils Composition, and Bioactivity of the Essential Oils of <i>Plectranthus barbatus</i> , <i>P. neochilus</i> , and <i>P. ornatus</i> Grown in Portugal. <i>Chemistry and Biodiversity</i> , 2014, 11, 719-732.	1.0	25
10	Yield and chemical composition of the essential oil of Moroccan chamomile [<i>Cladanthus mixtus</i> (L.) Chevall.] growing wild at different sites in Morocco. <i>Flavour and Fragrance Journal</i> , 2013, 28, 360-366.	1.2	15
11	Glandular trichomes, histochemical localization of secretion, and essential oil composition in <i>Plectranthus grandidentatus</i> growing in Portugal. <i>Flavour and Fragrance Journal</i> , 2013, 28, 393-401.	1.2	11
12	Propolis volatiles characterisation from acaricide-treated and -untreated beehives maintained at Algarve (Portugal). <i>Natural Product Research</i> , 2013, 27, 743-749.	1.0	15
13	Genomic characterization, molecular cloning and expression analysis of two terpene synthases from <i>Thymus caespititius</i> (Lamiaceae). <i>Planta</i> , 2013, 238, 191-204.	1.6	41
14	Chemical Composition and Antioxidant Activity of Essential Oils from <i>Cinnamodendron dinisii</i> Schwacke and <i>Siparuna guianensis</i> Aublet. <i>Antioxidants</i> , 2013, 2, 384-397.	2.2	61
15	Chemical variability of the essential oils from <i>Rosa canina</i> L. and <i>Rosa sempervirens</i> L. flowers collected at Tunisia. <i>Journal of Essential Oil Research</i> , 2012, 24, 475-480.	1.3	8
16	Chemical analyses of the essential oils from leaves of <i>Mikania glauca</i> Mart. ex Baker. <i>Journal of Essential Oil Research</i> , 2012, 24, 599-604.	1.3	5
17	Bioassays Against Pinewood Nematode: Assessment of a Suitable Dilution Agent and Screening for Bioactive Essential Oils. <i>Molecules</i> , 2012, 17, 12312-12329.	1.7	30
18	<i>Salvia officinalis</i> L. essential oils: effect of hydrodistillation time on the chemical composition, antioxidant and antimicrobial activities. <i>Natural Product Research</i> , 2011, 25, 526-541.	1.0	48

#	ARTICLE	IF	CITATIONS
19	Antimicrobial activity, cytotoxicity and intracellular growth inhibition of Portuguese Thymus essential oils. <i>Revista Brasileira De Farmacognosia</i> , 2011, 21, 1012-1024.	0.6	18
20	ISSR molecular characterization and leaf volatiles analysis of <i>Pittosporum undulatum</i> Vent. naturalized in the Azores archipelago (Portugal). <i>Industrial Crops and Products</i> , 2011, 33, 710-719.	2.5	19
21	Volatile and molecular analysis of <i>Juniperus brevifolia</i> (Seub.) Antoine, an Azorean endemic species. <i>Biochemical Systematics and Ecology</i> , 2010, 38, 621-629.	0.6	7
22	Antioxidant activity of six Portuguese thyme species essential oils. <i>Flavour and Fragrance Journal</i> , 2010, 25, 150-155.	1.2	60
23	Volatiles from <i>Plicanthus hirtellus</i> (F. Weber) R.M. Schust. and <i>Radula boryana</i> (F. Weber) Nees (Hepaticae) grown in São Tomé e Príncipe Archipelago. <i>Flavour and Fragrance Journal</i> , 2010, 25, 219-222.	1.2	13
24	Composition and antioxidant activity of <i>Thymus vulgaris</i> volatiles: Comparison between supercritical fluid extraction and hydrodistillation. <i>Journal of Separation Science</i> , 2010, 33, 2211-2218.	1.3	69
25	Volatiles from <i>Thymbra</i> and <i>Thymus</i> species of the Western Mediterranean Basin, Portugal and Macaronesia. <i>Natural Product Communications</i> , 2010, 5, 1934578X1000500.	0.2	10
26	<i>Foeniculum vulgare</i> Essential Oils: Chemical Composition, Antioxidant and Antimicrobial Activities. <i>Natural Product Communications</i> , 2010, 5, 1934578X1000500.	0.2	44
27	Herbicidal Activity of Volatiles from Coriander, Winter Savory, Cotton Lavender, and Thyme Isolated by Hydrodistillation and Supercritical Fluid Extraction. <i>Journal of Agricultural and Food Chemistry</i> , 2010, 58, 11007-11013.	2.4	51
28	Nematicidal activity of essential oils and volatiles derived from Portuguese aromatic flora against the pinewood nematode, <i>Bursaphelenchus xylophilus</i> . <i>Journal of Nematology</i> , 2010, 42, 8-16.	0.4	47
29	Volatiles from <i>Thymbra</i> and <i>Thymus</i> species of the western Mediterranean basin, Portugal and Macaronesia. <i>Natural Product Communications</i> , 2010, 5, 1465-76.	0.2	17
30	Menthol and Geraniol Biotransformation and Glycosylation Capacity of <i>Levisticum officinale</i> Hairy Roots. <i>Planta Medica</i> , 2009, 75, 387-391.	0.7	30
31	Volatile and molecular characterization of two Portuguese endemic species: <i>Angelica lignescens</i> and <i>Melanoselinum decipiens</i> . <i>Biochemical Systematics and Ecology</i> , 2009, 37, 98-105.	0.6	12
32	Assessment of the essential oil composition of <i>Tornabenea annua</i> , <i>Tornabenea insularis</i> and <i>Tornabenea tenuissima</i> fruits from Cape Verde Islands. <i>Biochemical Systematics and Ecology</i> , 2009, 37, 474-478.	0.6	4
33	A combined approach using RAPD, ISSR and volatile analysis for the characterization of <i>Thymus caespitius</i> from Flores, Corvo and Graciosa islands (Azores, Portugal). <i>Biochemical Systematics and Ecology</i> , 2009, 37, 670-677.	0.6	35
34	<i>Chaerophyllum azoricum</i> Trel. grown in the Azores archipelago, Portugal: evaluation of the genetic diversity using molecular markers and comparison with volatile oils profiles. <i>Flavour and Fragrance Journal</i> , 2009, 24, 259-265.	1.2	4
35	Liverwort <i>Radula</i> species from Portugal: chemotaxonomical evaluation of volatiles composition. <i>Flavour and Fragrance Journal</i> , 2009, 24, 316-325.	1.2	15
36	Enrichment of the thymoquinone content in volatile oil from <i>Satureja montana</i> using supercritical fluid extraction. <i>Journal of Separation Science</i> , 2009, 32, 328-334.	1.3	43

#	ARTICLE	IF	CITATIONS
37	Supercritical fluid extraction of the volatile oil from <i>Santolina chamaecyparissus</i> . Journal of Separation Science, 2009, 32, 3215-3222.	1.3	24
38	Biotransformation of menthol and geraniol by hairy root cultures of <i>Anethum graveolens</i> : effect on growth and volatile components. Biotechnology Letters, 2009, 31, 897-903.	1.1	24
39	Antioxidant Activities of the Supercritical and Conventional <i>Satureja montana</i> Extracts. Journal of Food Science, 2009, 74, C713-7.	1.5	33
40	Antioxidant Capacity of the Essential Oils From <i>Lavandula luisieri</i> , <i>L. stoechas</i> subsp. <i>lusitanica</i> , <i>L. stoechas</i> subsp. <i>lusitanica</i> x <i>L. luisieri</i> and <i>L. viridis</i> Grown in Algarve (Portugal). Journal of Essential Oil Research, 2009, 21, 327-336.	1.3	41
41	Volatile compounds from the symbiotic system <i>Azolla filiculoides</i> <i>Anabaena azollae</i> bacteria. Plant Biosystems, 2009, 143, 268-274.	0.8	6
42	Nitrogen stress induction on <i>Levisticum officinale</i> hairy roots grown in darkness and under photoperiod conditions: effect on growth and volatile components. Biotechnology Letters, 2008, 30, 1265-1270.	1.1	7
43	Factors affecting secondary metabolite production in plants: volatile components and essential oils. Flavour and Fragrance Journal, 2008, 23, 213-226.	1.2	882
44	Chemotaxonomy of <i>Hypericum</i> genus from Portugal: Geographical distribution and essential oils composition of <i>Hypericum perforatum</i> , <i>Hypericum humifusum</i> , <i>Hypericum linarifolium</i> and <i>Hypericum pulchrum</i> . Biochemical Systematics and Ecology, 2008, 36, 40-50.	0.6	48
45	Genetic diversity and chemical polymorphism of <i>Thymus caespitius</i> from Pico, São Jorge and Terceira islands (Azores). Biochemical Systematics and Ecology, 2008, 36, 790-797.	0.6	47
46	Portuguese <i>Thymbra</i> and <i>Thymus</i> Species Volatiles: Chemical Composition and Biological Activities. Current Pharmaceutical Design, 2008, 14, 3120-3140.	0.9	124
47	Assessment of the Antioxidant Ability of <i>Thymus albicans</i> , <i>T. mastichina</i> , <i>T. camphoratus</i> and <i>T. carnosus</i> Essential Oils by TBARS and Micellar Model systems. Natural Product Communications, 2007, 2, 1934578X0700200.	0.2	11
48	<i>Pittosporum undulatum</i> Vent. grown in Portugal: secretory structures, seasonal variation and enantiomeric composition of its essential oil. Flavour and Fragrance Journal, 2007, 22, 1-9.	1.2	17
49	Composition of the leaf, flower and fruit volatile oils of <i>Pittosporum tobira</i> (Thunb.) W. T. Aiton grown in three locations in Portugal. Flavour and Fragrance Journal, 2007, 22, 311-316.	1.2	8
50	Supercritical carbon dioxide extraction of volatiles from <i>Satureja fruticosa</i> Bâguinot. Flavour and Fragrance Journal, 2007, 22, 438-442.	1.2	21
51	Essential oil composition of <i>Pterospartum tridentatum</i> grown in Portugal. Food Chemistry, 2007, 102, 1083-1088.	4.2	17
52	Essential Oil Composition and Glandular Trichomes of <i>Marrubium vulgare</i> L. Growing Wild in Algeria. Journal of Essential Oil Research, 2006, 18, 369-373.	1.3	28
53	Chemical Composition, Leaf Trichome Types and Biological Activities of the Essential Oils of Four Related <i>Salvia</i> Species Indigenous to Southern Africa. Journal of Essential Oil Research, 2006, 18, 72-79.	1.3	59
54	Biological Activities and Composition of <i>Salvia muirii</i> L. Bol. Essential Oil. Journal of Essential Oil Research, 2006, 18, 48-51.	1.3	11

#	ARTICLE	IF	CITATIONS
55	Simple gas chromatographic method for the stereodifferentiation of methyl nilate, a chiral \pm -methyl- β -hydroxy ester. <i>Journal of Chromatography A</i> , 2006, 1108, 225-230.	1.8	3
56	<i>Asterella africana</i> (Mont.) A. Evans grown on Madeira and in mainland Portugal: morphological data and composition of the essential oil. <i>Flavour and Fragrance Journal</i> , 2006, 21, 534-538.	1.2	7
57	Constituents of the Essential Oil of Sea Fennel (<i>Crithmum maritimum</i> L.) Growing Wild in Turkey. <i>Journal of Medicinal Food</i> , 2006, 9, 128-130.	0.8	21
58	<i>Origanum glandulosum</i> Desf. grown wild in Algeria: essential oil composition and glycosidic bound volatiles. <i>Flavour and Fragrance Journal</i> , 2005, 20, 209-212.	1.2	26
59	Comparison of the essential oil composition of four <i>Plagiochila</i> species: <i>P. bifaria</i> , <i>P. maderensis</i> , <i>P. retrorsa</i> and <i>P. stricta</i> . <i>Flavour and Fragrance Journal</i> , 2005, 20, 703-709.	1.2	15
60	Growth and essential oil composition of hairy root cultures of <i>Levisticum officinale</i> W.D.J. Koch (lovage). <i>Plant Science</i> , 2005, 168, 1089-1096.	1.7	56
61	Chemical polymorphism of populations of <i>Thymus caespititius</i> grown on the islands Corvo, Flores, São Miguel and Terceira (Azores) and on Madeira, assessed by analysis of their essential oils. <i>Plant Science</i> , 2005, 169, 1112-1117.	1.7	26
62	<i>Thymus carnosus</i> Boiss.: Effect of Harvesting Period, Collection Site and Type of Plant Material on Essential Oil Composition. <i>Journal of Essential Oil Research</i> , 2005, 17, 422-426.	1.3	17
63	Antibacterial and Antioxidant Activities of Essential Oils Isolated from <i>Thymbra capitata</i> L. (Cav.) and <i>Origanum vulgare</i> L.. <i>Journal of Agricultural and Food Chemistry</i> , 2005, 53, 8162-8168.	2.4	146
64	Micromorphology of trichomes and composition of essential oil of <i>Teucrium capitatum</i> . <i>Flavour and Fragrance Journal</i> , 2004, 19, 336-340.	1.2	42
65	Effect of the volatile constituents isolated from <i>Thymus albicans</i> , <i>Th. mastichina</i> , <i>Th. carnosus</i> and <i>Thymbra capitata</i> in sunflower oil. <i>Molecular Nutrition and Food Research</i> , 2003, 47, 397-402.	0.0	29
66	Antimicrobial activity of essential oils isolated from Portuguese endemic species of <i>Thymus</i> . <i>Letters in Applied Microbiology</i> , 2003, 36, 35-40.	1.0	154
67	Chemical polymorphism of the essential oils from populations of <i>Thymus caespititius</i> grown on the islands Pico, Faial and Graciosa (Azores). <i>Phytochemical Analysis</i> , 2003, 14, 228-231.	1.2	19
68	Effect of the essential volatile oils isolated from <i>Thymbra capitata</i> (L.) Cav. on olive and sunflower oils. <i>Grasas Y Aceites</i> , 2003, 54, .	0.3	13
69	Composition of the essential oil of <i>Juniperus cedrus</i> Webb & Berth. grown on Madeira. <i>Flavour and Fragrance Journal</i> , 2002, 17, 111-114.	1.2	30
70	Composition of the essential oil and micromorphology of trichomes of <i>Teucrium salviastrum</i> , an endemic species from Portugal. <i>Flavour and Fragrance Journal</i> , 2002, 17, 287-291.	1.2	29
71	Title is missing!. <i>Biotechnology Letters</i> , 2002, 24, 1031-1036.	1.1	34
72	Essential oils from Azorean <i>Laurus azorica</i> . <i>Phytochemistry</i> , 2001, 57, 245-250.	1.4	33

#	ARTICLE	IF	CITATIONS
73	Essential oil composition of <i>Thymus lotocephalus</i> G. Lázpez & R. Morales, collected during flowering and vegetative phases. <i>Flavour and Fragrance Journal</i> , 2001, 16, 417-421.	1.2	12
74	Glandular trichomes and essential oils of <i>Helichrysum stoechas</i> . <i>Israel Journal of Plant Sciences</i> , 2001, 49, 115-122.	0.3	23
75	In vitro evaluation of antioxidant activity of essential oils and their components. <i>Flavour and Fragrance Journal</i> , 2000, 15, 12-16.	1.2	112
76	Essential oils from seven populations of <i>Juniperus brevifolia</i> (Seub.) Antoine, an endemic species of the Azores. <i>Flavour and Fragrance Journal</i> , 2000, 15, 31-39.	1.2	18
77	Chemical polymorphism of the essential oils from populations of <i>Thymus caespititius</i> grown on the island S. Jorge (Azores). <i>Phytochemistry</i> , 2000, 55, 241-246.	1.4	41
78	Title is missing!. <i>Biotechnology Letters</i> , 1999, 21, 859-864.	1.1	16
79	Composition of the essential oil of <i>Hypericum foliosum</i> Aiton from five Azorean islands. <i>Flavour and Fragrance Journal</i> , 1999, 14, 283-286.	1.2	30
80	Composition of the essential oil of <i>Chaerophyllum azoricum</i> Trel., an endemic species of the Azores archipelago. <i>Flavour and Fragrance Journal</i> , 1999, 14, 287-289.	1.2	23
81	Composition of the essential oil of <i>Melanoselinum decipiens</i> (Schrad. & Wendl.) Hoffm., an endemic species of the Madeira and Azores Archipelagos. <i>Flavour and Fragrance Journal</i> , 1998, 13, 90-92.	1.2	2
82	Antimicrobial and antioxidant properties of some commercial essential oils. <i>Flavour and Fragrance Journal</i> , 1998, 13, 235-244.	1.2	449
83	<i>Plectranthus madagascariensis</i> : Morphology of the Glandular Trichomes, Essential Oil Composition, and Its Biological Activity. <i>International Journal of Plant Sciences</i> , 1998, 159, 31-38.	0.6	65
84	Morphology and distribution of trichomes in two endemic <i>Teucrium</i> species of Macaronesia. <i>Acta Botanica Gallica</i> , 1997, 144, 363-369.	0.9	6
85	Composition of the Essential Oil of <i>Monizia edulis</i> Lowe, an Endemic Species of the Madeira Archipelago. <i>Flavour and Fragrance Journal</i> , 1997, 12, 29-31.	1.2	3
86	Composition of the essential oil of <i>Teucrium haenseleri</i> Boiss.. <i>Flavour and Fragrance Journal</i> , 1997, 12, 355-357.	1.2	9
87	Composition of the Essential Oil of <i>Teucrium heterophyllum</i> L'Hér. Grown on Madeira. <i>Flavour and Fragrance Journal</i> , 1996, 11, 129-132.	1.2	15
88	The Essential Oils of Two Endemic <i>Argyranthemum</i> Species of the Madeira Archipelago: <i>A. pinnatifidum</i> (L. fil.) Lowe spp. <i>pinnatifidum</i> and <i>A. haemotomma</i> (Lowe) Lowe. <i>Flavour and Fragrance Journal</i> , 1996, 11, 211-214.	1.2	2
89	Biotransformation of monoterpenes and sesquiterpenes by cell suspension cultures of <i>Achillea millefolium</i> L. ssp. <i>millefolium</i> . <i>Biotechnology Letters</i> , 1996, 18, 863-868.	1.1	24
90	Composition of the Essential Oil of <i>Lavandula pinnata</i> L. fil. var. <i>pinnata</i> grown on madeira. <i>Flavour and Fragrance Journal</i> , 1995, 10, 93-96.	1.2	20

#	ARTICLE	IF	CITATIONS
91	Composition of the essential oil of <i>Micromeria varia</i> Benth. ssp. <i>thymoides</i> (Sol. ex Lowe) Párez var. <i>thymoides</i> , an endemic species of the madeira archipelago. <i>Flavour and Fragrance Journal</i> , 1995, 10, 199-202.	1.2	5
92	Accumulation of stress metabolites in cell suspension cultures of <i>Hyoscyamus albus</i> . <i>Phytochemistry</i> , 1994, 35, 371-375.	1.4	12
93	Composition of the essential oil of <i>artemisia argentea</i> L., an endemic species of the madeira archipelago. <i>Flavour and Fragrance Journal</i> , 1994, 9, 229-232.	1.2	10
94	The essential oils of two endemic Portuguese thyme species: <i>Thymus capitellatus</i> Hoffmanns. & Link and <i>T. lotocephalus</i> G. López & R. Morales. <i>Flavour and Fragrance Journal</i> , 1993, 8, 53-57.	1.2	12
95	The essential oils of two endemic <i>Teucrium</i> species from Madeira: <i>T. abutiloides</i> L'Hér. and <i>T. betonicum</i> L'Hér. <i>Flavour and Fragrance Journal</i> , 1993, 8, 277-280.	1.2	13
96	Composition of the Essential Oils from Two Populations of <i>Achillea millefolium</i> L. ssp. <i>millefolium</i> . <i>Journal of Chromatographic Science</i> , 1992, 30, 392-395.	0.7	22
97	Seasonal Variation in the Composition of the Essential Oil of <i>Crithmum maritimum</i> L.. <i>Flavour and Fragrance Journal</i> , 1992, 7, 147-150.	1.2	27
98	Composition of the Essential Oils from Leaves and Flowers of <i>Achillea millefolium</i> L. ssp. <i>millefolium</i> . <i>Flavour and Fragrance Journal</i> , 1992, 7, 219-222.	1.2	45
99	Analysis of the Essential Oil of <i>Crithmum maritimum</i> L.. <i>Journal of Essential Oil Research</i> , 1991, 3, 313-316.	1.3	25