

David Shiers

List of Publications by Citations

Source: <https://exaly.com/author-pdf/8393115/david-shiers-publications-by-citations.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

31
papers

1,114
citations

10
h-index

33
g-index

36
ext. papers

1,550
ext. citations

7
avg, IF

3.72
L-index

#	Paper	IF	Citations
31	The Lancet Psychiatry Commission: a blueprint for protecting physical health in people with mental illness. <i>Lancet Psychiatry, the</i> , 2019 , 6, 675-712	23.3	411
30	Excess mortality in persons with severe mental disorders: a multilevel intervention framework and priorities for clinical practice, policy and research agendas. <i>World Psychiatry</i> , 2017 , 16, 30-40	14.4	296
29	Assessment and treatment of physical health problems among people with schizophrenia: national cross-sectional study. <i>British Journal of Psychiatry</i> , 2014 , 205, 473-7	5.4	84
28	Structured lifestyle education for people with schizophrenia, schizoaffective disorder and first-episode psychosis (STEPWISE): randomised controlled trial. <i>British Journal of Psychiatry</i> , 2019 , 214, 63-73	5.4	46
27	Antipsychotic drugs versus cognitive behavioural therapy versus a combination of both in people with psychosis: a randomised controlled pilot and feasibility study. <i>Lancet Psychiatry, the</i> , 2018 , 5, 411-423	23.3	35
26	Structured lifestyle education to support weight loss for people with schizophrenia, schizoaffective disorder and first episode psychosis: the STEPWISE RCT. <i>Health Technology Assessment</i> , 2018 , 22, 1-160	4.4	24
25	Antipsychotic medication versus psychological intervention versus a combination of both in adolescents with first-episode psychosis (MAPS): a multicentre, three-arm, randomised controlled pilot and feasibility study. <i>Lancet Psychiatry, the</i> , 2020 , 7, 788-800	23.3	23
24	STEPWISE - STRUCTURED lifestyle Education for People With SchizophrEnia: a study protocol for a randomised controlled trial. <i>Trials</i> , 2016 , 17, 475	2.8	10
23	STRUCTURED lifestyle education for people With SchizophrEnia (STEPWISE): mixed methods process evaluation of a group-based lifestyle education programme to support weight loss in people with schizophrenia. <i>BMC Psychiatry</i> , 2019 , 19, 358	4.2	9
22	Adherence to NICE guidance on lifestyle advice for people with schizophrenia: a survey. <i>BJPsych Bulletin</i> , 2017 , 41, 137-144	1.2	8
21	Early psychosis for the non-specialist doctor. <i>BMJ, The</i> , 2017 , 357, j4578	5.9	8
20	COVID-19 and UK family carers: policy implications. <i>Lancet Psychiatry, the</i> , 2021 , 8, 929-936	23.3	8
19	Improving the physical health of people with severe mental illness: Don't just screen, intervene. <i>InnovAiT</i> , 2019 , 12, 203-210	0	6
18	Study protocol for a randomised controlled trial of CBT vs antipsychotics vs both in 14-18-year-olds: Managing Adolescent first episode Psychosis: a feasibility study (MAPS). <i>Trials</i> , 2019 , 20, 395	2.8	6
17	Living with diabetes alongside a severe mental illness: A qualitative exploration with people with severe mental illness, family members and healthcare staff. <i>Diabetic Medicine</i> , 2021 , 38, e14562	3.5	6
16	Physical Health Problems in Psychosis: Is It Time to Consider the Views of Family Carers?. <i>Frontiers in Psychiatry</i> , 2018 , 9, 668	5	4
15	Exploring Severe Mental Illness and Diabetes: Protocol for a Longitudinal, Observational, and Qualitative Mixed Methods Study. <i>JMIR Research Protocols</i> , 2019 , 8, 13407	2	3

14	"I don't think I took her fears seriously": Exploring the experiences of family members of individuals at-risk of developing psychosis over 12 months. <i>Clinical Psychology and Psychotherapy</i> , 2020 , 27, 965-976 ^{2.9}	2
13	Applying economic models to estimate local economic benefits of improved coverage of early intervention for psychosis. <i>Microbial Biotechnology</i> , 2019 , 13, 1424-1430	3.3 2
12	Impact of severe mental illness on healthcare use and health outcomes for people with type 2 diabetes: a longitudinal observational study in England. <i>British Journal of General Practice</i> , 2021 , 71, e565-e573 ^{1.6}	2
11	Psychological intervention, antipsychotic medication or a combined treatment for adolescents with a first episode of psychosis: the MAPS feasibility three-arm RCT. <i>Health Technology Assessment</i> , 2021 , 25, 1-124	4.4 2
10	Exploring the facilitators, barriers, and strategies for self-management in adults living with severe mental illness, with and without long-term conditions: A qualitative evidence synthesis. <i>PLoS ONE</i> , 2021 , 16, e0258937	3.7 1
9	Identifying determinants of diabetes risk and outcomes for people with severe mental illness: a mixed-methods study. <i>Health Services and Delivery Research</i> , 2021 , 9, 1-194	1.5 1
8	Exploring determinants of self-management in adults with severe mental illness: a qualitative evidence synthesis. <i>BJPsych Open</i> , 2021 , 7, S13-S13	5 1
7	Still 'being bothered about Billy': managing the physical health of people with severe mental illness. <i>British Journal of General Practice</i> , 2021 , 71, 373-376	1.6 1
6	Healthcare resource use and costs for people with type 2 diabetes mellitus with and without severe mental illness in England: longitudinal matched-cohort study using the Clinical Practice Research Datalink.. <i>British Journal of Psychiatry</i> , 2021 , 1-8	5.4 1
5	The DIAMONDS intervention to support self-management of type 2 diabetes in people with severe mental illness: Study protocol for a single-group feasibility study. <i>SSM Mental Health</i> , 2022 , 2, 100086	0
4	Stand-alone cognitive behavioural therapy is not in clinical equipoise with antipsychotic treatment - Author's reply. <i>Lancet Psychiatry</i> , 2018 , 5, 540-541	23.3
3	Authors' reply. <i>British Journal of Psychiatry</i> , 2016 , 208, 398-9	5.4
2	Study protocol for the development and internal validation of Schizophrenia Prediction of Resistance to Treatment (SPIRIT): a clinical tool for predicting risk of treatment resistance to antipsychotics in first-episode schizophrenia.. <i>BMJ Open</i> , 2022 , 12, e056420	3
1	Multiple versus single risk behaviour interventions for people with severe mental illness: a network meta-analysis and qualitative synthesis 2022 , 10, 1-186	