
Jannice Schau Slettemeås

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8398475/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Emergence of AmpC-producing Escherichia coli in the broiler production chain in a country with a low antimicrobial usage profile. Veterinary Microbiology, 2014, 171, 315-320.	1.9	65
2	Integron, Plasmid and Host Strain Characteristics of Escherichia coli from Humans and Food Included in the Norwegian Antimicrobial Resistance Monitoring Programs. PLoS ONE, 2015, 10, e0128797.	2.5	42
3	Plasmid and Host Strain Characteristics of Escherichia coli Resistant to Extended-Spectrum Cephalosporins in the Norwegian Broiler Production. PLoS ONE, 2016, 11, e0154019.	2.5	37
4	Staphylococcus cornubiensis sp. nov., a member of the Staphylococcus intermedius Group (SIG). International Journal of Systematic and Evolutionary Microbiology, 2018, 68, 3404-3408.	1.7	31
5	Clonal spread of Escherichia coli resistant to cephalosporins and quinolones in the Nordic broiler production. Veterinary Microbiology, 2018, 213, 123-128.	1.9	22
6	Further Diversity of the <i>Staphylococcus Intermedius</i> Group and Heterogeneity in the <i>Mbo</i> I Restriction Site Used for <i>Staphylococcus Pseudintermedius</i> Species Identification. Journal of Veterinary Diagnostic Investigation, 2010, 22, 756-759.	1.1	19
7	Occurrence and characterization of quinolone resistant Escherichia coli from Norwegian turkey meat and complete sequence of an IncX1 plasmid encoding qnrS1. PLoS ONE, 2019, 14, e0212936.	2.5	18
8	Dissemination of Quinolone-Resistant Escherichia coli in the Norwegian Broiler and Pig Production Chains and Possible Persistence in the Broiler Production Environment. Applied and Environmental Microbiology, 2020, 86, .	3.1	16
9	Occurrence of quinolone resistant E. coli originating from different animal species in Norway. Veterinary Microbiology, 2018, 217, 25-31.	1.9	12
10	Comparative Genome Analyses of Wild Type- and Quinolone Resistant Escherichia coli Indicate Dissemination of QREC in the Norwegian Broiler Breeding Pyramid. Frontiers in Microbiology, 2020, 11, 938.	3.5	5
11	Actinobacillus pleuropneumoniae Eradication with Enrofloxacin May Lead to Dissemination and Long-Term Persistence of Quinolone Resistant Escherichia coli in Pig Herds. Antibiotics, 2020, 9, 910.	3.7	4
12	Complete Genome Sequences of 12 Quinolone-Resistant Escherichia coli Strains Containing qnrS1 Based on Hybrid Assemblies. Microbiology Resource Announcements, 2021, 10, .	0.6	1
13	Closed Genome Sequences of Providencia alcalifaciens Isolates from Dogs. Microbiology Resource Announcements, 2022, 11, e0095521.	0.6	1
14	Longitudinal Sampling Reveals Persistence of and Genetic Diversity in Extended-Spectrum Cephalosporin-Resistant Escherichia coli From Norwegian Broiler Production. Frontiers in Microbiology, 2021, 12, 795127.	3.5	1
15	A European multicenter evaluation study to investigate the performance on commercially available selective agar plates for the detection of carbapenemase producing Enterobacteriaceae. Journal of Microbiological Methods, 2022, 193, 106418.	1.6	0