

Miquel Quer

List of Publications by Citations

Source: <https://exaly.com/author-pdf/8408544/miquel-quer-publications-by-citations.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

188
papers

4,148
citations

33
h-index

57
g-index

217
ext. papers

4,957
ext. citations

2.9
avg, IF

4.99
L-index

#	Paper	IF	Citations
188	HPV Involvement in Head and Neck Cancers: Comprehensive Assessment of Biomarkers in 3680 Patients. <i>Journal of the National Cancer Institute</i> , 2016 , 108, djv403	9.7	441
187	Second neoplasm in patients with head and neck cancer. <i>Head and Neck</i> , 1999 , 21, 204-10	4.2	235
186	Proposal for revision of the European Laryngological Society classification of endoscopic cordectomies. <i>European Archives of Oto-Rhino-Laryngology</i> , 2007 , 264, 499-504	3.5	205
185	The role of type of tobacco and type of alcoholic beverage in oral carcinogenesis. <i>International Journal of Cancer</i> , 2004 , 108, 741-9	7.5	186
184	Distant metastases in head and neck cancer patients who achieved loco-regional control. <i>Head and Neck</i> , 2000 , 22, 680-6	4.2	177
183	An anatomical study of anastomoses between the laryngeal nerves. <i>Laryngoscope</i> , 1999 , 109, 983-7	3.6	131
182	Endoscopic supraglottic laryngectomy: a proposal for a classification by the working committee on nomenclature, European Laryngological Society. <i>European Archives of Oto-Rhino-Laryngology</i> , 2009 , 266, 993-8	3.5	75
181	Is the external laryngeal nerve an exclusively motor nerve? The cricothyroid connection branch. <i>Laryngoscope</i> , 2003 , 113, 525-9	3.6	73
180	Classification of parotidectomies: a proposal of the European Salivary Gland Society. <i>European Archives of Oto-Rhino-Laryngology</i> , 2016 , 273, 3307-12	3.5	70
179	Inclusion of extracapsular spread in the pTNM classification system: a proposal for patients with head and neck carcinoma. <i>JAMA Otolaryngology - Head and Neck Surgery</i> , 2013 , 139, 483-8	3.9	69
178	Influence of the persistence of tobacco and alcohol use in the appearance of second neoplasm in patients with a head and neck cancer. A case-control study. <i>Cancer Causes and Control</i> , 2009 , 20, 645-52	2.8	59
177	Pretreatment count of peripheral neutrophils, monocytes, and lymphocytes as independent prognostic factor in patients with head and neck cancer. <i>Head and Neck</i> , 2017 , 39, 219-226	4.2	53
176	Variability in nerve patterns of the adductor muscle group supplied by the recurrent laryngeal nerve. <i>Laryngoscope</i> , 2005 , 115, 358-62	3.6	52
175	Salvage surgery after locoregional failure in head and neck carcinoma patients treated with chemoradiotherapy. <i>European Archives of Oto-Rhino-Laryngology</i> , 2011 , 268, 295-301	3.5	50
174	Carotid blowout syndrome: modern trends in management. <i>Cancer Management and Research</i> , 2018 , 10, 5617-5628	3.6	50
173	Human papillomavirus as prognostic marker with rising prevalence in neck squamous cell carcinoma of unknown primary: A retrospective multicentre study. <i>European Journal of Cancer</i> , 2017 , 74, 73-81	7.5	47
172	European Laryngological Society: ELS recommendations for the follow-up of patients treated for laryngeal cancer. <i>European Archives of Oto-Rhino-Laryngology</i> , 2014 , 271, 2469-79	3.5	46

171	Enhanced cell migration and apoptosis resistance may underlie the association between high SERPINE1 expression and poor outcome in head and neck carcinoma patients. <i>Oncotarget</i> , 2015 , 6, 29016-33	3.3	45
170	Endoscopic laser surgery in the treatment of radiation failure of early laryngeal carcinoma. <i>Head and Neck</i> , 2000 , 22, 520-3	4.2	44
169	Variability of the nerve supply patterns of the human posterior cricoarytenoid muscle. <i>Laryngoscope</i> , 2003 , 113, 602-6	3.6	42
168	Results of salvage surgery for local or regional recurrence after larynx preservation with induction chemotherapy and radiotherapy. <i>Head and Neck</i> , 2001 , 23, 733-8	4.2	41
167	Hypothyroidism in patients treated with total laryngectomy. A multivariate study. <i>European Archives of Oto-Rhino-Laryngology</i> , 2002 , 259, 193-6	3.5	40
166	Prostaglandin E(2) pathway in head and neck squamous cell carcinoma. <i>Head and Neck</i> , 2008 , 30, 1175-81	4.2	39
165	Double positivity for HPV-DNA/p16 is the biomarker with strongest diagnostic accuracy and prognostic value for human papillomavirus related oropharyngeal cancer patients. <i>Oral Oncology</i> , 2018 , 78, 137-144	4.4	38
164	Head and Neck Cancer: A Review of the Impact of Treatment Delay on Outcome. <i>Advances in Therapy</i> , 2018 , 35, 153-160	4.1	38
163	Prevention of stomal recurrence. <i>Head and Neck</i> , 1996 , 18, 54-9	4.2	37
162	Ku70 predicts response and primary tumor recurrence after therapy in locally advanced head and neck cancer. <i>International Journal of Cancer</i> , 2008 , 123, 1068-79	7.5	36
161	Incidence and significance of clinically unsuspected thyroid tissue in lymph nodes found during neck dissection in head and neck carcinoma patients. <i>Laryngoscope</i> , 2005 , 115, 470-4	3.6	36
160	Selective dissection of levels II-III with intraoperative control of the upper and middle jugular nodes: a therapeutic option for the N0 neck. <i>Head and Neck</i> , 2001 , 23, 441-6	4.2	36
159	Surgical options in benign parotid tumors: a proposal for classification. <i>European Archives of Oto-Rhino-Laryngology</i> , 2017 , 274, 3825-3836	3.5	36
158	The effect of waiting time on local control and survival in head and neck carcinoma patients treated with radiotherapy. <i>Radiotherapy and Oncology</i> , 2003 , 66, 277-81	5.3	34
157	Management of recurrent head and neck cancer: variables related to salvage surgery. <i>European Archives of Oto-Rhino-Laryngology</i> , 2016 , 273, 4417-4424	3.5	34
156	Expression of IL-1 β correlates with distant metastasis in patients with head and neck squamous cell carcinoma. <i>Oncotarget</i> , 2015 , 6, 37398-409	3.3	32
155	Interaction between head and neck squamous cell carcinoma cells and fibroblasts in the biosynthesis of PGE2. <i>Journal of Lipid Research</i> , 2012 , 53, 630-42	6.3	31
154	Montgomery salivary bypass tube in the reconstruction of the hypopharynx. Cost-benefit study. <i>Annals of Otology, Rhinology and Laryngology</i> , 1999 , 108, 864-8	2.1	31

153	Management of the facial nerve in parotid cancer: preservation or resection and reconstruction. <i>European Archives of Oto-Rhino-Laryngology</i> , 2018 , 275, 2615-2626	3.5	31
152	Metachronous second primary tumours in the aerodigestive tract in patients with early stage head and neck squamous cell carcinomas. <i>European Archives of Oto-Rhino-Laryngology</i> , 2005 , 262, 905-9	3.5	30
151	Potential structures that could be confused with a nonrecurrent inferior laryngeal nerve: an anatomic study. <i>Laryngoscope</i> , 2008 , 118, 56-60	3.6	29
150	Initial treatment of the early stages (I, II) of supraglottic squamous cell carcinoma: partial laryngectomy versus radiotherapy. <i>European Archives of Oto-Rhino-Laryngology</i> , 2000 , 257, 512-6	3.5	28
149	Supracricoid laryngectomy as salvage surgery after failure of radiation therapy. <i>European Archives of Oto-Rhino-Laryngology</i> , 2007 , 264, 809-14	3.5	27
148	Contemporary management of primary parapharyngeal space tumors. <i>Head and Neck</i> , 2019 , 41, 522-535	4.2	27
147	Total or subtotal glossectomy with microsurgical reconstruction: functional and oncological results. <i>Microsurgery</i> , 2011 , 31, 517-23	2.1	26
146	Laryngeal carcinoma in patients without a history of tobacco and alcohol use. <i>Head and Neck</i> , 1997 , 19, 200-4	4.2	26
145	Pretreatment peripheral blood leukocytes are independent predictors of survival in oral cavity cancer. <i>Cancer</i> , 2020 , 126, 994-1003	6.4	26
144	Validation of the pathological classification of lymph node metastasis for head and neck tumors according to the 8th edition of the TNM Classification of Malignant Tumors. <i>Oral Oncology</i> , 2017 , 70, 29-33	4.4	25
143	The role of HPV on the risk of second primary neoplasia in patients with oropharyngeal carcinoma. <i>Oral Oncology</i> , 2017 , 64, 37-43	4.4	25
142	Selective neck dissection in surgically treated head and neck squamous cell carcinoma patients with a clinically positive neck: Systematic review. <i>European Journal of Surgical Oncology</i> , 2018 , 44, 395-403	3.6	25
141	Oropharyngeal and laryngeal sensory innervation in the pathophysiology of swallowing disorders and sensory stimulation treatments. <i>Annals of the New York Academy of Sciences</i> , 2016 , 1380, 104-120	6.5	25
140	Lymphadenopathy caused by Mycobacterium colombiense. <i>Journal of Clinical Microbiology</i> , 2008 , 46, 1885-7	9.7	25
139	Polymorphous adenocarcinoma of the salivary glands: reappraisal and update. <i>European Archives of Oto-Rhino-Laryngology</i> , 2018 , 275, 1681-1695	3.5	25
138	Analysis of the Clinical Relevance of Histological Classification of Benign Epithelial Salivary Gland Tumours. <i>Advances in Therapy</i> , 2019 , 36, 1950-1974	4.1	24
137	Prognostic and predictive factors in recurrent and/or metastatic head and neck squamous cell carcinoma: A review of the literature. <i>Critical Reviews in Oncology/Hematology</i> , 2019 , 137, 84-91	7	24
136	Gene expression signatures and molecular markers associated with clinical outcome in locally advanced head and neck carcinoma. <i>Carcinogenesis</i> , 2012 , 33, 1707-16	4.6	24

135	Function preservation in stage III squamous laryngeal carcinoma: results with an induction chemotherapy protocol. <i>Laryngoscope</i> , 1995 , 105, 822-6	3.6	23
134	Second, third, and fourth head and neck tumors. A progressive decrease in survival. <i>Head and Neck</i> , 2012 , 34, 1716-9	4.2	22
133	Primary tracheoesophageal puncture vs esophageal speech. <i>JAMA Otolaryngology</i> , 1992 , 118, 188-90		22
132	Results of an organ preservation protocol with induction chemotherapy and radiotherapy in patients with locally advanced pyriform sinus carcinoma. <i>European Archives of Oto-Rhino-Laryngology</i> , 2002 , 259, 32-6	3.5	21
131	A Novel Role For Nanog As An Early Cancer Risk Marker In Patients With Laryngeal Precancerous Lesions. <i>Scientific Reports</i> , 2017 , 7, 11110	4.9	20
130	Can cure be achieved in patients with head and neck carcinomas? The problem of second neoplasm. <i>Expert Review of Anticancer Therapy</i> , 2001 , 1, 125-33	3.5	20
129	The risk of second primary tumors in head and neck cancer: A systematic review. <i>Head and Neck</i> , 2020 , 42, 456-466	4.2	20
128	Prognostic role of MMP-9 expression in head and neck carcinoma patients treated with radiotherapy or chemoradiotherapy. <i>Oral Oncology</i> , 2013 , 49, 322-5	4.4	19
127	Pharyngocutaneous fistula after total laryngectomy: multivariate analysis of risk factors and a severity-based classification proposal. <i>European Archives of Oto-Rhino-Laryngology</i> , 2019 , 276, 143-151	3.5	18
126	Postoperative staging of the neck dissection using extracapsular spread and lymph node ratio as prognostic factors in HPV-negative head and neck squamous cell carcinoma patients. <i>Oral Oncology</i> , 2018 , 77, 37-42	4.4	17
125	Influence of age on laryngeal carcinoma. <i>Annals of Otolaryngology, Rhinology and Laryngology</i> , 1998 , 107, 164-9	2.1	17
124	Prognostic capacity of Systemic Inflammation Response Index (SIRI) in patients with head and neck squamous cell carcinoma. <i>Head and Neck</i> , 2020 , 42, 336-343	4.2	16
123	Identification of somatic VHL gene mutations in sporadic head and neck paragangliomas in association with activation of the HIF-1/ miR-210 signaling pathway. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013 , 98, E1661-6	5.6	15
122	Foramen thyroideum: a comparative study in embryos, fetuses, and adults. <i>Laryngoscope</i> , 1997 , 107, 1146-50	3.6	15
121	External validation of a risk group defined by recursive partitioning analysis in patients with head and neck carcinoma treated with surgery and postoperative radiotherapy. <i>Head and Neck</i> , 2007 , 29, 815-21	4.2	15
120	How much does it cost to preserve a larynx? An economic study. <i>European Archives of Oto-Rhino-Laryngology</i> , 2000 , 257, 72-6	3.5	15
119	Salvage surgery after local recurrence in patients with head and neck carcinoma treated with chemoradiotherapy or bioradiotherapy. <i>Auris Nasus Larynx</i> , 2015 , 42, 145-9	2.2	14
118	Results of an organ preservation protocol with induction chemotherapy and radiotherapy in patients with locally advanced laryngeal carcinoma. <i>European Archives of Oto-Rhino-Laryngology</i> , 2005 , 262, 93-8	3.5	14

117	Surgical management of intrathoracic goitres. <i>European Archives of Oto-Rhino-Laryngology</i> , 2019 , 276, 305-314	3.5	14
116	How phenotype guides management of non-conventional squamous cell carcinomas of the larynx?. <i>European Archives of Oto-Rhino-Laryngology</i> , 2017 , 274, 2709-2726	3.5	13
115	CKMT1 and NCOA1 expression as a predictor of clinical outcome in patients with advanced-stage head and neck squamous cell carcinoma. <i>Head and Neck</i> , 2016 , 38 Suppl 1, E1392-403	4.2	13
114	Epidemiologic characteristics of squamous head and neck carcinoma patients. Results of a hospital register. <i>Acta Otorrinolaringológica Española</i> , 2019 , 70, 272-278	0.9	13
113	Expression of the CXCL12/CXCR4 chemokine axis predicts regional control in head and neck squamous cell carcinoma. <i>European Archives of Oto-Rhino-Laryngology</i> , 2016 , 273, 4525-4533	3.5	12
112	Prostacyclin-synthase expression in head and neck carcinoma patients and its prognostic value in the response to radiotherapy. <i>Journal of Pathology</i> , 2015 , 235, 125-35	9.4	12
111	Stomal recurrence in head and neck cancer patients with temporary tracheostomy. <i>Auris Nasus Larynx</i> , 2014 , 41, 467-70	2.2	12
110	Delayed carotid blow-out syndrome: a new complication of chemoradiotherapy treatment in pharyngolaryngeal carcinoma. <i>Journal of Laryngology and Otology</i> , 2012 , 126, 1189-91	1.8	12
109	Mutation analysis of the SDHD gene in four kindreds with familial paraganglioma: description of one novel germline mutation. <i>Diagnostic Molecular Pathology</i> , 2005 , 14, 109-14		12
108	Effect of serpinE1 overexpression on the primary tumor and lymph node, and lung metastases in head and neck squamous cell carcinoma. <i>Head and Neck</i> , 2019 , 41, 429-439	4.2	12
107	Immunohistochemical Expression of Cortactin and Focal Adhesion Kinase Predicts Recurrence Risk and Laryngeal Cancer Risk Beyond Histologic Grading. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2018 , 27, 805-813	4	11
106	Development and (pre-) clinical assessment of a novel surgical tool for primary and secondary tracheoesophageal puncture with immediate voice prosthesis insertion, the Provox Vega Puncture Set. <i>European Archives of Oto-Rhino-Laryngology</i> , 2013 , 270, 255-62	3.5	11
105	The prognostic value of pretreatment platelet count in patients with head and neck squamous cell carcinoma. <i>Auris Nasus Larynx</i> , 2017 , 44, 313-318	2.2	11
104	High RAB25 expression is associated with good clinical outcome in patients with locally advanced head and neck squamous cell carcinoma. <i>Cancer Medicine</i> , 2013 , 2, 950-63	4.8	11
103	Risk of third and fourth tumors in patients with head and neck cancer. <i>Head and Neck</i> , 2010 , 32, 1467-72	4.2	11
102	Detection of Distant Metastases in Head and Neck Cancer: Changing Landscape. <i>Advances in Therapy</i> , 2018 , 35, 161-172	4.1	10
101	Prognostic significance of extranodal extension in head and neck squamous cell carcinoma cN0 patients with occult metastatic neck nodes. <i>Acta Otorrinolaringológica Española</i> , 2018 , 69, 156-164	0.9	10
100	Resultados de un protocolo de preservaci3n de 3gano en pacientes con tumores localmente avanzados de la laringe. <i>Acta Otorrinolaringológica Española</i> , 2009 , 60, 176-185	0.9	10

99	Prognostic role of intraparotid lymph node metastasis in primary parotid cancer: Systematic review. <i>Head and Neck</i> , 2021 , 43, 997-1008	4.2	10
98	Impact of total laryngectomy on return to work. <i>Acta Otorrinolaringológica Española</i> , 2018 , 69, 74-79	0.9	9
97	Overexpression of the nuclear factor-kappa B (p65) in association with local failure in patients with head and neck carcinoma undergoing radiotherapy or chemoradiotherapy. <i>Head and Neck</i> , 2013 , 35, 370-375	4.2	8
96	Parotidectomies in benign parotid tumours: "Sant Pau" surgical extension classification. <i>Acta Otorrinolaringológica Española</i> , 2010 , 61, 1-5	0.9	8
95	Modalidades de preservación de laringe en carcinomas de laringe e hipofaringe. <i>Acta Otorrinolaringológica Española</i> , 2007 , 58, 476-482	0.9	8
94	Comparison of the Radiation Therapy Oncology Group recursive partitioning classification and Union Internationale Contre le Cancer TNM classification for patients with head and neck carcinoma. <i>Head and Neck</i> , 2005 , 27, 248-57	4.2	8
93	Relationship between response to induction chemotherapy and disease control in patients with advanced laryngeal carcinoma included in an organ preservation protocol. <i>European Archives of Oto-Rhino-Laryngology</i> , 2017 , 274, 2581-2587	3.5	8
92	External validation of sTWEAK as a prognostic noninvasive biomarker for head and neck squamous cell carcinoma. <i>Head and Neck</i> , 2016 , 38 Suppl 1, E1358-63	4.2	8
91	Management of the Neck in Well-Differentiated Thyroid Cancer. <i>Current Oncology Reports</i> , 2020 , 23, 1	6.3	8
90	Prognostic significance of extracapsular spread in isolated neck recurrences in head and neck squamous cell carcinoma patients. <i>European Archives of Oto-Rhino-Laryngology</i> , 2017 , 274, 527-533	3.5	7
89	Parotid incidentaloma identified by positron emission/computed tomography: when to consider diagnoses other than warthin tumor. <i>International Archives of Otorhinolaryngology</i> , 2015 , 19, 112-5	1.5	7
88	Reconstruction of oral cavity defects with FAMM (facial artery musculomucosal) flaps. Our experience. <i>Acta Otorrinolaringológica Española</i> , 2015 , 66, 275-80	0.9	7
87	Long-Term Hearing Outcomes following Stapedotomy in Patients with Otosclerosis and Preoperative Small Air-Bone Gap. <i>Audiology and Neuro-Otology</i> , 2017 , 22, 350-355	2.2	7
86	Ceratocricoid muscle: an embryological and anatomical study. <i>Clinical Anatomy</i> , 2009 , 22, 463-70	2.5	7
85	Recurrent laryngeal squamous cell carcinoma: rTNM versus composite laryngeal recurrence staging system. Proposal for a modification of the CLRSS to improve patient classification. <i>Head and Neck</i> , 2008 , 30, 939-45	4.2	7
84	Treatment of neck nodes after induction chemotherapy in patients with primary advanced tumours. <i>European Archives of Oto-Rhino-Laryngology</i> , 2000 , 257, 521-5	3.5	7
83	Competing mortality in oropharyngeal carcinoma according to human papillomavirus status. <i>Head and Neck</i> , 2019 , 41, 1328-1334	4.2	7
82	Elective neck dissection during salvage surgery after radiotherapy in patients with head and neck squamous cell carcinoma. <i>Acta Otorhinolaryngologica Italica</i> , 2018 , 38, 86-93	2.8	7

81	Risk of onset of second neoplasms and successive neoplasms in patients with a head and neck index tumour. <i>Acta Otorrinolaringológica Española</i> , 2020 , 71, 9-15	0.9	6
80	Elective treatment of the neck for second primary tumors of the head and neck. <i>European Archives of Oto-Rhino-Laryngology</i> , 2014 , 271, 1187-90	3.5	6
79	Blood transfusions in laryngeal cancer: effect on prognosis. <i>Head and Neck</i> , 1996 , 18, 218-24	4.2	6
78	Repair of post-laryngectomy pharyngocutaneous fistulas using a pectoralis major flap. <i>Brazilian Journal of Otorhinolaryngology</i> , 2019 , 85, 351-356	1.6	6
77	Weighted lymph node ratio: New tool in the assessment of postoperative staging of the neck dissection in HPV-negative head and neck squamous cell carcinoma patients. <i>Head and Neck</i> , 2020 , 42, 2912-2919	4.2	5
76	Histologically negative specimens after induction therapy: frequency and impact on survival. <i>Head and Neck</i> , 2000 , 22, 808-13	4.2	5
75	The combined use of EFS, GPX2, and SPRR1A expression could distinguish favorable from poor clinical outcome among epithelial-like head and neck carcinoma subtypes. <i>Head and Neck</i> , 2019 , 41, 1830-1845	4.2	5
74	European Laryngological Society position paper on laryngeal dysplasia Part II: diagnosis, treatment, and follow-up. <i>European Archives of Oto-Rhino-Laryngology</i> , 2021 , 278, 1723-1732	3.5	5
73	Patients with locally advanced hypopharyngeal carcinoma. Results over a 30-year period. <i>Acta Otorrinolaringológica Española</i> , 2019 , 70, 315-326	0.9	4
72	Salvage surgery in head and neck cancer: External validation of predictors of disease-specific survival. <i>Oral Oncology</i> , 2020 , 109, 104876	4.4	4
71	Comparison of chi-squared automatic interaction detection classification trees vs TNM classification for patients with head and neck squamous cell carcinoma. <i>JAMA Otolaryngology</i> , 2012 , 138, 272-9		4
70	Oncological results of salvage laryngectomy in patients with laryngeal carcinoma. <i>Acta Otorrinolaringológica Española</i> , 2020 , 71, 70-77	0.9	4
69	Does age influence disease-specific survival in patients with squamous cell carcinomas of the head and neck?. <i>Journal of Surgical Oncology</i> , 2020 , 121, 1058-1066	2.8	3
68	CXCR4-targeted nanotoxins induce GSDME-dependent pyroptosis in head and neck squamous cell carcinoma.. <i>Journal of Experimental and Clinical Cancer Research</i> , 2022 , 41, 49	12.8	3
67	Self-assembling protein nanocarrier for selective delivery of cytotoxic polypeptides to CXCR4+ head and neck squamous cell carcinoma tumors. <i>Acta Pharmaceutica Sinica B</i> , 2021 ,	15.5	3
66	Prognostic value of CD45 transcriptional expression in head and neck cancer. <i>European Archives of Oto-Rhino-Laryngology</i> , 2018 , 275, 225-232	3.5	3
65	How Phenotype Guides Management of the Most Common Malignant Salivary Neoplasms of the Larynx?. <i>Advances in Therapy</i> , 2017 , 34, 813-825	4.1	2
64	Risk of Onset of Second Neoplasms and Successive Neoplasms in Patients With a Head and Neck Index Tumour. <i>Acta Otorrinolaringologica (English Edition)</i> , 2020 , 71, 9-15	0.1	2

63	Impact of Total Laryngectomy on Return to Work. <i>Acta Otorrinolaringologica (English Edition)</i> , 2018 , 69, 74-79	0.1	2
62	Indications and results of extended total laryngectomy with en-bloc resection of overlying cervical skin. <i>European Archives of Oto-Rhino-Laryngology</i> , 2019 , 276, 3179-3184	3.5	2
61	Results of an organ conservation protocol in patients with locally advanced laryngeal tumours. <i>Acta Otorrinolaringologica (English Edition)</i> , 2009 , 60, 176-185	0.1	2
60	Prognostic value of the nodal yield in elective neck dissections in patients with head and neck carcinomas. <i>European Archives of Oto-Rhino-Laryngology</i> , 2021 , 1	3.5	2
59	Salvage carbon dioxide transoral laser microsurgery for laryngeal cancer after (chemo)radiotherapy: a European Laryngological Society consensus statement. <i>European Archives of Oto-Rhino-Laryngology</i> , 2021 , 278, 4373-4381	3.5	2
58	Prevention of stomal recurrence 1996 , 18, 54		2
57	Voice Outcome After Carbon Dioxide Transoral Laser Microsurgery for Glottic Cancer According to the European Laryngological Society Classification of Cordectomy Types - A Systematic Review.. <i>Journal of Voice</i> , 2022 ,	1.9	2
56	Prognostic Significance of Extranodal Extension in Head and Neck Squamous Cell Carcinoma cN0 Patients With Occult Metastatic Neck Nodes. <i>Acta Otorrinolaringologica (English Edition)</i> , 2018 , 69, 156-164	0.1	1
55	Aetiology and Treatment of Vocal Fold Paralysis: Retrospective Study of 108 Patients. <i>Acta Otorrinolaringologica (English Edition)</i> , 2014 , 65, 225-230	0.1	1
54	Parotidectomies in benign parotid tumours: Bant Paußurgical extension classification. <i>Acta Otorrinolaringologica (English Edition)</i> , 2010 , 61, 1-5	0.1	1
53	Risk of distant metastases in head and neck carcinoma patients and myeloperoxidase (MPO) expression.. <i>Journal of Clinical Oncology</i> , 2016 , 34, 6067-6067	2.2	1
52	Prognostic significance of albumin levels prior to treatment in patients with head and neck squamous cell carcinoma. <i>Acta Otorrinolaringologica Española</i> , 2020 , 71, 204-211	0.9	1
51	The aspartate aminotransaminase/alanine aminotransaminase (De Ritis) ratio predicts sensitivity to radiotherapy in head and neck carcinoma patients. <i>Head and Neck</i> , 2021 , 43, 2091-2100	4.2	1
50	Trends in disease-specific survival of head and neck squamous cell carcinoma patients treated in a single institution over a 30-year period. <i>Oral Oncology</i> , 2021 , 115, 105184	4.4	1
49	Predictive value of transcriptional expression of Krüppel-like factor-6 (KLF6) in head and neck carcinoma patients treated with radiotherapy. <i>Clinical and Translational Oncology</i> , 2021 , 23, 2507-2512	3.6	1
48	Reply to the letter to the editor concerning: 'Classification of parotidectomies: a proposal of the European Salivary Gland Society'. <i>European Archives of Oto-Rhino-Laryngology</i> , 2016 , 273, 3453	3.5	1
47	European Laryngological Society position paper on laryngeal dysplasia Part I: aetiology and pathological classification. <i>European Archives of Oto-Rhino-Laryngology</i> , 2021 , 278, 1717-1722	3.5	1
46	Prognostic role of extracapsular spread in planned neck dissection after chemoradiotherapy. <i>Head and Neck</i> , 2018 , 40, 2514-2520	4.2	1

45	Absence of disruptive TP53 mutations in high-risk human papillomavirus-driven neck squamous cell carcinoma of unknown primary. <i>Head and Neck</i> , 2019 , 41, 3833-3841	4.2	o
44	Trasplante de la laringe: ¿una opción terapéutica para el siglo XXI? Revisión de la literatura. <i>Acta Otorrinolaringológica Española</i> , 2008 , 59, 127-138	0.9	o
43	Organ Preservation in Laryngeal and Hypopharyngeal Cancer. <i>Acta Otorrinolaringologica (English Edition)</i> , 2007 , 58, 476-482	0.1	o
42	Predictive capacity of IL-8 expression in head and neck squamous carcinoma patients treated with radiotherapy or chemoradiotherapy. <i>Acta Otorrinolaringológica Española</i> , 2020 , 72, 337-337	0.9	o
41	Oral cavity colon adenocarcinoma metastases: case report with surgical approach and review of more than 30 years literature. <i>Oral and Maxillofacial Surgery</i> , 2021 , 25, 99-101	1.6	o
40	Intraoperative monitoring of the facial nerve during parotid gland surgery in Otolaryngology services - Head and Neck Surgery. <i>Acta Otorrinolaringológica Española</i> , 2021 , 72, 158-163	0.9	o
39	Prognostic capacity of the weighted lymph node ratio in head and neck squamous cell carcinoma patients treated with salvage neck dissection. <i>European Archives of Oto-Rhino-Laryngology</i> , 2021 , 278, 4005-4010	3.5	o
38	Analysis of Specific Survival and Local Control Through a Recursive Partitioning Analysis in Patients With Head and Neck Carcinoma. <i>Acta Otorrinolaringologica (English Edition)</i> , 2020 , 71, 131-139	0.1	
37	Oncological Results of Salvage Laryngectomy in Patients With Laryngeal Carcinoma. <i>Acta Otorrinolaringologica (English Edition)</i> , 2020 , 71, 70-77	0.1	
36	Epidemiologic Characteristics of Squamous Head and Neck Carcinoma Patients. Results of a Hospital Register. <i>Acta Otorrinolaringologica (English Edition)</i> , 2019 , 70, 272-278	0.1	
35	Pharyngocutaneous fistula in irradiated patients: systematic review and our experience.. <i>Journal of Laryngology and Otology</i> , 1-23	1.8	
34	Relationship between the transcriptional expression of PIM1 and local control in patients with head and neck squamous cell carcinomas treated with radiotherapy.. <i>European Archives of Oto-Rhino-Laryngology</i> , 2022 , 1	3.5	
33	Prognostic capacity of the lymph node ratio in squamous cell carcinomas of the head and neck. <i>Acta Otorrinolaringológica Española</i> , 2020 , 71, 265-274	0.9	
32	Differential Characteristics of Patients With Squamous Cell Carcinoma of the Head and Neck According to Gender. <i>Acta Otorrinolaringologica (English Edition)</i> , 2021 , 72, 359-369	0.1	
31	Predictive capacity of IL-8 expression in head and neck squamous carcinoma patients treated with radiotherapy or chemoradiotherapy. <i>Acta Otorrinolaringologica (English Edition)</i> , 2021 , 72, 337-343	0.1	
30	Low skeletal muscle mass assessed directly from the 3rd cervical vertebra can predict pharyngocutaneous fistula risk after total laryngectomy in the male population. <i>European Archives of Oto-Rhino-Laryngology</i> , 2021 , 279, 853	3.5	
29	Distant metastases in squamous-cell head and neck carcinoma (SHNC) patients who achieved loco-regional control (LRC). <i>Journal of Clinical Oncology</i> , 2004 , 22, 5519-5519	2.2	
28	Smoke and alcohol consumption as a risk factors in the development of second primary neoplasms (SPN) in head & neck cancer (HNC) patients. A case-control study. <i>Journal of Clinical Oncology</i> , 2004 , 22, 5582-5582	2.2	

27	A proposal of pathological classification (pTNM) of patients with head and neck cancer according to capsular rupture (pN- RC). <i>Journal of Clinical Oncology</i> , 2007 , 25, 16525-16525	2.2
26	Risk of second primary neoplasia in patients with oropharyngeal carcinoma: Role of HPV status in the outcome.. <i>Journal of Clinical Oncology</i> , 2019 , 37, e17544-e17544	2.2
25	Analysis of specific survival and local control through a recursive partitioning analysis in patients with head and neck carcinoma. <i>Acta Otorrinolaringológica Española</i> , 2020 , 71, 131-139	0.9
24	Prognostic value of nodal involvement in patients with oropharyngeal carcinoma according to the HPV status. <i>Acta Otorrinolaringológica Española</i> , 2020 , 71, 212-218	0.9
23	Expression of interleukin-1[IL1-] and risk of distant metastases (DM) in head and neck squamous cell carcinoma (HNSCC).. <i>Journal of Clinical Oncology</i> , 2014 , 32, 6078-6078	2.2
22	Expression of the CXCL12/CXCR4 chemokine axis as predictor of regional control in head and neck squamous cell carcinoma (HNSCC).. <i>Journal of Clinical Oncology</i> , 2015 , 33, e17093-e17093	2.2
21	Validation of the pathological classification of lymph node metastasis for head and neck tumors (HNSCC) according to the 8th edition of the TNM Classification of Malignant Tumors (8th TNM).. <i>Journal of Clinical Oncology</i> , 2017 , 35, 6014-6014	2.2
20	Total Laryngectomy 2009 , 229-243	
19	Progressive increasing in the risk of second and subsequent malignant tumors in patients with a head and neck cancer: A validation study with SEER data.. <i>Journal of Clinical Oncology</i> , 2012 , 30, 5595-5595	2.2
18	Gene expression as a predictor of radiotherapy or chemoradiotherapy response in head and neck squamous cell carcinoma patients: A molecular approach.. <i>Journal of Clinical Oncology</i> , 2012 , 30, 5590-5590	2.2
17	Prognostic Value of Nodal Involvement in Patients With Oropharyngeal Carcinoma According to the HPV Status. <i>Acta Otorrinolaringologica (English Edition)</i> , 2020 , 71, 212-218	0.1
16	Prognostic Significance of Albumin Levels Prior to Treatment in Patients With Head and Neck Squamous Cell Carcinoma. <i>Acta Otorrinolaringologica (English Edition)</i> , 2020 , 71, 204-211	0.1
15	Prognostic Capacity of the Lymph Node Ratio in Squamous Cell Carcinomas of the Head and Neck. <i>Acta Otorrinolaringologica (English Edition)</i> , 2020 , 71, 265-274	0.1
14	Oncological results of salvage treatment in patients with hypopharyngeal carcinoma. <i>Acta Otorrinolaringológica Española</i> , 2021 , 72, 85-91	0.9
13	Oncological Results of Salvage Treatment in Patients With Hypopharyngeal Carcinoma. <i>Acta Otorrinolaringologica (English Edition)</i> , 2021 , 72, 85-91	0.1
12	Organ preservation in patients with advanced laryngeal tumours. Results of induction chemotherapy versus chemoradiotherapy in actual clinical practice. <i>Acta Otorrinolaringológica Española</i> , 2021 , 72, 143-151	0.9
11	Organ Preservation in Patients With Advanced Laryngeal Tumours. Results of Induction Chemotherapy Versus Chemoradiotherapy in Actual Clinical Practice. <i>Acta Otorrinolaringologica (English Edition)</i> , 2021 , 72, 143-151	0.1
10	Pacientes fumadores de cigarrillos con carcinomas escamosos de cabeza y cuello. Características epidemiológicas y pronósticas diferenciales. <i>Acta Otorrinolaringológica Española</i> , 2021 , 72, 222-229	0.9

9	Head and neck squamous cell carcinoma in cigar smokers. Distinctive epidemiological and prognostic characteristics. <i>Acta Otorrinolaringologica (English Edition)</i> , 2021 , 72, 222-229	0.1
8	Reply to the letter "A proposal for a level for parapharyngeal extension of parotid gland". <i>European Archives of Oto-Rhino-Laryngology</i> , 2016 , 273, 3457	3.5
7	Patients With Locally Advanced Hypopharyngeal Carcinoma. Results Over a 30-year Period. <i>Acta Otorrinolaringologica (English Edition)</i> , 2019 , 70, 315-326	0.1
6	Organ preservation after treatment with induction chemotherapy in patients with locally advanced carcinomas (T3-T4) of oral cavity and oropharynx. <i>Acta Otorrinolaringologica Española</i> , 2021 , 72, 27-36	0.9
5	Organ Preservation After Treatment With Induction Chemotherapy in Patients With Locally Advanced Carcinomas (T3-T4) of Oral Cavity and Oropharynx. <i>Acta Otorrinolaringologica (English Edition)</i> , 2021 , 72, 27-36	0.1
4	Differential characteristics of patients with squamous cell carcinoma of the head and neck according to gender. <i>Acta Otorrinolaringologica Española</i> , 2021 , 72, 359-359	0.9
3	The Selective Role of Open and Endoscopic Approaches for Sinonasal Malignant Tumours.. <i>Advances in Therapy</i> , 2022 , 1	4.1
2	Distant metastases in squamous-cell head and neck carcinoma (SHNC) patients who achieved loco-regional control (LRC). <i>Journal of Clinical Oncology</i> , 2004 , 22, 5519-5519	2.2
1	Smoke and alcohol consumption as a risk factors in the development of second primary neoplasms (SPN) in head & neck cancer (HNC) patients. A case-control study. <i>Journal of Clinical Oncology</i> , 2004 , 22, 5582-5582	2.2