

Miquel Quer

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8408544/miquel-quer-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

188
papers

4,148
citations

33
h-index

57
g-index

217
ext. papers

4,957
ext. citations

2.9
avg, IF

4.99
L-index

#	Paper	IF	Citations
188	Relationship between the transcriptional expression of PIM1 and local control in patients with head and neck squamous cell carcinomas treated with radiotherapy.. <i>European Archives of Oto-Rhino-Laryngology</i> , 2022 , 1	3.5	
187	CXCR4-targeted nanotoxins induce GSDME-dependent pyroptosis in head and neck squamous cell carcinoma.. <i>Journal of Experimental and Clinical Cancer Research</i> , 2022 , 41, 49	12.8	3
186	The Selective Role of Open and Endoscopic Approaches for Sinonasal Malignant Tumours.. <i>Advances in Therapy</i> , 2022 , 1	4.1	
185	Voice Outcome After Carbon Dioxide Transoral Laser Microsurgery for Glottic Cancer According to the European Laryngological Society Classification of Cordectomy Types - A Systematic Review.. <i>Journal of Voice</i> , 2022 ,	1.9	2
184	Differential Characteristics of Patients With Squamous Cell Carcinoma of the Head and Neck According to Gender. <i>Acta Otorrinolaringologica (English Edition)</i> , 2021 , 72, 359-369	0.1	
183	Predictive capacity of IL-8 expression in head and neck squamous carcinoma patients treated with radiotherapy or chemoradiotherapy. <i>Acta Otorrinolaringologica (English Edition)</i> , 2021 , 72, 337-343	0.1	
182	Self-assembling protein nanocarrier for selective delivery of cytotoxic polypeptides to CXCR4+ head and neck squamous cell carcinoma tumors. <i>Acta Pharmaceutica Sinica B</i> , 2021 ,	15.5	3
181	Low skeletal muscle mass assessed directly from the 3rd cervical vertebra can predict pharyngocutaneous fistula risk after total laryngectomy in the male population. <i>European Archives of Oto-Rhino-Laryngology</i> , 2021 , 279, 853	3.5	
180	Oncological results of salvage treatment in patients with hypopharyngeal carcinoma. <i>Acta Otorrinolaringologica Española</i> , 2021 , 72, 85-91	0.9	
179	The aspartate aminotransaminase/alanine aminotransaminase (De Ritis) ratio predicts sensitivity to radiotherapy in head and neck carcinoma patients. <i>Head and Neck</i> , 2021 , 43, 2091-2100	4.2	1
178	Oncological Results of Salvage Treatment in Patients With Hypopharyngeal Carcinoma. <i>Acta Otorrinolaringologica (English Edition)</i> , 2021 , 72, 85-91	0.1	
177	Trends in disease-specific survival of head and neck squamous cell carcinoma patients treated in a single institution over a 30-year period. <i>Oral Oncology</i> , 2021 , 115, 105184	4.4	1
176	Prognostic value of the nodal yield in elective neck dissections in patients with head and neck carcinomas. <i>European Archives of Oto-Rhino-Laryngology</i> , 2021 , 1	3.5	2
175	Organ preservation in patients with advanced laryngeal tumours. Results of induction chemotherapy versus chemoradiotherapy in actual clinical practice. <i>Acta Otorrinolaringologica Española</i> , 2021 , 72, 143-151	0.9	
174	Organ Preservation in Patients With Advanced Laryngeal Tumours. Results of Induction Chemotherapy Versus Chemoradiotherapy in Actual Clinical Practice. <i>Acta Otorrinolaringologica (English Edition)</i> , 2021 , 72, 143-151	0.1	
173	Predictive value of transcriptional expression of Krüppel-like factor-6 (KLF6) in head and neck carcinoma patients treated with radiotherapy. <i>Clinical and Translational Oncology</i> , 2021 , 23, 2507-2512	3.6	1
172	Pacientes fumadores de cigarrillos con carcinomas escamosos de cabeza y cuello. Características epidemiológicas y pronósticas diferenciales. <i>Acta Otorrinolaringologica Española</i> , 2021 , 72, 222-229	0.9	

171	Head and neck squamous cell carcinoma in cigar smokers. Distinctive epidemiological and prognostic characteristics. <i>Acta Otorrinolaringologica (English Edition)</i> , 2021 , 72, 222-229	0.1	
170	Oral cavity colon adenocarcinoma metastases: case report with surgical approach and review of more than 30 years literature. <i>Oral and Maxillofacial Surgery</i> , 2021 , 25, 99-101	1.6	0
169	European Laryngological Society position paper on laryngeal dysplasia Part II: diagnosis, treatment, and follow-up. <i>European Archives of Oto-Rhino-Laryngology</i> , 2021 , 278, 1723-1732	3.5	5
168	European Laryngological Society position paper on laryngeal dysplasia Part I: aetiology and pathological classification. <i>European Archives of Oto-Rhino-Laryngology</i> , 2021 , 278, 1717-1722	3.5	1
167	Intraoperative monitoring of the facial nerve during parotid gland surgery in Otolaryngology services - Head and Neck Surgery. <i>Acta Otorrinolaringologica Española</i> , 2021 , 72, 158-163	0.9	0
166	Prognostic role of intraparotid lymph node metastasis in primary parotid cancer: Systematic review. <i>Head and Neck</i> , 2021 , 43, 997-1008	4.2	10
165	Prognostic capacity of the weighted lymph node ratio in head and neck squamous cell carcinoma patients treated with salvage neck dissection. <i>European Archives of Oto-Rhino-Laryngology</i> , 2021 , 278, 4005-4010	3.5	0
164	Organ preservation after treatment with induction chemotherapy in patients with locally advanced carcinomas (T3-T4) of oral cavity and oropharynx. <i>Acta Otorrinolaringologica Española</i> , 2021 , 72, 27-36	0.9	
163	Organ Preservation After Treatment With Induction Chemotherapy in Patients With Locally Advanced Carcinomas (T3-T4) of Oral Cavity and Oropharynx. <i>Acta Otorrinolaringologica (English Edition)</i> , 2021 , 72, 27-36	0.1	
162	Differential characteristics of patients with squamous cell carcinoma of the head and neck according to gender. <i>Acta Otorrinolaringologica Española</i> , 2021 , 72, 359-359	0.9	
161	Salvage carbon dioxide transoral laser microsurgery for laryngeal cancer after (chemo)radiotherapy: a European Laryngological Society consensus statement. <i>European Archives of Oto-Rhino-Laryngology</i> , 2021 , 278, 4373-4381	3.5	2
160	Risk of Onset of Second Neoplasms and Successive Neoplasms in Patients With a Head and Neck Index Tumour. <i>Acta Otorrinolaringologica (English Edition)</i> , 2020 , 71, 9-15	0.1	2
159	Analysis of Specific Survival and Local Control Through a Recursive Partitioning Analysis in Patients With Head and Neck Carcinoma. <i>Acta Otorrinolaringologica (English Edition)</i> , 2020 , 71, 131-139	0.1	
158	Oncological Results of Salvage Laryngectomy in Patients With Laryngeal Carcinoma. <i>Acta Otorrinolaringologica (English Edition)</i> , 2020 , 71, 70-77	0.1	
157	Does age influence disease-specific survival in patients with squamous cell carcinomas of the head and neck?. <i>Journal of Surgical Oncology</i> , 2020 , 121, 1058-1066	2.8	3
156	Weighted lymph node ratio: New tool in the assessment of postoperative staging of the neck dissection in HPV-negative head and neck squamous cell carcinoma patients. <i>Head and Neck</i> , 2020 , 42, 2912-2919	4.2	5
155	Salvage surgery in head and neck cancer: External validation of predictors of disease-specific survival. <i>Oral Oncology</i> , 2020 , 109, 104876	4.4	4
154	Risk of onset of second neoplasms and successive neoplasms in patients with a head and neck index tumour. <i>Acta Otorrinolaringologica Española</i> , 2020 , 71, 9-15	0.9	6

153	Predictive capacity of IL-8 expression in head and neck squamous carcinoma patients treated with radiotherapy or chemoradiotherapy. <i>Acta Otorrinolaringológica Española</i> , 2020 , 72, 337-337	0.9	0
152	Prognostic capacity of the lymph node ratio in squamous cell carcinomas of the head and neck. <i>Acta Otorrinolaringológica Española</i> , 2020 , 71, 265-274	0.9	
151	Analysis of specific survival and local control through a recursive partitioning analysis in patients with head and neck carcinoma. <i>Acta Otorrinolaringológica Española</i> , 2020 , 71, 131-139	0.9	
150	Prognostic value of nodal involvement in patients with oropharyngeal carcinoma according to the HPV status. <i>Acta Otorrinolaringológica Española</i> , 2020 , 71, 212-218	0.9	
149	Oncological results of salvage laryngectomy in patients with laryngeal carcinoma. <i>Acta Otorrinolaringológica Española</i> , 2020 , 71, 70-77	0.9	4
148	Prognostic significance of albumin levels prior to treatment in patients with head and neck squamous cell carcinoma. <i>Acta Otorrinolaringológica Española</i> , 2020 , 71, 204-211	0.9	1
147	Pretreatment peripheral blood leukocytes are independent predictors of survival in oral cavity cancer. <i>Cancer</i> , 2020 , 126, 994-1003	6.4	26
146	The risk of second primary tumors in head and neck cancer: A systematic review. <i>Head and Neck</i> , 2020 , 42, 456-466	4.2	20
145	Prognostic capacity of Systemic Inflammation Response Index (SIRI) in patients with head and neck squamous cell carcinoma. <i>Head and Neck</i> , 2020 , 42, 336-343	4.2	16
144	Prognostic Value of Nodal Involvement in Patients With Oropharyngeal Carcinoma According to the HPV Status. <i>Acta Otorrinolaringologica (English Edition)</i> , 2020 , 71, 212-218	0.1	
143	Prognostic Significance of Albumin Levels Prior to Treatment in Patients With Head and Neck Squamous Cell Carcinoma. <i>Acta Otorrinolaringologica (English Edition)</i> , 2020 , 71, 204-211	0.1	
142	Prognostic Capacity of the Lymph Node Ratio in Squamous Cell Carcinomas of the Head and Neck. <i>Acta Otorrinolaringologica (English Edition)</i> , 2020 , 71, 265-274	0.1	
141	Management of the Neck in Well-Differentiated Thyroid Cancer. <i>Current Oncology Reports</i> , 2020 , 23, 1	6.3	8
140	Patients with locally advanced hypopharyngeal carcinoma. Results over a 30-year period. <i>Acta Otorrinolaringológica Española</i> , 2019 , 70, 315-326	0.9	4
139	Analysis of the Clinical Relevance of Histological Classification of Benign Epithelial Salivary Gland Tumours. <i>Advances in Therapy</i> , 2019 , 36, 1950-1974	4.1	24
138	Prognostic and predictive factors in recurrent and/or metastatic head and neck squamous cell carcinoma: A review of the literature. <i>Critical Reviews in Oncology/Hematology</i> , 2019 , 137, 84-91	7	24
137	Absence of disruptive TP53 mutations in high-risk human papillomavirus-driven neck squamous cell carcinoma of unknown primary. <i>Head and Neck</i> , 2019 , 41, 3833-3841	4.2	0
136	Indications and results of extended total laryngectomy with en-bloc resection of overlying cervical skin. <i>European Archives of Oto-Rhino-Laryngology</i> , 2019 , 276, 3179-3184	3.5	2

135	Epidemiologic Characteristics of Squamous Head and Neck Carcinoma Patients. Results of a Hospital Register. <i>Acta Otorrinolaringologica (English Edition)</i> , 2019 , 70, 272-278	0.1	
134	Risk of second primary neoplasia in patients with oropharyngeal carcinoma: Role of HPV status in the outcome.. <i>Journal of Clinical Oncology</i> , 2019 , 37, e17544-e17544	2.2	
133	The combined use of EFS, GPX2, and SPRR1A expression could distinguish favorable from poor clinical outcome among epithelial-like head and neck carcinoma subtypes. <i>Head and Neck</i> , 2019 , 41, 1830-1845	4.2	5
132	Patients With Locally Advanced Hypopharyngeal Carcinoma. Results Over a 30-year Period. <i>Acta Otorrinolaringologica (English Edition)</i> , 2019 , 70, 315-326	0.1	
131	Pharyngocutaneous fistula after total laryngectomy: multivariate analysis of risk factors and a severity-based classification proposal. <i>European Archives of Oto-Rhino-Laryngology</i> , 2019 , 276, 143-151	3.5	18
130	Epidemiologic characteristics of squamous head and neck carcinoma patients. Results of a hospital register. <i>Acta Otorrinolaringologica Española</i> , 2019 , 70, 272-278	0.9	13
129	Surgical management of intrathoracic goitres. <i>European Archives of Oto-Rhino-Laryngology</i> , 2019 , 276, 305-314	3.5	14
128	Competing mortality in oropharyngeal carcinoma according to human papillomavirus status. <i>Head and Neck</i> , 2019 , 41, 1328-1334	4.2	7
127	Repair of post-laryngectomy pharyngocutaneous fistulas using a pectoralis major flap. <i>Brazilian Journal of Otorhinolaryngology</i> , 2019 , 85, 351-356	1.6	6
126	Contemporary management of primary parapharyngeal space tumors. <i>Head and Neck</i> , 2019 , 41, 522-535	4.2	27
125	Effect of serpinE1 overexpression on the primary tumor and lymph node, and lung metastases in head and neck squamous cell carcinoma. <i>Head and Neck</i> , 2019 , 41, 429-439	4.2	12
124	Double positivity for HPV-DNA/p16 is the biomarker with strongest diagnostic accuracy and prognostic value for human papillomavirus related oropharyngeal cancer patients. <i>Oral Oncology</i> , 2018 , 78, 137-144	4.4	38
123	Immunohistochemical Expression of Cortactin and Focal Adhesion Kinase Predicts Recurrence Risk and Laryngeal Cancer Risk Beyond Histologic Grading. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2018 , 27, 805-813	4	11
122	Head and Neck Cancer: A Review of the Impact of Treatment Delay on Outcome. <i>Advances in Therapy</i> , 2018 , 35, 153-160	4.1	38
121	Detection of Distant Metastases in Head and Neck Cancer: Changing Landscape. <i>Advances in Therapy</i> , 2018 , 35, 161-172	4.1	10
120	Selective neck dissection in surgically treated head and neck squamous cell carcinoma patients with a clinically positive neck: Systematic review. <i>European Journal of Surgical Oncology</i> , 2018 , 44, 395-403	3.6	25
119	Postoperative staging of the neck dissection using extracapsular spread and lymph node ratio as prognostic factors in HPV-negative head and neck squamous cell carcinoma patients. <i>Oral Oncology</i> , 2018 , 77, 37-42	4.4	17
118	Impact of Total Laryngectomy on Return to Work. <i>Acta Otorrinolaringologica (English Edition)</i> , 2018 , 69, 74-79	0.1	2

117	Impact of total laryngectomy on return to work. <i>Acta Otorrinolaringológica Española</i> , 2018 , 69, 74-79	0.9	9
116	Prognostic significance of extranodal extension in head and neck squamous cell carcinoma cN0 patients with occult metastatic neck nodes. <i>Acta Otorrinolaringológica Española</i> , 2018 , 69, 156-164	0.9	10
115	Prognostic Significance of Extranodal Extension in Head and Neck Squamous Cell Carcinoma cN0 Patients With Occult Metastatic Neck Nodes. <i>Acta Otorrinolaringologica (English Edition)</i> , 2018 , 69, 156-164	0.9	1
114	Prognostic value of CD45 transcriptional expression in head and neck cancer. <i>European Archives of Oto-Rhino-Laryngology</i> , 2018 , 275, 225-232	3.5	3
113	Elective neck dissection during salvage surgery after radiotherapy in patients with head and neck squamous cell carcinoma. <i>Acta Otorhinolaryngologica Italica</i> , 2018 , 38, 86-93	2.8	7
112	Carotid blowout syndrome: modern trends in management. <i>Cancer Management and Research</i> , 2018 , 10, 5617-5628	3.6	50
111	Management of the facial nerve in parotid cancer: preservation or resection and reconstruction. <i>European Archives of Oto-Rhino-Laryngology</i> , 2018 , 275, 2615-2626	3.5	31
110	Prognostic role of extracapsular spread in planned neck dissection after chemoradiotherapy. <i>Head and Neck</i> , 2018 , 40, 2514-2520	4.2	1
109	Polymorphous adenocarcinoma of the salivary glands: reappraisal and update. <i>European Archives of Oto-Rhino-Laryngology</i> , 2018 , 275, 1681-1695	3.5	25
108	Human papillomavirus as prognostic marker with rising prevalence in neck squamous cell carcinoma of unknown primary: A retrospective multicentre study. <i>European Journal of Cancer</i> , 2017 , 74, 73-81	7.5	47
107	How Phenotype Guides Management of the Most Common Malignant Salivary Neoplasms of the Larynx?. <i>Advances in Therapy</i> , 2017 , 34, 813-825	4.1	2
106	Prognostic significance of extracapsular spread in isolated neck recurrences in head and neck squamous cell carcinoma patients. <i>European Archives of Oto-Rhino-Laryngology</i> , 2017 , 274, 527-533	3.5	7
105	Validation of the pathological classification of lymph node metastasis for head and neck tumors according to the 8th edition of the TNM Classification of Malignant Tumors. <i>Oral Oncology</i> , 2017 , 70, 29-33	4.4	25
104	How phenotype guides management of non-conventional squamous cell carcinomas of the larynx?. <i>European Archives of Oto-Rhino-Laryngology</i> , 2017 , 274, 2709-2726	3.5	13
103	The role of HPV on the risk of second primary neoplasia in patients with oropharyngeal carcinoma. <i>Oral Oncology</i> , 2017 , 64, 37-43	4.4	25
102	A Novel Role For Nanog As An Early Cancer Risk Marker In Patients With Laryngeal Precancerous Lesions. <i>Scientific Reports</i> , 2017 , 7, 11110	4.9	20
101	Pretreatment count of peripheral neutrophils, monocytes, and lymphocytes as independent prognostic factor in patients with head and neck cancer. <i>Head and Neck</i> , 2017 , 39, 219-226	4.2	53
100	The prognostic value of pretreatment platelet count in patients with head and neck squamous cell carcinoma. <i>Auris Nasus Larynx</i> , 2017 , 44, 313-318	2.2	11

99	Long-Term Hearing Outcomes following Stapedotomy in Patients with Otosclerosis and Preoperative Small Air-Bone Gap. <i>Audiology and Neuro-Otology</i> , 2017 , 22, 350-355	2.2	7
98	Relationship between response to induction chemotherapy and disease control in patients with advanced laryngeal carcinoma included in an organ preservation protocol. <i>European Archives of Oto-Rhino-Laryngology</i> , 2017 , 274, 2581-2587	3.5	8
97	Surgical options in benign parotid tumors: a proposal for classification. <i>European Archives of Oto-Rhino-Laryngology</i> , 2017 , 274, 3825-3836	3.5	36
96	Validation of the pathological classification of lymph node metastasis for head and neck tumors (HNSCC) according to the 8th edition of the TNM Classification of Malignant Tumors (8th TNM).. <i>Journal of Clinical Oncology</i> , 2017 , 35, 6014-6014	2.2	
95	Oropharyngeal and laryngeal sensory innervation in the pathophysiology of swallowing disorders and sensory stimulation treatments. <i>Annals of the New York Academy of Sciences</i> , 2016 , 1380, 104-120	6.5	25
94	Expression of the CXCL12/CXCR4 chemokine axis predicts regional control in head and neck squamous cell carcinoma. <i>European Archives of Oto-Rhino-Laryngology</i> , 2016 , 273, 4525-4533	3.5	12
93	CKMT1 and NCOA1 expression as a predictor of clinical outcome in patients with advanced-stage head and neck squamous cell carcinoma. <i>Head and Neck</i> , 2016 , 38 Suppl 1, E1392-403	4.2	13
92	HPV Involvement in Head and Neck Cancers: Comprehensive Assessment of Biomarkers in 3680 Patients. <i>Journal of the National Cancer Institute</i> , 2016 , 108, djv403	9.7	441
91	Classification of parotidectomies: a proposal of the European Salivary Gland Society. <i>European Archives of Oto-Rhino-Laryngology</i> , 2016 , 273, 3307-12	3.5	70
90	Risk of distant metastases in head and neck carcinoma patients and myeloperoxidase (MPO) expression.. <i>Journal of Clinical Oncology</i> , 2016 , 34, 6067-6067	2.2	1
89	External validation of sTWEAK as a prognostic noninvasive biomarker for head and neck squamous cell carcinoma. <i>Head and Neck</i> , 2016 , 38 Suppl 1, E1358-63	4.2	8
88	Reply to the letter "A proposal for a level for parapharyngeal extension of parotid gland". <i>European Archives of Oto-Rhino-Laryngology</i> , 2016 , 273, 3457	3.5	
87	Management of recurrent head and neck cancer: variables related to salvage surgery. <i>European Archives of Oto-Rhino-Laryngology</i> , 2016 , 273, 4417-4424	3.5	34
86	Reply to the letter to the editor concerning: 'Classification of parotidectomies: a proposal of the European Salivary Gland Society'. <i>European Archives of Oto-Rhino-Laryngology</i> , 2016 , 273, 3453	3.5	1
85	Parotid incidentaloma identified by positron emission/computed tomography: when to consider diagnoses other than warthin tumor. <i>International Archives of Otorhinolaryngology</i> , 2015 , 19, 112-5	1.5	7
84	Reconstruction of oral cavity defects with FAMM (facial artery musculomucosal) flaps. Our experience. <i>Acta Otorrinolaringológica Española</i> , 2015 , 66, 275-80	0.9	7
83	Salvage surgery after local recurrence in patients with head and neck carcinoma treated with chemoradiotherapy or bioradiotherapy. <i>Auris Nasus Larynx</i> , 2015 , 42, 145-9	2.2	14
82	Prostacyclin-synthase expression in head and neck carcinoma patients and its prognostic value in the response to radiotherapy. <i>Journal of Pathology</i> , 2015 , 235, 125-35	9.4	12

81	Enhanced cell migration and apoptosis resistance may underlie the association between high SERPINE1 expression and poor outcome in head and neck carcinoma patients. <i>Oncotarget</i> , 2015 , 6, 29016-33	3.3	45
80	Expression of IL-1 α correlates with distant metastasis in patients with head and neck squamous cell carcinoma. <i>Oncotarget</i> , 2015 , 6, 37398-409	3.3	32
79	Expression of the CXCL12/CXCR4 chemokine axis as predictor of regional control in head and neck squamous cell carcinoma (HNSCC).. <i>Journal of Clinical Oncology</i> , 2015 , 33, e17093-e17093	2.2	
78	European Laryngological Society: ELS recommendations for the follow-up of patients treated for laryngeal cancer. <i>European Archives of Oto-Rhino-Laryngology</i> , 2014 , 271, 2469-79	3.5	46
77	Aetiology and Treatment of Vocal Fold Paralysis: Retrospective Study of 108 Patients. <i>Acta Otorrinolaringologica (English Edition)</i> , 2014 , 65, 225-230	0.1	1
76	Stomal recurrence in head and neck cancer patients with temporary tracheostomy. <i>Auris Nasus Larynx</i> , 2014 , 41, 467-70	2.2	12
75	Elective treatment of the neck for second primary tumors of the head and neck. <i>European Archives of Oto-Rhino-Laryngology</i> , 2014 , 271, 1187-90	3.5	6
74	Expression of interleukin-1 α and risk of distant metastases (DM) in head and neck squamous cell carcinoma (HNSCC).. <i>Journal of Clinical Oncology</i> , 2014 , 32, 6078-6078	2.2	
73	Overexpression of the nuclear factor-kappa B (p65) in association with local failure in patients with head and neck carcinoma undergoing radiotherapy or chemoradiotherapy. <i>Head and Neck</i> , 2013 , 35, 370-4	4.2	8
72	Identification of somatic VHL gene mutations in sporadic head and neck paragangliomas in association with activation of the HIF-1 α /miR-210 signaling pathway. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013 , 98, E1661-6	5.6	15
71	Development and (pre-) clinical assessment of a novel surgical tool for primary and secondary tracheoesophageal puncture with immediate voice prosthesis insertion, the Provox Vega Puncture Set. <i>European Archives of Oto-Rhino-Laryngology</i> , 2013 , 270, 255-62	3.5	11
70	Prognostic role of MMP-9 expression in head and neck carcinoma patients treated with radiotherapy or chemoradiotherapy. <i>Oral Oncology</i> , 2013 , 49, 322-5	4.4	19
69	Inclusion of extracapsular spread in the pTNM classification system: a proposal for patients with head and neck carcinoma. <i>JAMA Otolaryngology - Head and Neck Surgery</i> , 2013 , 139, 483-8	3.9	69
68	High RAB25 expression is associated with good clinical outcome in patients with locally advanced head and neck squamous cell carcinoma. <i>Cancer Medicine</i> , 2013 , 2, 950-63	4.8	11
67	Interaction between head and neck squamous cell carcinoma cells and fibroblasts in the biosynthesis of PGE2. <i>Journal of Lipid Research</i> , 2012 , 53, 630-42	6.3	31
66	Second, third, and fourth head and neck tumors. A progressive decrease in survival. <i>Head and Neck</i> , 2012 , 34, 1716-9	4.2	22
65	Delayed carotid blow-out syndrome: a new complication of chemoradiotherapy treatment in pharyngolaryngeal carcinoma. <i>Journal of Laryngology and Otology</i> , 2012 , 126, 1189-91	1.8	12
64	Gene expression signatures and molecular markers associated with clinical outcome in locally advanced head and neck carcinoma. <i>Carcinogenesis</i> , 2012 , 33, 1707-16	4.6	24

63	Comparison of chi-squared automatic interaction detection classification trees vs TNM classification for patients with head and neck squamous cell carcinoma. <i>JAMA Otolaryngology</i> , 2012 , 138, 272-9		4
62	Progressive increasing in the risk of second and subsequent malignant tumors in patients with a head and neck cancer: A validation study with SEER data.. <i>Journal of Clinical Oncology</i> , 2012 , 30, 5595-5595 ²		
61	Gene expression as a predictor of radiotherapy or chemoradiotherapy response in head and neck squamous cell carcinoma patients: A molecular approach.. <i>Journal of Clinical Oncology</i> , 2012 , 30, 5590-5590 ²		
60	Salvage surgery after locoregional failure in head and neck carcinoma patients treated with chemoradiotherapy. <i>European Archives of Oto-Rhino-Laryngology</i> , 2011 , 268, 295-301	3.5	50
59	Total or subtotal glossectomy with microsurgical reconstruction: functional and oncological results. <i>Microsurgery</i> , 2011 , 31, 517-23	2.1	26
58	Parotidectomies in benign parotid tumours: Bant Pau surgical extension classification. <i>Acta Otorrinolaringologica (English Edition)</i> , 2010 , 61, 1-5	0.1	1
57	Parotidectomies in benign parotid tumours: "Sant Pau" surgical extension classification. <i>Acta Otorrinolaringologica Española</i> , 2010 , 61, 1-5	0.9	8
56	Risk of third and fourth tumors in patients with head and neck cancer. <i>Head and Neck</i> , 2010 , 32, 1467-72	4.2	11
55	Resultados de un protocolo de preservaci3n de hgano en pacientes con tumores localmente avanzados de la laringe. <i>Acta Otorrinolaringologica Española</i> , 2009 , 60, 176-185	0.9	10
54	Ceratocricoid muscle: an embryological and anatomical study. <i>Clinical Anatomy</i> , 2009 , 22, 463-70	2.5	7
53	Endoscopic supraglottic laryngectomy: a proposal for a classification by the working committee on nomenclature, European Laryngological Society. <i>European Archives of Oto-Rhino-Laryngology</i> , 2009 , 266, 993-8	3.5	75
52	Influence of the persistence of tobacco and alcohol use in the appearance of second neoplasm in patients with a head and neck cancer. A case-control study. <i>Cancer Causes and Control</i> , 2009 , 20, 645-52	2.8	59
51	Results of an organ conservation protocol in patients with locally advanced laryngeal tumours. <i>Acta Otorrinolaringologica (English Edition)</i> , 2009 , 60, 176-185	0.1	2
50	Total Laryngectomy 2009 , 229-243		
49	Potential structures that could be confused with a nonrecurrent inferior laryngeal nerve: an anatomic study. <i>Laryngoscope</i> , 2008 , 118, 56-60	3.6	29
48	Trasplante de la laringe: ¿una opci3n terap3utica para el siglo XXI? Revisi3n de la literatura. <i>Acta Otorrinolaringologica Española</i> , 2008 , 59, 127-138	0.9	0
47	Ku70 predicts response and primary tumor recurrence after therapy in locally advanced head and neck cancer. <i>International Journal of Cancer</i> , 2008 , 123, 1068-79	7.5	36
46	Lymphadenopathy caused by Mycobacterium colombiense. <i>Journal of Clinical Microbiology</i> , 2008 , 46, 1885-7	9.7	25

45	Recurrent laryngeal squamous cell carcinoma: rTNM versus composite laryngeal recurrence staging system. Proposal for a modification of the CLRSS to improve patient classification. <i>Head and Neck</i> , 2008 , 30, 939-45	4.2	7
44	Prostaglandin E(2) pathway in head and neck squamous cell carcinoma. <i>Head and Neck</i> , 2008 , 30, 1175-81	4.2	39
43	Organ Preservation in Laryngeal and Hypopharyngeal Cancer. <i>Acta Otorrinolaringologica (English Edition)</i> , 2007 , 58, 476-482	0.1	0
42	External validation of a risk group defined by recursive partitioning analysis in patients with head and neck carcinoma treated with surgery and postoperative radiotherapy. <i>Head and Neck</i> , 2007 , 29, 815-21	4.2	15
41	Modalidades de preservaci3n de 3rgano en carcinomas de laringe e hipofaringe. <i>Acta Otorrinolaringologica Espa3ola</i> , 2007 , 58, 476-482	0.9	8
40	Supracricoid laryngectomy as salvage surgery after failure of radiation therapy. <i>European Archives of Oto-Rhino-Laryngology</i> , 2007 , 264, 809-14	3.5	27
39	Proposal for revision of the European Laryngological Society classification of endoscopic cordectomies. <i>European Archives of Oto-Rhino-Laryngology</i> , 2007 , 264, 499-504	3.5	205
38	A proposal of pathological classification (pTNM) of patients with head and neck cancer according to capsular rupture (pN- RC). <i>Journal of Clinical Oncology</i> , 2007 , 25, 16525-16525	2.2	
37	Mutation analysis of the SDHD gene in four kindreds with familial paraganglioma: description of one novel germline mutation. <i>Diagnostic Molecular Pathology</i> , 2005 , 14, 109-14		12
36	Comparison of the Radiation Therapy Oncology Group recursive partitioning classification and Union Internationale Contre le Cancer TNM classification for patients with head and neck carcinoma. <i>Head and Neck</i> , 2005 , 27, 248-57	4.2	8
35	Variability in nerve patterns of the adductor muscle group supplied by the recurrent laryngeal nerve. <i>Laryngoscope</i> , 2005 , 115, 358-62	3.6	52
34	Incidence and significance of clinically unsuspected thyroid tissue in lymph nodes found during neck dissection in head and neck carcinoma patients. <i>Laryngoscope</i> , 2005 , 115, 470-4	3.6	36
33	Results of an organ preservation protocol with induction chemotherapy and radiotherapy in patients with locally advanced laryngeal carcinoma. <i>European Archives of Oto-Rhino-Laryngology</i> , 2005 , 262, 93-8	3.5	14
32	Metachronous second primary tumours in the aerodigestive tract in patients with early stage head and neck squamous cell carcinomas. <i>European Archives of Oto-Rhino-Laryngology</i> , 2005 , 262, 905-9	3.5	30
31	The role of type of tobacco and type of alcoholic beverage in oral carcinogenesis. <i>International Journal of Cancer</i> , 2004 , 108, 741-9	7.5	186
30	Distant metastases in squamous-cell head and neck carcinoma (SHNC) patients who achieved loco-regional control (LRC). <i>Journal of Clinical Oncology</i> , 2004 , 22, 5519-5519	2.2	
29	Smoke and alcohol consumption as a risk factors in the development of second primary neoplasms (SPN) in head & neck cancer (HNC) patients. A case-control study. <i>Journal of Clinical Oncology</i> , 2004 , 22, 5582-5582	2.2	
28	Distant metastases in squamous-cell head and neck carcinoma (SHNC) patients who achieved loco-regional control (LRC). <i>Journal of Clinical Oncology</i> , 2004 , 22, 5519-5519	2.2	

27	Smoke and alcohol consumption as a risk factors in the development of second primary neoplasms (SPN) in head & neck cancer (HNC) patients. A case-control study. <i>Journal of Clinical Oncology</i> , 2004 , 22, 5582-5582	2.2	
26	Is the external laryngeal nerve an exclusively motor nerve? The cricothyroid connection branch. <i>Laryngoscope</i> , 2003 , 113, 525-9	3.6	73
25	Variability of the nerve supply patterns of the human posterior cricoarytenoid muscle. <i>Laryngoscope</i> , 2003 , 113, 602-6	3.6	42
24	The effect of waiting time on local control and survival in head and neck carcinoma patients treated with radiotherapy. <i>Radiotherapy and Oncology</i> , 2003 , 66, 277-81	5.3	34
23	Results of an organ preservation protocol with induction chemotherapy and radiotherapy in patients with locally advanced pyriform sinus carcinoma. <i>European Archives of Oto-Rhino-Laryngology</i> , 2002 , 259, 32-6	3.5	21
22	Hypothyroidism in patients treated with total laryngectomy. A multivariate study. <i>European Archives of Oto-Rhino-Laryngology</i> , 2002 , 259, 193-6	3.5	40
21	Results of salvage surgery for local or regional recurrence after larynx preservation with induction chemotherapy and radiotherapy. <i>Head and Neck</i> , 2001 , 23, 733-8	4.2	41
20	Selective dissection of levels II-III with intraoperative control of the upper and middle jugular nodes: a therapeutic option for the N0 neck. <i>Head and Neck</i> , 2001 , 23, 441-6	4.2	36
19	Can cure be achieved in patients with head and neck carcinomas? The problem of second neoplasm. <i>Expert Review of Anticancer Therapy</i> , 2001 , 1, 125-33	3.5	20
18	Endoscopic laser surgery in the treatment of radiation failure of early laryngeal carcinoma. <i>Head and Neck</i> , 2000 , 22, 520-3	4.2	44
17	Distant metastases in head and neck cancer patients who achieved loco-regional control. <i>Head and Neck</i> , 2000 , 22, 680-6	4.2	177
16	Histologically negative specimens after induction therapy: frequency and impact on survival. <i>Head and Neck</i> , 2000 , 22, 808-13	4.2	5
15	How much does it cost to preserve a larynx? An economic study. <i>European Archives of Oto-Rhino-Laryngology</i> , 2000 , 257, 72-6	3.5	15
14	Initial treatment of the early stages (I, II) of supraglottic squamous cell carcinoma: partial laryngectomy versus radiotherapy. <i>European Archives of Oto-Rhino-Laryngology</i> , 2000 , 257, 512-6	3.5	28
13	Treatment of neck nodes after induction chemotherapy in patients with primary advanced tumours. <i>European Archives of Oto-Rhino-Laryngology</i> , 2000 , 257, 521-5	3.5	7
12	Montgomery salivary bypass tube in the reconstruction of the hypopharynx. Cost-benefit study. <i>Annals of Otology, Rhinology and Laryngology</i> , 1999 , 108, 864-8	2.1	31
11	An anatomical study of anastomoses between the laryngeal nerves. <i>Laryngoscope</i> , 1999 , 109, 983-7	3.6	131
10	Second neoplasm in patients with head and neck cancer. <i>Head and Neck</i> , 1999 , 21, 204-10	4.2	235

9	Influence of age on laryngeal carcinoma. <i>Annals of Otolaryngology, Rhinology and Laryngology</i> , 1998 , 107, 164-9	2.1	17
8	Foramen thyroideum: a comparative study in embryos, fetuses, and adults. <i>Laryngoscope</i> , 1997 , 107, 1146-50	3.6	15
7	Laryngeal carcinoma in patients without a history of tobacco and alcohol use. <i>Head and Neck</i> , 1997 , 19, 200-4	4.2	26
6	Prevention of stomal recurrence. <i>Head and Neck</i> , 1996 , 18, 54-9	4.2	37
5	Blood transfusions in laryngeal cancer: effect on prognosis. <i>Head and Neck</i> , 1996 , 18, 218-24	4.2	6
4	Prevention of stomal recurrence 1996 , 18, 54		2
3	Function preservation in stage III squamous laryngeal carcinoma: results with an induction chemotherapy protocol. <i>Laryngoscope</i> , 1995 , 105, 822-6	3.6	23
2	Primary tracheoesophageal puncture vs esophageal speech. <i>JAMA Otolaryngology</i> , 1992 , 118, 188-90		22
1	Pharyngocutaneous fistula in irradiated patients: systematic review and our experience.. <i>Journal of Laryngology and Otolaryngology</i> , 1-23	1.8	