

List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/8409898/xucong-lin-publications-by-citations.pdf>  
**Version:** 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.  
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

77 papers	1,345 citations	23 h-index	31 g-index
79 ext. papers	1,485 ext. citations	4.3 avg, IF	4.51 L-index

#	Paper	IF	Citations
77	Determination of pyrethroid pesticide residues in vegetables by pressurized capillary electrochromatography. <i>Talanta</i> , <b>2006</b> , 69, 97-102	6.2	86
76	Preparation and evaluation of a neutral methacrylate-based monolithic column for hydrophilic interaction stationary phase by pressurized capillary electrochromatography. <i>Journal of Chromatography A</i> , <b>2009</b> , 1216, 4611-7	4.5	47
75	Novel highly hydrophilic methacrylate-based monolithic column with mixed-mode of hydrophilic and strong cation-exchange interactions for pressurized capillary electrochromatography. <i>Journal of Chromatography A</i> , <b>2011</b> , 1218, 4671-7	4.5	45
74	Investigation of enantiomer recognition of molecularly imprinted polymeric monoliths in pressurized capillary electrochromatography screening the amino acids and their derivatives. <i>Journal of Chromatography A</i> , <b>2009</b> , 1216, 5320-6	4.5	44
73	Electrochromatographic characterization of methacrylate-based monolith with mixed mode of hydrophilic and weak electrostatic interactions by pressurized capillary electrochromatography. <i>Journal of Chromatography A</i> , <b>2008</b> , 1190, 365-71	4.5	41
72	On-line coupling of pressurized capillary electrochromatography with end-column amperometric detection for analysis of estrogens. <i>Electrophoresis</i> , <b>2005</b> , 26, 2342-50	3.6	39
71	Preparation of a mixed-mode hydrophilic interaction/anion-exchange polymeric monolithic stationary phase for capillary liquid chromatography of polar analytes. <i>Journal of Chromatography A</i> , <b>2009</b> , 1216, 801-6	4.5	38
70	Methacrylate-based monolithic column with mixed-mode hydrophilic interaction/strong cation-exchange stationary phase for capillary liquid chromatography and pressure-assisted CEC. <i>Electrophoresis</i> , <b>2008</b> , 29, 4055-65	3.6	38
69	End-column chemiluminescence detection for pressurized capillary electrochromatographic analysis of norepinephrine and epinephrine. <i>Journal of Chromatography A</i> , <b>2007</b> , 1170, 118-21	4.5	35
68	Separation of structurally related estrogens using isocratic elution pressurized capillary electrochromatography. <i>Journal of Chromatography A</i> , <b>2005</b> , 1092, 258-62	4.5	35
67	Aptamer-based polyhedral oligomeric silsesquioxane (POSS)-containing hybrid affinity monolith prepared via a "one-pot" process for selective extraction of ochratoxin A. <i>Journal of Chromatography A</i> , <b>2018</b> , 1563, 37-46	4.5	35
66	Preparation and characterization of hybrid-silica monolithic column with mixed-mode of hydrophilic and strong anion-exchange interactions for pressurized capillary electrochromatography. <i>Journal of Chromatography A</i> , <b>2012</b> , 1239, 56-63	4.5	31
65	Sulfoalkylbetaine-based monolithic column with mixed-mode of hydrophilic interaction and strong anion-exchange stationary phase for capillary electrochromatography. <i>Electrophoresis</i> , <b>2010</b> , 31, 2997-3005	3.6	31
64	Separation and determination of five major opium alkaloids with mixed mode of hydrophilic/cation-exchange monolith by pressurized capillary electrochromatography. <i>Journal of Separation Science</i> , <b>2007</b> , 30, 3011-7	3.4	31
63	Development of a new method for analysis of Sudan dyes by pressurized CEC with amperometric detection. <i>Electrophoresis</i> , <b>2007</b> , 28, 1696-703	3.6	29
62	Preparation and evaluation of a sulfoalkylbetaine-based zwitterionic monolithic column for CEC of polar analytes. <i>Electrophoresis</i> , <b>2009</b> , 30, 2702-10	3.6	26
61	Rapid fabrication of ionic liquid-functionalized monolithic column via in-situ urea-formaldehyde polycondensation for pressurized capillary electrochromatography. <i>Journal of Chromatography A</i> , <b>2016</b> , 1449, 100-8	4.5	26

60	Electroneutral silica-based hybrid monolith for hydrophilic interaction capillary electrochromatography. <i>Journal of Chromatography A</i> , <b>2012</b> , 1260, 174-82	4.5	25
59	On-column coaxial flow chemiluminescence detection for underivatized amino acids by pressurized capillary electrochromatography using a monolithic column. <i>Analytical Chemistry</i> , <b>2006</b> , 78, 5322-8	7.8	25
58	A facile AuNPs@aptamer-modified mercaptosiloxane-based hybrid affinity monolith with an unusually high coverage density of aptamer for on-column selective extraction of ochratoxin A. <i>Analyst, The</i> , <b>2018</b> , 143, 5210-5217	5	25
57	Urea-formaldehyde monolithic column for hydrophilic in-tube solid-phase microextraction of aminoglycosides. <i>Journal of Chromatography A</i> , <b>2017</b> , 1485, 24-31	4.5	24
56	Preparation and evaluation of highly hydrophilic aptamer-based hybrid affinity monolith for on-column specific discrimination of ochratoxin A. <i>Talanta</i> , <b>2019</b> , 200, 193-202	6.2	24
55	An aptamer@AuNP-modified POSS-polyethylenimine hybrid affinity monolith with a high aptamer coverage density for sensitive and selective recognition of ochratoxin A. <i>Journal of Materials Chemistry B</i> , <b>2018</b> , 6, 1965-1972	7.3	24
54	Rapid capillary electrochromatographic profiling of phytohormones on a hydrophilic interaction/strong anion-exchange mixed-mode monolith. <i>Analyst, The</i> , <b>2013</b> , 138, 635-41	5	23
53	Monolithic column with double mixed-modes of hydrophilic interaction/cation-exchange and reverse-phase/cation-exchange stationary phase for pressurized capillary electrochromatography. <i>Electrophoresis</i> , <b>2008</b> , 29, 928-35	3.6	23
52	PVC matrix membrane sensor for fluorescent determination of phosphate. <i>Talanta</i> , <b>2006</b> , 70, 32-6	6.2	23
51	Branched polyethyleneimine-bonded tentacle-type polymer stationary phase for peptides and proteins separations by open-tubular capillary electrochromatography. <i>Journal of Separation Science</i> , <b>2011</b> , 34, 3383-91	3.4	22
50	Vinylbenzyl quaternary ammonium-based polymeric monolith with hydrophilic interaction/strong anion exchange mixed-mode for pressurized capillary electrochromatography. <i>Journal of Chromatography A</i> , <b>2013</b> , 1316, 104-11	4.5	21
49	Preparation and characterization of a molecularly imprinted monolithic column for pressure-assisted CEC separation of nitroimidazole drugs. <i>Electrophoresis</i> , <b>2010</b> , 31, 2822-30	3.6	20
48	Detection of trans-fatty acids by high performance liquid chromatography coupled with in-tube solid-phase microextraction using hydrophobic polymeric monolith. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , <b>2017</b> , 1040, 214-221	3.2	19
47	Pressurized capillary electrochromatography with indirect amperometric detection for analysis of organophosphorus pesticide residues. <i>Analyst, The</i> , <b>2010</b> , 135, 2150-6	5	18
46	Silver nanoparticles-coated monolithic column for in-tube solid-phase microextraction of monounsaturated fatty acid methyl esters. <i>Journal of Chromatography A</i> , <b>2019</b> , 1585, 19-26	4.5	18
45	Towards online specific recognition and sensitive analysis of bisphenol A by using AuNPs@aptamer hybrid-silica affinity monolithic column with LC-MS. <i>Talanta</i> , <b>2020</b> , 219, 121275	6.2	17
44	Phenylaminopropyl silica monolithic column for pressure assisted capillary electrochromatography. <i>Journal of Chromatography A</i> , <b>2006</b> , 1117, 170-5	4.5	17
43	Dipyridyl-immobilized ionic liquid type hybrid silica monolith for hydrophilic interaction electrochromatography. <i>Electrophoresis</i> , <b>2013</b> , 34, 3091-9	3.6	16

42	Towards high-efficient online specific discrimination of zearalenone by using gold nanoparticles@aptamer-based affinity monolithic column. <i>Journal of Chromatography A</i> , <b>2020</b> , 1620, 461026	4.5	15
41	Separation and determination of structurally related free bile acids by pressurized capillary electrochromatography coupled to laser induced fluorescence detection. <i>Analytical Methods</i> , <b>2010</b> , 2, 1927	3.2	15
40	Quantification of domoic acid in shellfish tissues by pressurized capillary electrochromatography. <i>Journal of Separation Science</i> , <b>2009</b> , 32, 2117-22	3.4	15
39	Sodium hyaluronate-functionalized urea-formaldehyde monolithic column for hydrophilic in-tube solid-phase microextraction of melamine. <i>Journal of Chromatography A</i> , <b>2017</b> , 1515, 54-61	4.5	14
38	Capillary liquid chromatography using a hydrophilic/cation-exchange monolithic column with a dynamically modified cationic surfactant. <i>Journal of Chromatography A</i> , <b>2009</b> , 1216, 7728-31	4.5	14
37	Online high-efficient specific detection of zearalenone in rice by using high-loading aptamer affinity hydrophilic monolithic column coupled with HPLC. <i>Talanta</i> , <b>2020</b> , 219, 121309	6.2	14
36	Highly hydrophilic polyhedral oligomeric silsesquioxane (POSS)-containing aptamer-modified affinity hybrid monolith for efficient on-column discrimination with low nonspecific adsorption. <i>Analyst, The</i> , <b>2019</b> , 144, 1555-1564	5	13
35	Preparation of aptamer-bound polyamine affinity monolithic column via a facile triazine-bridged strategy and application to on-column specific discrimination of ochratoxin A. <i>Journal of Separation Science</i> , <b>2019</b> , 42, 2272-2279	3.4	13
34	Polyhedral oligomeric silsesquioxane (POSS)-based multifunctional organic-silica hybrid monoliths. <i>Analyst, The</i> , <b>2013</b> , 138, 5555-8	5	13
33	Separation of Polar and Basic Compounds in Hydrophilic Interaction Pressurized CEC Using Diethylenetriaminopropyl Silica Monolithic Columns. <i>Chromatographia</i> , <b>2006</b> , 64, 267-272	2.1	13
32	Glycin-bonded silica monolithic column as zwitterionic stationary phase for hydrophilic interaction pressurized capillary electrochromatography. <i>Journal of Separation Science</i> , <b>2009</b> , 32, 2767-75	3.4	12
31	Analysis of phenolic xenoestrogens by pressurized CEC with amperometric detection. <i>Electrophoresis</i> , <b>2010</b> , 31, 1011-8	3.6	12
30	Analysis of phenoxyl-type N-methylcarbamate pesticide residues in vegetables by capillary zone electrophoresis with pre-column hydrolysis and amperometric detection. <i>Journal of Chromatographic Science</i> , <b>2008</b> , 46, 615-21	1.4	12
29	Biomimetic Synthesis of Ag Se Quantum Dots with Enhanced Photothermal Properties and as "Gatekeepers" to Cap Mesoporous Silica Nanoparticles for Chemo-Photothermal Therapy. <i>Chemistry - an Asian Journal</i> , <b>2019</b> , 14, 155-161	4.5	12
28	A facile versatile polymeric monolith for multiple separations. <i>Analyst, The</i> , <b>2013</b> , 138, 771-4	5	10
27	Triamine-bonded stationary phase for open tubular capillary electrochromatography. <i>Journal of Separation Science</i> , <b>2010</b> , 33, 3184-93	3.4	10
26	Evaluation of band broadening in chemiluminescence detection coupled to pressurized capillary electrochromatography with an off-column coaxial flow interface. <i>Electrophoresis</i> , <b>2008</b> , 29, 401-9	3.6	8
25	Rapid analysis of trace levels of flavins by pressurized capillary electrochromatography-laser induced fluorescence detection with sulfonated N-octadecyl methacrylate monolith. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , <b>2010</b> , 53, 1324-31	3.5	7

24	Photocatalytic reduction for graphene oxide by PbTiO with high polarizability and its electrocatalytic application in pyrrole detection. <i>Journal of Colloid and Interface Science</i> , <b>2020</b> , 560, 502-509	3.3	7
23	Quinine-modified polymer monolithic column with reversed-phase /strong anion-exchange mixed-mode for pressurized capillary electrochromatography. <i>Electrophoresis</i> , <b>2018</b> , 39, 1504-1511	3.6	6
22	Phenylalanine functionalized zwitterionic monolith for hydrophobic interaction electrochromatography. <i>Electrophoresis</i> , <b>2013</b> , 34, 3293-9	3.6	6
21	Phenylaminopropyl-functionalized stationary phase for open-tubular capillary electrochromatography of alkaloids and aromatic acids. <i>Journal of Separation Science</i> , <b>2011</b> , 34, 2337-44	3.4	6
20	Integration of fluorescence/photoacoustic imaging and targeted chemo/photothermal therapy with Ag <sub>2</sub> Se@BSA-RGD nanodots. <i>New Journal of Chemistry</i> , <b>2020</b> , 44, 4850-4857	3.6	5
19	Sensitive profiling of trace neurotoxin domoic acid by pressurized capillary electrochromatography with laser-induced fluorescence detection. <i>RSC Advances</i> , <b>2017</b> , 7, 53778-53784	3.7	5
18	Sensitive capillary electrophoretic profiling of nicotine and nor nicotine in mushrooms with amperometric detection. <i>Electrophoresis</i> , <b>2013</b> , 34, 2033-40	3.6	5
17	Preparation of a neutral porous monolith and its evaluation in pressurized capillary electrochromatography with neutral and charged solutes. <i>Electrophoresis</i> , <b>2010</b> , 31, 1674-80	3.6	5
16	Silica-based zwitterionic monolithic stationary phase for separation of neutral and ionized solutes using pressurized CEC. <i>Journal of Separation Science</i> , <b>2010</b> , 33, 1625-32	3.4	4
15	Online specific recognition of mycotoxins using aptamer-grafted ionic affinity monolith with mixed-mode mechanism. <i>Journal of Chromatography A</i> , <b>2021</b> , 1639, 461930	4.5	4
14	Sensitive amperometric detection for capillary electrophoresis of phenol carbamates with in-line thermal hydrolysis strategy. <i>Electrophoresis</i> , <b>2019</b> , 40, 1648-1655	3.6	3
13	A polymer monolith for hydrophilic and dynamically surfactant-modified reversed-phase capillary electrochromatography. <i>RSC Advances</i> , <b>2013</b> , 3, 21888	3.7	3
12	Off-Column Amperometric Detection for Pressurized Capillary Electrochromatography. <i>Chromatographia</i> , <b>2010</b> , 71, 659-665	2.1	3
11	Highly efficient preparation of ECD-based chiral monolithic column by "one-pot" hydroxymethyl polycondensation for enantioseparation in capillary liquid chromatography. <i>Journal of Chromatography A</i> , <b>2020</b> , 1616, 460781	4.5	3
10	Facile preparation of stainless steel microextraction fiber via in situ growth of metal-organic framework UiO-66 and its application to sensitive analysis of polycyclic musks. <i>Journal of Separation Science</i> , <b>2020</b> , 43, 2240-2246	3.4	2
9	Gold Nanoparticle-Decorated Porous Silica for Surface-Enhanced Raman Scattering-Based Detection of Trace Molecules in Liquid Phase. <i>ACS Applied Nano Materials</i> ,	5.6	2
8	Heteroporous 3D covalent organic framework-based magnetic nanospheres for sensitive detection of bisphenol A. <i>Talanta</i> , <b>2021</b> , 231, 122343	6.2	2
7	Aptamer-functionalized metal-organic framework-based electrospun nanofibrous composite coating fiber for specific recognition of ultratrace microcystin in water. <i>Journal of Chromatography A</i> , <b>2021</b> , 1656, 462542	4.5	2

6	Aptamer-functionalized metal-organic framework-coated nanofibers with multi-affinity sites for highly sensitive, selective recognition of ultra-trace microcystin-LR. <i>Talanta</i> , <b>2022</b> , 236, 122880	6.2	2
5	A facile aptamer immobilization strategy to fabricate a robust affinity monolith for highly specific in-tube solid-phase microextraction. <i>Analyst, The</i> , <b>2021</b> , 146, 5732-5739	5	2
4	Towards highly specific aptamer-affinity monolithic column by efficient UV light-initiated polymerization in "one-pot". <i>Analytica Chimica Acta</i> , <b>2021</b> , 1165, 338517	6.6	1
3	Develop potential multi-target drugs by self-assembly of quercetin with amino acids and metal ion to achieve significant efficacy in anti-Alzheimer's disease. <i>Nano Research</i> , 1	10	1
2	In situ photo-initiated polymerized oligonucleotide-functionalized hydrophilic capillary affinity monolith for highly selective in-tube microextraction of ochratoxin A mycotoxin. <i>Mikrochimica Acta</i> , <b>2021</b> , 188, 341	5.8	0
1	Facile DNA adsorption enabling ammonium-based hydrophilic affinity monolithic column for high-performance online selective microextraction of ochratoxin A. <i>Analytica Chimica Acta</i> , <b>2021</b> , 1185, 339077	6.6	