Caryl E Sortwell

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8412436/publications.pdf

Version: 2024-02-01

53 papers

2,100 citations

257101 24 h-index 243296 44 g-index

66 all docs 66
docs citations

66 times ranked 2522 citing authors

#	Article	IF	CITATIONS
1	BDNF rs6265 Genotype Influences Outcomes of Pharmacotherapy and Subthalamic Nucleus Deep Brain Stimulation in Early-Stage Parkinson's Disease. Neuromodulation, 2022, 25, 846-853.	0.4	6
2	Behavioral phenotyping of a rat model of the BDNF Val66Met polymorphism reveals selective impairment of fear memory. Translational Psychiatry, 2022, 12, 93.	2.4	16
3	Dopaminergic Positron Emission Tomography Imaging in the Alphaâ€Synuclein Preformed Fibril Model Reveals Similarities to Early Parkinson's Disease. Movement Disorders, 2022, 37, 1739-1748.	2.2	8
4	Beta2-adrenoreceptor agonist clenbuterol produces transient decreases in alpha-synuclein mRNA but no long-term reduction in protein. Npj Parkinson's Disease, 2022, 8, .	2.5	6
5	Leveraging the preformed fibril model to distinguish between alpha-synuclein inclusion- and nigrostriatal degeneration-associated immunogenicity. Neurobiology of Disease, 2022, 171, 105804.	2.1	12
6	The BDNF Val66Met polymorphism (rs6265) enhances dopamine neuron graft efficacy and side-effect liability in rs6265 knock-in rats. Neurobiology of Disease, 2021, 148, 105175.	2.1	13
7	Striatal Afferent BDNF Is Disrupted by Synucleinopathy and Partially Restored by STN DBS. Journal of Neuroscience, 2021, 41, 2039-2052.	1.7	16
8	Stereotaxic Intracranial Delivery of Chemicals, Proteins or Viral Vectors to Study Parkinson's Disease. Journal of Visualized Experiments, 2021, , .	0.2	0
9	α-Synuclein antisense oligonucleotides as a disease-modifying therapy for Parkinson's disease. JCI Insight, 2021, 6, .	2.3	60
10	Synucleinopathy-associated pathogenesis in Parkinson's disease and the potential for brain-derived neurotrophic factor. Npj Parkinson's Disease, 2021, 7, 35.	2.5	26
11	Preformed fibrils generated from mouse alpha-synuclein produce more inclusion pathology in rats than fibrils generated from rat alpha-synuclein. Parkinsonism and Related Disorders, 2021, 89, 41-47.	1.1	7
12	Next-Generation Diamond Electrodes for Neurochemical Sensing: Challenges and Opportunities. Micromachines, 2021, 12, 128.	1.4	15
13	BDNF rs6265 Variant Alters Outcomes with Levodopa in Early-Stage Parkinson's Disease. Neurotherapeutics, 2020, 17, 1785-1795.	2.1	12
14	Striatal Nurr1, but not FosB expression links a levodopa-induced dyskinesia phenotype to genotype in Fisher 344 vs. Lewis hemiparkinsonian rats. Experimental Neurology, 2020, 330, 113327.	2.0	10
15	Striatal Nurr1 Facilitates the Dyskinetic State and Exacerbates Levodopa-Induced Dyskinesia in a Rat Model of Parkinson's Disease. Journal of Neuroscience, 2020, 40, 3675-3691.	1.7	15
16	Developmental exposure to the organochlorine pesticide dieldrin causes male-specific exacerbation of α-synuclein-preformed fibril-induced toxicity and motor deficits. Neurobiology of Disease, 2020, 141, 104947.	2.1	24
17	Time course and magnitude of alpha-synuclein inclusion formation and nigrostriatal degeneration in the rat model of synucleinopathy triggered by intrastriatal \hat{l}_{\pm} -synuclein preformed fibrils. Neurobiology of Disease, 2019, 130, 104525.	2.1	67
18	Generation of Alpha-Synuclein Preformed Fibrils from Monomers and Use In Vivo. Journal of Visualized Experiments, 2019 , , .	0.2	29

#	Article	IF	CITATIONS
19	Genetic silencing of striatal CaV1.3 prevents and ameliorates levodopa dyskinesia. Movement Disorders, 2019, 34, 697-707.	2.2	19
20	BDNF provides many routes toward STN DBSâ€mediated disease modification. Movement Disorders, 2019, 34, 22-34.	2.2	20
21	Impact of the Aged Brain Environment on Gene Therapy for Parkinson's Disease. , 2018, , 647-657.		0
22	Lewy body-like alpha-synuclein inclusions trigger reactive microgliosis prior to nigral degeneration. Journal of Neuroinflammation, 2018, 15, 129.	3.1	131
23	Quality Over Quantity: Advantages of Using Alpha-Synuclein Preformed Fibril Triggered Synucleinopathy to Model Idiopathic Parkinson's Disease. Frontiers in Neuroscience, 2018, 12, 621.	1.4	32
24	Bdnf variant is associated with milder motor symptom severity in early-stage Parkinson's disease. Parkinsonism and Related Disorders, 2018, 53, 70-75.	1.1	13
25	Best Practices for Generating and Using Alpha-Synuclein Pre-Formed Fibrils to Model Parkinson's Disease in Rodents. Journal of Parkinson's Disease, 2018, 8, 303-322.	1.5	151
26	Silencing Alpha Synuclein in Mature Nigral Neurons Results in Rapid Neuroinflammation and Subsequent Toxicity. Frontiers in Molecular Neuroscience, 2018, 11, 36.	1.4	75
27	Glycomic and Proteomic Changes in Aging Brain Nigrostriatal Pathway. Molecular and Cellular Proteomics, 2018, 17, 1778-1787.	2.5	27
28	Alpha-Synuclein mRNA Is Not Increased in Sporadic PD and Alpha-Synuclein Accumulation Does Not Block GDNF Signaling in Parkinson's Disease and Disease Models. Molecular Therapy, 2017, 25, 2231-2235.	3.7	49
29	Rationally Engineered AAV Capsids Improve Transduction and Volumetric Spread in the CNS. Molecular Therapy - Nucleic Acids, 2017, 8, 184-197.	2.3	48
30	Subthalamic Nucleus Deep Brain Stimulation Employs trkB Signaling for Neuroprotection and Functional Restoration. Journal of Neuroscience, 2017, 37, 6786-6796.	1.7	41
31	Can STN DBS protect both nigral somata and innervation of the striatum?. Annals of Neurology, 2017, 82, 855-855.	2.8	1
32	Nortriptyline inhibits aggregation and neurotoxicity of alpha-synuclein by enhancing reconfiguration of the monomeric form. Neurobiology of Disease, 2017, 106, 191-204.	2.1	28
33	Subthalamic Nucleus Deep Brain Stimulation Does Not Modify the Functional Deficits or Axonopathy Induced by Nigrostriatal α-Synuclein Overexpression. Scientific Reports, 2017, 7, 16356.	1.6	24
34	Impact of age and vector construct on striatal and nigral transgene expression. Molecular Therapy - Methods and Clinical Development, 2016, 3, 16082.	1.8	21
35	Viral Vector-Based Modeling of Neurodegenerative Disorders: Parkinson's Disease. Methods in Molecular Biology, 2016, 1382, 367-382.	0.4	19
36	The Longitudinal Transcriptomic Response of the Substantia Nigra to Intrastriatal 6-Hydroxydopamine Reveals Significant Upregulation of Regeneration-Associated Genes. PLoS ONE, 2015, 10, e0127768.	1.1	18

3

#	Article	IF	Citations
37	Loss of VGLUT3 Produces Circadian-Dependent Hyperdopaminergia and Ameliorates Motor Dysfunction and l-Dopa-Mediated Dyskinesias in a Model of Parkinson's Disease. Journal of Neuroscience, 2015, 35, 14983-14999.	1.7	53
38	Recombinant adenoassociated virus $2/5$ -mediated gene transfer is reduced in the aged rat midbrain. Neurobiology of Aging, 2015, 36, 1110-1120.	1.5	22
39	Tricyclic antidepressant treatment evokes regional changes in neurotrophic factors over time within the intact and degenerating nigrostriatal system. Experimental Neurology, 2015, 266, 11-21.	2.0	22
40	Intrastriatal injection of pre-formed mouse α-synuclein fibrils into rats triggers α-synuclein pathology and bilateral nigrostriatal degeneration. Neurobiology of Disease, 2015, 82, 185-199.	2.1	285
41	Interrogating the aged striatum: Robust survival of grafted dopamine neurons in aging rats produces inferior behavioral recovery and evidence of impaired integration. Neurobiology of Disease, 2015, 77, 191-203.	2.1	24
42	Chronic Amitriptyline Treatment Attenuates Nigrostriatal Degeneration and Significantly Alters Trophic Support in a Rat Model of Parkinsonism. Neuropsychopharmacology, 2015, 40, 874-883.	2.8	21
43	High-Frequency Stimulation of the Rat Entopeduncular Nucleus Does Not Provide Functional or Morphological Neuroprotection from 6-Hydroxydopamine. PLoS ONE, 2015, 10, e0133957.	1.1	14
44	Morphological and Behavioral Impact of AAV2/5-Mediated Overexpression of Human Wildtype Alpha-Synuclein in the Rat Nigrostriatal System. PLoS ONE, 2013, 8, e81426.	1.1	70
45	Striatal Pleiotrophin Overexpression Provides Functional and Morphological Neuroprotection in the 6-Hydroxydopamine Model. Molecular Therapy, 2012, 20, 544-554.	3.7	27
46	Endogenous neural precursors influence grafted neural stem cells and contribute to neuroprotection in the parkinsonian rat. European Journal of Neuroscience, 2012, 35, 883-895.	1.2	17
47	Subthalamic Nucleus Stimulation Increases Brain Derived Neurotrophic Factor in the Nigrostriatal System and Primary Motor Cortex. Journal of Parkinson's Disease, 2011, 1, 123-136.	1.5	64
48	Subthalamic nucleus stimulation increases brain derived neurotrophic factor in the nigrostriatal system and primary motor cortex. Journal of Parkinson's Disease, 2011, 1, 123-36.	1.5	44
49	Impact of dendritic spine preservation in medium spiny neurons on dopamine graft efficacy and the expression of dyskinesias in parkinsonian rats. European Journal of Neuroscience, 2010, 31, 478-490.	1.2	54
50	Focal not widespread grafts induce novel dyskinetic behavior in parkinsonian rats. Neurobiology of Disease, 2006, 21, 165-180.	2.1	93
51	Diminished Survival of Mesencephalic Dopamine Neurons Grafted into Aged Hosts Occurs during the Immediate Postgrafting Interval. Experimental Neurology, 2001, 169, 23-29.	2.0	62
52	Diminished Viability, Growth, and Behavioral Efficacy of Fetal Dopamine Neuron Grafts in Aging Rats with Long-Term Dopamine Depletion: An Argument for Neurotrophic Supplementation. Journal of Neuroscience, 1999, 19, 5563-5573.	1.7	83
53	Therapeutic Potential of Nerve Growth Factors in Parkinson??s Disease. Drugs and Aging, 1999, 14, 261-287.	1.3	74