

# Sven Ove gren

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/8413362/sven-ove-ogren-publications-by-year.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

120  
papers

5,139  
citations

44  
h-index

64  
g-index

126  
ext. papers

5,493  
ext. citations

4.6  
avg, IF

5.14  
L-index

#	Paper	IF	Citations
120	Injection of galanin into the dorsal hippocampus impairs emotional memory independent of 5-HT receptor activation. <i>Behavioural Brain Research</i> , <b>2021</b> , 405, 113178	3.4	
119	Atypical but not typical antipsychotic drugs ameliorate phencyclidine-induced emotional memory impairments in mice. <i>European Neuropsychopharmacology</i> , <b>2019</b> , 29, 616-628	1.2	5
118	Neuropeptide and Small Transmitter Coexistence: Fundamental Studies and Relevance to Mental Illness. <i>Frontiers in Neural Circuits</i> , <b>2018</b> , 12, 106	3.5	53
117	Nociceptin and the NOP receptor in aversive learning in mice. <i>European Neuropsychopharmacology</i> , <b>2017</b> , 27, 1298-1307	1.2	5
116	The fast-off hypothesis revisited: A functional kinetic study of antipsychotic antagonism of the dopamine D2 receptor. <i>European Neuropsychopharmacology</i> , <b>2016</b> , 26, 467-76	1.2	30
115	Involvement of the Striatal Medium Spiny Neurons of the Direct Pathway in the Motor Stimulant Effects of Phencyclidine. <i>International Journal of Neuropsychopharmacology</i> , <b>2016</b> , 19,	5.8	4
114	60 years of advances in neuropsychopharmacology for improving brain health, renewed hope for progress. <i>European Neuropsychopharmacology</i> , <b>2015</b> , 25, 591-8	1.2	13
113	Analysis of mechanisms for memory enhancement using novel and potent 5-HT1A receptor ligands. <i>European Neuropsychopharmacology</i> , <b>2015</b> , 25, 1314-23	1.2	5
112	Replacing SNAP-25b with SNAP-25a expression results in metabolic disease. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2015</b> , 112, E4326-35	11.5	25
111	The role of the serotonin receptor subtypes 5-HT1A and 5-HT7 and its interaction in emotional learning and memory. <i>Frontiers in Pharmacology</i> , <b>2015</b> , 6, 162	5.6	81
110	Neural Stem Cell Transplant-Induced Effect on Neurogenesis and Cognition in Alzheimer Tg2576 Mice Is Inhibited by Concomitant Treatment with Amyloid-Lowering or Cholinergic $\alpha$ 7 Nicotinic Receptor Drugs. <i>Neural Plasticity</i> , <b>2015</b> , 2015, 370432	3.3	38
109	Learning from the past and looking to the future: Emerging perspectives for improving the treatment of psychiatric disorders. <i>European Neuropsychopharmacology</i> , <b>2015</b> , 25, 599-656	1.2	86
108	Typical and atypical antipsychotics do not differ markedly in their reversibility of antagonism of the dopamine D2 receptor. <i>International Journal of Neuropsychopharmacology</i> , <b>2014</b> , 17, 149-55	5.8	15
107	GABA(A) receptor activation in the CA1 area of the dorsal hippocampus impairs consolidation of conditioned contextual fear in C57BL/6J mice. <i>Behavioural Brain Research</i> , <b>2013</b> , 238, 160-9	3.4	26
106	Classical Neurotransmitters and Neuropeptides <b>2013</b> , 1835-1841		1
105	Central 5-HT1A receptor-mediated modulation of heart rate dynamics and its adjustment by conditioned and unconditioned fear in mice. <i>British Journal of Pharmacology</i> , <b>2013</b> , 170, 859-70	8.6	8
104	Prenatal exposure to carbamazepine reduces hippocampal and cortical neuronal cell population in new-born and young mice without detectable effects on learning and memory. <i>PLoS ONE</i> , <b>2013</b> , 8, e80497	3.7	11

103	5-HT1A and 5-HT7 receptor crosstalk in the regulation of emotional memory: implications for effects of selective serotonin reuptake inhibitors. <i>Neuropharmacology</i> , <b>2012</b> , 63, 1150-60	5.5	44
102	Prenatal immune activation interacts with genetic Nurr1 deficiency in the development of attentional impairments. <i>Journal of Neuroscience</i> , <b>2012</b> , 32, 436-51	6.6	93
101	Modeling Parkinson's disease genetics: altered function of the dopamine system in Adh4 knockout mice. <i>Behavioural Brain Research</i> , <b>2011</b> , 217, 439-45	3.4	8
100	Ethanol and acetaldehyde exposure induces specific epigenetic modifications in the prodynorphin gene promoter in a human neuroblastoma cell line. <i>FASEB Journal</i> , <b>2011</b> , 25, 1069-75	0.9	33
99	Rapid increase of Nurr1 mRNA expression in limbic and cortical brain structures related to coping with depression-like behavior in mice. <i>Journal of Neuroscience Research</i> , <b>2010</b> , 88, 2284-93	4.4	26
98	The selective 5-HT(1A) receptor antagonist NAD-299 increases acetylcholine release but not extracellular glutamate levels in the frontal cortex and hippocampus of awake rat. <i>European Neuropsychopharmacology</i> , <b>2010</b> , 20, 487-500	1.2	13
97	Neuropeptides in learning and memory processes with focus on galanin. <i>European Journal of Pharmacology</i> , <b>2010</b> , 626, 9-17	5.3	66
96	Galanin, galanin receptor subtypes and depression-like behaviour. <i>Exs</i> , <b>2010</b> , 102, 163-81		22
95	Nogo receptor 1 regulates formation of lasting memories. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2009</b> , 106, 20476-81	11.5	72
94	The nociceptin system and hippocampal cognition in mice: a pharmacological and genetic analysis. <i>Brain Research</i> , <b>2009</b> , 1305 Suppl, S7-19	3.7	26
93	5-HT(1A) and NMDA receptors interact in the rat medial septum and modulate hippocampal-dependent spatial learning. <i>Hippocampus</i> , <b>2009</b> , 19, 1187-98	3.5	31
92	Assessing aversive emotional states through the heart in mice: implications for cardiovascular dysregulation in affective disorders. <i>Neuroscience and Biobehavioral Reviews</i> , <b>2009</b> , 33, 181-90	9	33
91	Neonatal infection with neurotropic influenza A virus affects working memory and expression of type III Nrg1 in adult mice. <i>Brain, Behavior, and Immunity</i> , <b>2009</b> , 23, 733-41	16.6	28
90	Bidirectional modulation of classical fear conditioning in mice by 5-HT(1A) receptor ligands with contrasting intrinsic activities. <i>Neuropharmacology</i> , <b>2009</b> , 57, 567-76	5.5	23
89	Effects of repeated treatment of phencyclidine on cognition and gene expression in C57BL/6 mice. <i>International Journal of Neuropsychopharmacology</i> , <b>2009</b> , 12, 243-55	5.8	26
88	EGb761 protects against nigrostriatal dopaminergic neurotoxicity in 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-induced Parkinsonism in mice: role of oxidative stress. <i>European Journal of Neuroscience</i> , <b>2008</b> , 28, 41-50	3.5	63
87	5-HT7 receptor stimulation by 8-OH-DPAT counteracts the impairing effect of 5-HT(1A) receptor stimulation on contextual learning in mice. <i>European Journal of Pharmacology</i> , <b>2008</b> , 596, 107-10	5.3	55
86	Blockade of 5-HT 1B receptors facilitates contextual aversive learning in mice by disinhibition of cholinergic and glutamatergic neurotransmission. <i>Neuropharmacology</i> , <b>2008</b> , 54, 1041-50	5.5	25

85	Evaluation of exploration and risk assessment in pre-weaning mice using the novel cage test. <i>Physiology and Behavior</i> , <b>2008</b> , 93, 139-47	3.5	24
84	The role of 5-HT(1A) receptors in learning and memory. <i>Behavioural Brain Research</i> , <b>2008</b> , 195, 54-77	3.4	231
83	Repeated low dose of phencyclidine administration impairs spatial learning in mice: blockade by clozapine but not by haloperidol. <i>European Neuropsychopharmacology</i> , <b>2008</b> , 18, 486-97	1.2	29
82	An ancient duplication of exon 5 in the Snap25 gene is required for complex neuronal development/function. <i>PLoS Genetics</i> , <b>2008</b> , 4, e1000278	6	58
81	Differential role of galanin receptors in the regulation of depression-like behavior and monoamine/stress-related genes at the cell body level. <i>Neuropsychopharmacology</i> , <b>2008</b> , 33, 2573-85	8.7	80
80	Effects of naltrexone and acamprosate on alcohol-induced NGFI-A expression in mouse brain. <i>Neurochemical Research</i> , <b>2008</b> , 33, 2062-9	4.6	5
79	Differential involvement of the dorsal hippocampus in passive avoidance in C57bl/6J and DBA/2J mice. <i>Hippocampus</i> , <b>2008</b> , 18, 11-9	3.5	67
78	The effects of methylmercury on motor activity are sex- and age-dependent, and modulated by genetic deletion of adenosine receptors and caffeine administration. <i>Toxicology</i> , <b>2007</b> , 241, 119-33	4.4	32
77	The neuropeptide galanin as an in vivo modulator of brain 5-HT1A receptors: possible relevance for affective disorders. <i>Physiology and Behavior</i> , <b>2007</b> , 92, 172-9	3.5	25
76	Activation of the brain 5-HT2C receptors causes hypolocomotion without anxiogenic-like cardiovascular adjustments in mice. <i>Neuropharmacology</i> , <b>2007</b> , 52, 949-57	5.5	26
75	Modification of inherent and drug-induced dopaminergic activity after exposure to benzo(alpha)pyrene. <i>NeuroToxicology</i> , <b>2007</b> , 28, 860-7	4.4	19
74	Galanin enhances and a galanin antagonist attenuates depression-like behaviour in the rat. <i>European Neuropsychopharmacology</i> , <b>2007</b> , 17, 64-9	1.2	46
73	Effects of the 5-HT1B receptor antagonist NAS-181 on extracellular levels of acetylcholine, glutamate and GABA in the frontal cortex and ventral hippocampus of awake rats: a microdialysis study. <i>European Neuropsychopharmacology</i> , <b>2007</b> , 17, 580-6	1.2	15
72	Big dynorphin, a prodynorphin-derived peptide produces NMDA receptor-mediated effects on memory, anxiolytic-like and locomotor behavior in mice. <i>Neuropsychopharmacology</i> , <b>2006</b> , 31, 1928-37	8.7	56
71	5-Hydroxytryptamine 1A receptor blockade facilitates aversive learning in mice: interactions with cholinergic and glutamatergic mechanisms. <i>Journal of Pharmacology and Experimental Therapeutics</i> , <b>2006</b> , 316, 581-91	4.7	82
70	Galanin receptor antagonists : a potential novel pharmacological treatment for mood disorders. <i>CNS Drugs</i> , <b>2006</b> , 20, 633-54	6.7	46
69	Reduced ethanol response in the alcohol-preferring RHA rats and neuropeptide mRNAs in relevant structures. <i>European Journal of Neuroscience</i> , <b>2006</b> , 23, 531-40	3.5	27
68	Analysis of the role of 5-HT1A receptors in spatial and aversive learning in the rat. <i>Neuropharmacology</i> , <b>2005</b> , 48, 830-52	5.5	73

67	Behavioural characterisation of young adult transgenic mice overexpressing galanin under the PDGF-B promoter. <i>Regulatory Peptides</i> , <b>2005</b> , 125, 67-78		40
66	Increased phencyclidine-induced hyperactivity following cortical cholinergic denervation. <i>NeuroReport</i> , <b>2005</b> , 16, 1815-9	1.7	11
65	Gene expression changes in brains of mice exposed to a maternal virus infection. <i>NeuroReport</i> , <b>2005</b> , 16, 1111-5	1.7	19
64	Dissociation of temporal dynamics of heart rate and blood pressure responses elicited by conditioned fear but not acoustic startle. <i>Behavioral Neuroscience</i> , <b>2005</b> , 119, 55-65	2.1	17
63	Distribution of galanin in the brain of a galanin-overexpressing transgenic mouse. <i>Neuropeptides</i> , <b>2005</b> , 39, 293-8	3.3	8
62	Behavioural characterisation of transgenic mice overexpressing galanin under the PDGF-B promoter. <i>Neuropeptides</i> , <b>2005</b> , 39, 299-304	3.3	33
61	Time-dependent involvement of the dorsal hippocampus in trace fear conditioning in mice. <i>Hippocampus</i> , <b>2005</b> , 15, 418-26	3.5	150
60	Modulation of neurotransmitter release and metabolism <b>2005</b> , 47-58		
59	Impeded interaction between Schwann cells and axons in the absence of laminin alpha4. <i>Journal of Neuroscience</i> , <b>2005</b> , 25, 3692-700	6.6	72
58	Corticotropin-releasing factor receptor 1 and central heart rate regulation in mice during expression of conditioned fear. <i>Journal of Pharmacology and Experimental Therapeutics</i> , <b>2005</b> , 312, 905-16	4.7	30
57	Evidence in locomotion test for the functional heterogeneity of ORL-1 receptors. <i>British Journal of Pharmacology</i> , <b>2004</b> , 141, 132-40	8.6	44
56	Chemical identity of 5-HT <sub>2A</sub> receptor immunoreactive neurons of the rat septal complex and dorsal hippocampus. <i>Brain Research</i> , <b>2004</b> , 1010, 156-65	3.7	48
55	Hypericum perforatum L (St John's wort) preferentially increases extracellular dopamine levels in the rat prefrontal cortex. <i>British Journal of Pharmacology</i> , <b>2004</b> , 142, 414-8	8.6	36
54	Enhanced hippocampal noradrenaline and serotonin release in galanin-overexpressing mice after repeated forced swimming test. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2004</b> , 101, 354-9	11.5	52
53	Central NPY receptor-mediated alteration of heart rate dynamics in mice during expression of fear conditioned to an auditory cue. <i>Regulatory Peptides</i> , <b>2004</b> , 120, 205-14		35
52	Distribution of galanin and galanin transcript in the brain of a galanin-overexpressing transgenic mouse. <i>Journal of Chemical Neuroanatomy</i> , <b>2004</b> , 28, 185-216	3.2	29
51	Disruption of EphA/ephrin-a signaling in the nigrostriatal system reduces dopaminergic innervation and dissociates behavioral responses to amphetamine and cocaine. <i>Molecular and Cellular Neurosciences</i> , <b>2004</b> , 26, 418-28	4.8	44
50	Behavioral and autonomic dynamics during contextual fear conditioning in mice. <i>Autonomic Neuroscience: Basic and Clinical</i> , <b>2004</b> , 115, 15-27	2.4	36

49	Selective 5-HT1A antagonists WAY 100635 and NAD-299 attenuate the impairment of passive avoidance caused by scopolamine in the rat. <i>Neuropsychopharmacology</i> , <b>2003</b> , 28, 253-64	8.7	81
48	Analysis of the role of the 5-HT1B receptor in spatial and aversive learning in the rat. <i>Neuropsychopharmacology</i> , <b>2003</b> , 28, 1642-55	8.7	24
47	Effects of prenatal exposure to methylmercury on dopamine-mediated locomotor activity and dopamine D2 receptor binding. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , <b>2003</b> , 367, 500-8	3.4	59
46	Decreased ethanol preference and wheel running in Nurr1-deficient mice. <i>European Journal of Neuroscience</i> , <b>2003</b> , 17, 2418-24	3.5	22
45	Galanin attenuates basal and antidepressant drug-induced increase of extracellular serotonin and noradrenaline levels in the rat hippocampus. <i>Neuroscience Letters</i> , <b>2003</b> , 339, 239-42	3.3	51
44	Local dopaminergic modulation of the motor activity induced by N-methyl-D-aspartate receptor stimulation in the ventral hippocampus. <i>Neuropsychopharmacology</i> , <b>2002</b> , 26, 737-43	8.7	17
43	Galanin is a potent in vivo modulator of mesencephalic serotonergic neurotransmission. <i>Neuropsychopharmacology</i> , <b>2002</b> , 27, 341-56	8.7	74
42	Some aspects on the anatomy and function of central cholecystinin systems. <i>Basic and Clinical Pharmacology and Toxicology</i> , <b>2002</b> , 91, 382-6		22
41	Facilitation of dopamine-mediated locomotor activity in adult rats following cholinergic denervation. <i>Experimental Neurology</i> , <b>2002</b> , 174, 96-108	5.7	16
40	Prolonged effects of intraventricular galanin on a 5-hydroxytryptamine(1A) receptor mediated function in the rat. <i>Neuroscience Letters</i> , <b>2001</b> , 299, 145-9	3.3	32
39	Adenosine/dopamine interaction: implications for the treatment of Parkinson's disease. <i>Parkinsonism and Related Disorders</i> , <b>2001</b> , 7, 235-241	3.6	102
38	Intraventricular galanin produces a time-dependent modulation of 5-HT1A receptors in the dorsal raphe of the rat. <i>NeuroReport</i> , <b>2000</b> , 11, 3943-8	1.7	32
37	Ontogeny of the motor inhibitory role of dopamine D(3) receptor subtype in rats. <i>European Journal of Pharmacology</i> , <b>2000</b> , 392, 35-9	5.3	6
36	Involvement of the 5-HT1A receptors in classical fear conditioning in C57BL/6J mice. <i>Journal of Neuroscience</i> , <b>2000</b> , 20, 8515-27	6.6	94
35	Decreased 5-HT transporter mRNA in neurons of the dorsal raphe nucleus and behavioral depression in the obese leptin-deficient ob/ob mouse. <i>Molecular Brain Research</i> , <b>2000</b> , 81, 51-61		101
34	Galanin and spatial learning in the rat. Evidence for a differential role for galanin in subregions of the hippocampal formation. <i>Neuropharmacology</i> , <b>2000</b> , 39, 1386-403	5.5	38
33	Galanin and learning. <i>Brain Research</i> , <b>1999</b> , 848, 174-82	3.7	37
32	Effect of N-methyl-D-aspartate on motor activity and in vivo adenosine striatal outflow in the rat. <i>European Journal of Pharmacology</i> , <b>1999</b> , 385, 15-9	5.3	30

31	A behavioral analysis of the spatial learning deficit induced by the NMDA receptor antagonist MK-801 (dizocilpine) in the rat. <i>Neuropsychopharmacology</i> , <b>1999</b> , 21, 414-26	8.7	72
30	Analysis of the 5-HT1A receptor involvement in passive avoidance in the rat. <i>British Journal of Pharmacology</i> , <b>1998</b> , 125, 499-509	8.6	44
29	Modulation of a 5-HT1A receptor-mediated behavioral response by the neuropeptide galanin. <i>Annals of the New York Academy of Sciences</i> , <b>1998</b> , 863, 442-4	6.5	2
28	Time-dependent effects of intrahippocampal galanin on spatial learning. Relationship to distribution and kinetics. <i>Annals of the New York Academy of Sciences</i> , <b>1998</b> , 863, 454-6	6.5	10
27	Intraventricular galanin modulates a 5-HT1A receptor-mediated behavioural response in the rat. <i>European Journal of Neuroscience</i> , <b>1998</b> , 10, 1230-40	3.5	41
26	Simultaneous determination of acetylcholine, choline and physostigmine in microdialysis samples from rat hippocampus by microbore liquid chromatography/electrochemistry on peroxidase redox polymer coated electrodes. <i>Journal of Neuroscience Methods</i> , <b>1998</b> , 83, 143-50	3	40
25	"Atypical" neuromodulatory profile of glutapyrone, a representative of a novel class of amino acid-containing dipeptide-mimicking 1,4-dihydropyridine (DHP) compounds: in vitro and in vivo studies. <i>European Neuropsychopharmacology</i> , <b>1998</b> , 8, 329-47	1.2	29
24	Dopamine D1 and D2 receptor-mediated acute and long-lasting behavioral effects of glial cell line-derived neurotrophic factor administered into the striatum. <i>Experimental Neurology</i> , <b>1998</b> , 154, 302-14	5.7	19
23	Differential effects of selective adenosine A1 and A2A receptor agonists on dopamine receptor agonist-induced behavioural responses in rats. <i>European Journal of Pharmacology</i> , <b>1998</b> , 347, 153-8	5.3	63
22	Dopamine receptor antagonists block nerve growth factor-induced hyperactivity. <i>European Journal of Pharmacology</i> , <b>1997</b> , 326, 1-5	5.3	7
21	Prolonged treatment with haloperidol and clozapine in the rat: differential effects on spontaneous and theophylline-induced motor activity. <i>Neuroscience Letters</i> , <b>1997</b> , 232, 21-4	3.3	1
20	Adenosine A2A agonists: a potential new type of atypical antipsychotic. <i>Neuropsychopharmacology</i> , <b>1997</b> , 17, 82-91	8.7	123
19	Dopamine D1 receptor-mediated facilitation of GABAergic neurotransmission in the rat strioentopenduncular pathway and its modulation by adenosine A1 receptor-mediated mechanisms. <i>European Journal of Neuroscience</i> , <b>1996</b> , 8, 1545-53	3.5	122
18	The Behavioural Pharmacology of Typical and Atypical Antipsychotic Drugs. <i>Handbook of Experimental Pharmacology</i> , <b>1996</b> , 225-266	3.2	18
17	Chronic haloperidol treatment leads to an increase in the intramembrane interaction between adenosine A2 and dopamine D2 receptors in the neostriatum. <i>Psychopharmacology</i> , <b>1994</b> , 116, 279-84	4.7	28
16	Effects of typical and atypical antipsychotic drugs on two-way active avoidance. Relationship to DA receptor blocking profile. <i>Psychopharmacology</i> , <b>1994</b> , 114, 383-91	4.7	46
15	Phencyclidine- and dizocilpine-induced hyperlocomotion are differentially mediated. <i>Neuropsychopharmacology</i> , <b>1994</b> , 11, 167-77	8.7	94
14	Differential effects of the putative galanin receptor antagonists M15 and M35 on striatal acetylcholine release. <i>European Journal of Pharmacology</i> , <b>1993</b> , 242, 59-64	5.3	41

13	Galanin stimulates striatal acetylcholine release via a mechanism unrelated to cholinergic receptor stimulation. <i>Regulatory Peptides</i> , <b>1993</b> , 45, 353-62		22
12	Galanin-evoked acetylcholine release in the rat striatum is blocked by the putative galanin antagonist M15. <i>Brain Research</i> , <b>1992</b> , 574, 317-9	3.7	27
11	Galanin stimulates acetylcholine release in the rat striatum. <i>Neuroscience Letters</i> , <b>1991</b> , 128, 253-6	3.3	32
10	Asphyctic lesion: proliferation of tyrosine hydroxylase-immunoreactive nerve cell bodies in the rat substantia nigra and functional changes in dopamine neurotransmission. <i>Brain Research</i> , <b>1991</b> , 543, 1-9	3.7	168
9	Galanin: Regulation of Cholinergic Function and Behaviour <b>1991</b> , 193-199		6
8	Potential antipsychotic agents. Part 8. Antidopaminergic properties of a potent series of 5-substituted (1S)-N-[(1-ethylpyrrolidin-2-yl)methyl]-2,3-dimethoxybenzamide. Synthesis via common lithio intermediates. <i>Helvetica Chimica Acta</i> , <b>1990</b> , 73, 417-425	2	17
7	The effects of p-chloroamphetamine, a depletor of brain serotonin, on the performance of rats in two types of positively reinforced complex spatial discrimination tasks. <i>Behavioral and Neural Biology</i> , <b>1989</b> , 52, 131-44		44
6	D1- and D2-receptor antagonists induce catalepsy via different efferent striatal pathways [corrected]. <i>Neuroscience Letters</i> , <b>1988</b> , 85, 333-8	3.3	64
5	The selective dopamine D2 receptor antagonist raclopride discriminates between dopamine-mediated motor functions. <i>Psychopharmacology</i> , <b>1986</b> , 90, 287-94	4.7	187
4	Serotonin receptor involvement in the avoidance learning deficit caused by p-chloroamphetamine-induced serotonin release. <i>Acta Physiologica Scandinavica</i> , <b>1986</b> , 126, 449-62		19
3	Central noradrenaline depletion attenuates amphetamine-induced locomotor behavior. <i>Neuroscience Letters</i> , <b>1986</b> , 64, 139-44	3.3	25
2	Role of serotonin in memory: facilitation by alaproclate and zimeldine. <i>Psychopharmacology</i> , <b>1984</b> , 84, 496-502	4.7	56
1	Inhibitors of neuronal monoamine uptake. 2. Selective inhibition of 5-hydroxytryptamine uptake by alpha-amino acid esters of phenethyl alcohols. <i>Journal of Medicinal Chemistry</i> , <b>1978</b> , 21, 448-56	8.3	86