

Elena Succurro

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8415805/elena-succurro-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

99
papers

2,237
citations

25
h-index

44
g-index

106
ext. papers

2,649
ext. citations

5.7
avg, IF

4.6
L-index

#	Paper	IF	Citations
99	Follow-up of women with a history of gestational diabetes in Italy: Are we missing an opportunity for primary prevention of type 2 diabetes and cardiovascular disease?. <i>Diabetes/Metabolism Research and Reviews</i> , 2021 , 37, e3411	7.5	
98	The Functional Polymorphism of rs9267551 Is an Independent Determinant of Arterial Stiffness.. <i>Frontiers in Cardiovascular Medicine</i> , 2021 , 8, 811431	5.4	
97	Pharmacological treatment of type 2 diabetes in elderly patients with heart failure: randomized trials and beyond. <i>Heart Failure Reviews</i> , 2021 , 1	5	3
96	Role of Vitamin D in Cardiovascular Diseases. <i>Endocrines</i> , 2021 , 2, 417-426	0.8	1
95	Sex-specific differences in left ventricular mass and myocardial energetic efficiency in non-diabetic, pre-diabetic and newly diagnosed type 2 diabetic subjects. <i>Cardiovascular Diabetology</i> , 2021 , 20, 60	8.7	3
94	Screening of postpartum diabetes in women with gestational diabetes: high-risk subgroups and areas for improvements-the STRONG observational study. <i>Acta Diabetologica</i> , 2021 , 58, 1187-1197	3.9	4
93	Hyperglycemia at 1h-OGTT in Pregnancy: A Reliable Predictor of Metabolic Outcomes?. <i>Frontiers in Endocrinology</i> , 2021 , 12, 612829	5.7	1
92	Nox2 up-regulation and hypoalbuminemia in patients with type 2 diabetes mellitus. <i>Free Radical Biology and Medicine</i> , 2021 , 168, 1-5	7.8	0
91	The TRIB3 R84 variant is associated with increased left ventricular mass in a sample of 2426 White individuals. <i>Cardiovascular Diabetology</i> , 2021 , 20, 115	8.7	0
90	Impaired Clinical Efficacy of Aspirin in Hypoalbuminemic Patients With Diabetes Mellitus. <i>Frontiers in Pharmacology</i> , 2021 , 12, 695961	5.6	1
89	Neutrophil degranulation biomarkers characterize restrictive echocardiographic pattern with diastolic dysfunction in patients with diabetes. <i>European Journal of Clinical Investigation</i> , 2021 , 51, e13640 ⁶	4.6	0
88	Depressed myocardial mechano-energetic efficiency in subjects with dysglycemia. <i>Diabetes Research and Clinical Practice</i> , 2021 , 177, 108883	7.4	0
87	Nonalcoholic fatty liver disease is associated with a decreased myocardial mechano-energetic efficiency. <i>Journal of Internal Medicine</i> , 2021 , 289, 221-231	10.8	5
86	The multifaceted spectrum of liver cirrhosis in older hospitalised patients: analysis of the REPOSI registry. <i>Age and Ageing</i> , 2021 , 50, 498-504	3	0
85	Effects of Intermittent Pneumatic Compression on Lower Limb Lymphedema in Patients with Type 2 Diabetes Mellitus: A Pilot Randomized Controlled Trial. <i>Medicina (Lithuania)</i> , 2021 , 57,	3.1	2
84	Why glycated albumin decreases in pregnancy? Evidences from a prospective study on physiological pregnancies of Caucasian women. <i>Clinica Chimica Acta</i> , 2021 , 520, 217-218	6.2	1
83	Non-alcoholic fatty liver disease is associated with cardiovascular disease in subjects with different glucose tolerance. <i>Diabetes/Metabolism Research and Reviews</i> , 2020 , 36, e3333	7.5	9

82	Reduction in Global Myocardial Glucose Metabolism in Subjects With 1-Hour Postload Hyperglycemia and Impaired Glucose Tolerance. <i>Diabetes Care</i> , 2020 , 43, 669-676	14.6	7
81	Association between Serum Mg Concentrations and Cardiovascular Organ Damage in a Cohort of Adult Subjects. <i>Nutrients</i> , 2020 , 12,	6.7	2
80	Italian recommendations for the diagnosis of gestational diabetes during COVID-19 pandemic: Position statement of the Italian Association of Clinical Diabetologists (AMD) and the Italian Diabetes Society (SID), diabetes, and pregnancy study group. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2020 , 30, 1418-1422	4.5	9
79	Different Patterns of Left Ventricular Hypertrophy in Metabolically Healthy and Insulin-Resistant Obese Subjects. <i>Nutrients</i> , 2020 , 12,	6.7	8
78	HDL cholesterol is an independent predictor of ß-cell function decline and incident type 2 diabetes: A longitudinal study. <i>Diabetes/Metabolism Research and Reviews</i> , 2020 , 36, e3289	7.5	9
77	Awareness about diabetes and pregnancy among diabetes specialists and fellows: The YoSID diabetes and pregnancy project. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2020 , 30, 1520-1524	4.5	2
76	Relative Risk of Cardiovascular Disease Is Higher in Women With Type 2 Diabetes, but Not in Those With Prediabetes, as Compared With Men. <i>Diabetes Care</i> , 2020 , 43, 3070-3078	14.6	5
75	Recommendations and management of hyperglycaemia in pregnancy during COVID-19 pandemic in Italy. <i>Diabetes Research and Clinical Practice</i> , 2020 , 166, 108345	7.4	9
74	Patterns of infections in older patients acutely admitted to medical wards: data from the REPOSI register. <i>Internal and Emergency Medicine</i> , 2019 , 14, 1347-1352	3.7	
73	Response to Letter to the Editor: "One-Hour Postload Hyperglycemia: Implications for Prediction and Prevention of Type 2 Diabetes". <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019 , 104, 676-677	5.6	1
72	Individuals With Prediabetes Display Different Age-Related Pathophysiological Characteristics. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019 , 104, 2911-2924	5.6	8
71	Hospital Care of Older Patients With COPD: Adherence to International Guidelines for Use of Inhaled Bronchodilators and Corticosteroids. <i>Journal of the American Medical Directors Association</i> , 2019 , 20, 1313-1317.e9	5.9	4
70	Elevated 1-h post-load plasma glucose is associated with right ventricular morphofunctional parameters in hypertensive patients. <i>Endocrine</i> , 2019 , 64, 525-535	4	2
69	Guidelines for the screening and diagnosis of gestational diabetes in Italy from 2010 to 2019: critical issues and the potential for improvement. <i>Acta Diabetologica</i> , 2019 , 56, 1159-1167	3.9	6
68	Prevalence of use and appropriateness of antidepressants prescription in acutely hospitalized elderly patients. <i>European Journal of Internal Medicine</i> , 2019 , 68, e7-e11	3.9	1
67	Are Circulating Mg Levels Associated with Glucose Tolerance Profiles and Incident Type 2 Diabetes?. <i>Nutrients</i> , 2019 , 11,	6.7	9
66	"MySweetGestation": A novel smartphone application for women with or at risk of diabetes during pregnancy. <i>Diabetes Research and Clinical Practice</i> , 2019 , 158, 107896	7.4	1
65	Relationships of surrogate indexes of insulin resistance with insulin sensitivity assessed by euglycemic hyperinsulinemic clamp and subclinical vascular damage. <i>BMJ Open Diabetes Research and Care</i> , 2019 , 7, e000911	4.5	33

64	One-hour post-load hyperglycemia combined with HbA1c identifies individuals with higher risk of cardiovascular diseases: Cross-sectional data from the CATAMERI study. <i>Diabetes/Metabolism Research and Reviews</i> , 2019 , 35, e3096	7.5	16
63	Major adverse cardiovascular events in non-valvular atrial fibrillation with chronic obstructive pulmonary disease: the ARAPACIS study. <i>Internal and Emergency Medicine</i> , 2018 , 13, 651-660	3.7	9
62	No effect on the short-term of a decrease in blood viscosity on insulin resistance. <i>Clinical Hemorheology and Microcirculation</i> , 2018 , 68, 45-50	2.5	3
61	Serum IgG2 levels are specifically associated with whole-body insulin-mediated glucose disposal in non-diabetic offspring of type 2 diabetic individuals: a cross-sectional study. <i>Scientific Reports</i> , 2018 , 8, 13616	4.9	5
60	One-Hour Postload Hyperglycemia: Implications for Prediction and Prevention of Type 2 Diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018 , 103, 3131-3143	5.6	23
59	Higher serum levels of uric acid are associated with a reduced insulin clearance in non-diabetic individuals. <i>Acta Diabetologica</i> , 2018 , 55, 835-842	3.9	17
58	Hemorheological alterations in adults with prediabetes identified by hemoglobin A1c levels. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2017 , 27, 601-608	4.5	9
57	Prevalence and Determinants of the Use of Lipid-Lowering Agents in a Population of Older Hospitalized Patients: the Findings from the REPOSI (REgistro POLiterapie Societ�Italiana di Medicina Interna) Study. <i>Drugs and Aging</i> , 2017 , 34, 311-319	4.7	7
56	Association between hemoglobin glycation index and hepatic steatosis in non-diabetic individuals. <i>Diabetes Research and Clinical Practice</i> , 2017 , 134, 53-61	7.4	17
55	Association between hemoglobin glycation index with insulin resistance and carotid atherosclerosis in non-diabetic individuals. <i>PLoS ONE</i> , 2017 , 12, e0175547	3.7	28
54	Liraglutide improves cardiac function in patients with type 2 diabetes and chronic heart failure. <i>Endocrine</i> , 2017 , 57, 464-473	4	41
53	Elevated hemoglobin glycation index identify non-diabetic individuals at increased risk of kidney dysfunction. <i>Oncotarget</i> , 2017 , 8, 79576-79586	3.3	10
52	Loss of Eyebrows and Eyelashes During Concomitant Treatment with Sitagliptin and Metformin. <i>Current Drug Safety</i> , 2017 , 12, 10-12	1.4	3
51	Adherence to antithrombotic therapy guidelines improves mortality among elderly patients with atrial fibrillation: insights from the REPOSI study. <i>Clinical Research in Cardiology</i> , 2016 , 105, 912-920	6.1	47
50	Response to the Letter: Comment to the letter by Marathe CS, Rayne CK, Jones KL, Horowitz M. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2016 , 101, L35	5.6	
49	Experiences on quantitative cardiac PET analysis 2016 ,		4
48	Frequency of 1-h post-load glucose ≥ 55 mg/dl among individuals with different glucose tolerance conditions. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2016 , 26, 439-41	4.5	6
47	Potential effects of current drug therapies on cognitive impairment in patients with type 2 diabetes. <i>Frontiers in Neuroendocrinology</i> , 2016 , 42, 76-92	8.9	29

46	One-Hour Postload Hyperglycemia Is a Stronger Predictor of Type 2 Diabetes Than Impaired Fasting Glucose. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015 , 100, 3744-51	5.6	78
45	Obese Patients With a Binge Eating Disorder Have an Unfavorable Metabolic and Inflammatory Profile. <i>Medicine (United States)</i> , 2015 , 94, e2098	1.8	60
44	Bilateral lower limbs edema with "wooden" character induced by insulin glargine treatment. <i>Acta Diabetologica</i> , 2015 , 52, 809-11	3.9	5
43	Rosacea-like facial rash related to metformin administration in a young woman. <i>BMC Pharmacology & Toxicology</i> , 2014 , 15, 3	2.6	8
42	Differences in insulin clearance between metabolically healthy and unhealthy obese subjects. <i>Acta Diabetologica</i> , 2014 , 51, 257-61	3.9	37
41	Elevated 1-h post-load plasma glucose levels in subjects with normal glucose tolerance are associated with unfavorable inflammatory profile. <i>Acta Diabetologica</i> , 2014 , 51, 927-32	3.9	26
40	Dietary patterns and 1-h post-load glucose in essential hypertension. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2014 , 24, 547-53	4.5	4
39	Association between noninvasive fibrosis markers and cardio-vascular organ damage among adults with hepatic steatosis. <i>PLoS ONE</i> , 2014 , 9, e104941	3.7	23
38	Multimorbidity and polypharmacy in the elderly: lessons from REPOSI. <i>Internal and Emergency Medicine</i> , 2014 , 9, 723-34	3.7	95
37	Insulin sensitivity, and β cell function in relation to hemoglobin A1C. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2014 , 24, 27-33	4.5	6
36	Increased carotid intima-media thickness in the physiologic range is associated with impaired postprandial glucose metabolism, insulin resistance and beta cell dysfunction. <i>Atherosclerosis</i> , 2013 , 229, 277-81	3.1	15
35	Insulin clearance is associated with carotid artery intima-media thickness. <i>Atherosclerosis</i> , 2013 , 229, 453-8	3.1	7
34	Nonalpine thyroid angiosarcoma in a patient with hashimoto thyroiditis. <i>Case Reports in Oncological Medicine</i> , 2013 , 2013, 901246	0.9	9
33	Reciprocal association of plasma IGF-1 and interleukin-6 levels with cardiometabolic risk factors in nondiabetic subjects. <i>Diabetes Care</i> 2008;31:1886-1888. <i>Diabetes Care</i> , 2013 , 36, 183-183	14.6	78
32	Decreased insulin clearance in individuals with elevated 1-h post-load plasma glucose levels. <i>PLoS ONE</i> , 2013 , 8, e77440	3.7	21
31	Cardiometabolic risk profiles and carotid atherosclerosis in individuals with prediabetes identified by fasting glucose, postchallenge glucose, and hemoglobin A1c criteria. <i>Diabetes Care</i> , 2012 , 35, 1144-9	14.6	67
30	Comparison of A1C, fasting and 2-h post-load plasma glucose criteria to diagnose diabetes in Italian Caucasians. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2012 , 22, 561-6	4.5	14
29	A functional variant of the dimethylarginine dimethylaminohydrolase-2 gene is associated with insulin sensitivity. <i>PLoS ONE</i> , 2012 , 7, e36224	3.7	16

28	Insulin sensitivity, β cell function, and incretin effect in individuals with elevated 1-hour postload plasma glucose levels. <i>Diabetes Care</i> , 2012 , 35, 868-72	14.6	58
27	IGF-1 levels link estimated glomerular filtration rate to insulin resistance in obesity: a study in obese, but metabolically healthy, subjects and obese, insulin-resistant subjects. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2011 , 21, 933-40	4.5	25
26	One-hour post-load plasma glucose levels are associated with elevated liver enzymes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2011 , 21, 713-8	4.5	25
25	The type 2 diabetes and insulin-resistance locus near IRS1 is a determinant of HDL cholesterol and triglycerides levels among diabetic subjects. <i>Atherosclerosis</i> , 2011 , 216, 157-60	3.1	22
24	Low insulin-like growth factor-1 levels are associated with anaemia in adult non-diabetic subjects. <i>Thrombosis and Haemostasis</i> , 2011 , 105, 365-70	7	20
23	Usefulness of hemoglobin A1c as a criterion to define the metabolic syndrome in a cohort of italian nondiabetic white subjects. <i>American Journal of Cardiology</i> , 2011 , 107, 1650-5	3	22
22	Nonalcoholic fatty liver disease is associated with low circulating levels of insulin-like growth factor-I. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2011 , 96, E1640-4	5.6	70
21	The TRIB3 R84 variant is associated with increased carotid intima-media thickness in vivo and with enhanced MAPK signalling in human endothelial cells. <i>Cardiovascular Research</i> , 2011 , 89, 184-92	9.9	23
20	One-hour postload plasma glucose levels and left ventricular mass in hypertensive patients. <i>Diabetes Care</i> , 2011 , 34, 1406-11	14.6	68
19	GRB10 gene and type 2 diabetes in Whites. <i>Journal of Internal Medicine</i> , 2010 , 267, 132-3	10.8	6
18	One-hour postload plasma glucose levels are associated with kidney dysfunction. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2010 , 5, 1922-7	6.9	61
17	Positive association between plasma IGF1 and high-density lipoprotein cholesterol levels in adult nondiabetic subjects. <i>European Journal of Endocrinology</i> , 2010 , 163, 75-80	6.5	23
16	Sudden progression from impaired glucose tolerance to type 2 diabetes after discontinuation of administration of anti-tumor necrosis factor-alpha antibody infliximab. <i>International Journal of Immunopathology and Pharmacology</i> , 2010 , 23, 961-3	3	11
15	An empirical index of insulin sensitivity from short IVGTT: validation against the minimal model and glucose clamp indices in patients with different clinical characteristics. <i>Diabetologia</i> , 2010 , 53, 144-52	10.3	61
14	TRIB3 R84 variant affects glucose homeostasis by altering the interplay between insulin sensitivity and secretion. <i>Diabetologia</i> , 2010 , 53, 1354-61	10.3	14
13	Acute rhabdomyolysis during treatment with amisulpride and metformin. <i>European Journal of Clinical Pharmacology</i> , 2010 , 66, 321-2	2.8	4
12	Differences in cardiovascular risk profile based on relationship between post-load plasma glucose and fasting plasma levels. <i>Diabetes/Metabolism Research and Reviews</i> , 2009 , 25, 351-6	7.5	6
11	IRS1 G972R polymorphism and type 2 diabetes: a paradigm for the difficult ascertainment of the contribution to disease susceptibility of low-frequency-low-risk variants. <i>Diabetologia</i> , 2009 , 52, 1852-7 ^{10.3}	10.3	30

10	IL-18 gene polymorphism and metabolic syndrome. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2009 , 19, e5-6	4.5	21
9	Low plasma insulin-like growth factor-1 levels are associated with reduced insulin sensitivity and increased insulin secretion in nondiabetic subjects. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2009 , 19, 713-9	4.5	41
8	Elevated one-hour post-load plasma glucose levels identifies subjects with normal glucose tolerance but early carotid atherosclerosis. <i>Atherosclerosis</i> , 2009 , 207, 245-9	3.1	120
7	Insulin secretion in metabolically obese, but normal weight, and in metabolically healthy but obese individuals. <i>Obesity</i> , 2008 , 16, 1881-6	8	110
6	Reciprocal association of plasma IGF-1 and interleukin-6 levels with cardiometabolic risk factors in nondiabetic subjects. <i>Diabetes Care</i> , 2008 , 31, 1886-8	14.6	43
5	Single-nucleotide polymorphism rs7754840 of CDKAL1 is associated with impaired insulin secretion in nondiabetic offspring of type 2 diabetic subjects and in a large sample of men with normal glucose tolerance. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008 , 93, 1924-30	5.6	71
4	Variance of the SGK1 gene is associated with insulin secretion in different European populations: results from the TUEF, EUGENE2, and METSIM studies. <i>PLoS ONE</i> , 2008 , 3, e3506	3.7	13
3	Metabolic and cardiovascular risk factors in subjects with impaired fasting glucose: the 100 versus 110 mg/dL threshold. <i>Diabetes/Metabolism Research and Reviews</i> , 2007 , 23, 547-50	7.5	21
2	Metabolically healthy but obese women have an intermediate cardiovascular risk profile between healthy nonobese women and obese insulin-resistant women. <i>Diabetes Care</i> , 2007 , 30, 2145-7	14.6	122
1	Plasma concentration of IGF-I is independently associated with insulin sensitivity in subjects with different degrees of glucose tolerance. <i>Diabetes Care</i> , 2005 , 28, 120-5	14.6	135