

Lina Wang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8425013/publications.pdf>

Version: 2024-02-01

11
papers

129
citations

1307543

7
h-index

1372553

10
g-index

12
all docs

12
docs citations

12
times ranked

156
citing authors

#	ARTICLE	IF	CITATIONS
1	Diagnostic value of dysregulated microribonucleic acids in the placenta and circulating exosomes in gestational diabetes mellitus. <i>Journal of Diabetes Investigation</i> , 2021, 12, 1490-1500.	2.4	24
2	Long noncoding RNA LINC00461 induced osteoarthritis progression by inhibiting miR-30a-5p. <i>Aging</i> , 2020, 12, 4111-4123.	3.1	19
3	Circulating exosomal hsa_circRNA_0039480 is highly expressed in gestational diabetes mellitus and may be served as a biomarker for early diagnosis of GDM. <i>Journal of Translational Medicine</i> , 2022, 20, 5.	4.4	17
4	Human chorionic gonadotropin and IL-35 contribute to the maintenance of peripheral immune tolerance during pregnancy through mediating the generation of IL-10+ or IL-35+ Breg cells. <i>Experimental Cell Research</i> , 2019, 383, 111513.	2.6	16
5	MiR-370 inhibits the oxidative stress and apoptosis of cardiac myocytes induced by hydrogen peroxide by targeting FOXO1. <i>Experimental and Therapeutic Medicine</i> , 2019, 18, 3025-3031.	1.8	14
6	Long non-coding RNA LOC283070 mediates the transition of LNCaP cells into androgen-independent cells possibly via CAMK1D. <i>American Journal of Translational Research (discontinued)</i> , 2016, 8, 5219-5234.	0.0	12
7	miR-137 suppresses cell growth and extracellular matrix degradation through regulating ADAMTS-5 in chondrocytes. <i>American Journal of Translational Research (discontinued)</i> , 2019, 11, 7027-7034.	0.0	8
8	Relationship between SIRT1 gene and adolescent depressive disorder with nonsuicidal self-injury behavior. <i>Medicine (United States)</i> , 2021, 100, e26747.	1.0	7
9	Shorter Leukocyte Telomere Length coupled with lower expression of Telomerase Genes in patients with Essential Hypertension. <i>International Journal of Medical Sciences</i> , 2020, 17, 2180-2186.	2.5	6
10	A pre-ablative thyroid-stimulating hormone with 30-70 mIU/L achieves better response to initial radioiodine remnant ablation in differentiated thyroid carcinoma patients. <i>Scientific Reports</i> , 2021, 11, 1348.	3.3	6
11	Patients with benign prostatic hyperplasia show shorter leukocyte telomere length but no association with telomerase gene polymorphisms in Han Chinese males. <i>International Journal of Clinical and Experimental Pathology</i> , 2020, 13, 2123-2129.	0.5	0