

# Shanqun Jiang

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8427069/publications.pdf>

Version: 2024-02-01

37  
papers

420  
citations

687363

13  
h-index

794594

19  
g-index

37  
all docs

37  
docs citations

37  
times ranked

608  
citing authors

#	ARTICLE	IF	CITATIONS
1	Associations of Two Common Polymorphisms in MTHFR Gene with Blood Lipids and Therapeutic Efficacy of Simvastatin. <i>Current Pharmaceutical Design</i> , 2022, 28, .	1.9	1
2	Effect of <i>ABCG1</i> gene DNA methylations on the lipid-lowering efficacy of simvastatin. <i>Pharmacogenomics</i> , 2021, 22, 27-39.	1.3	1
3	The effect of ABCA1 gene DNA methylation on blood pressure levels in a Chinese hyperlipidemic population. <i>Journal of Human Hypertension</i> , 2021, 35, 1139-1148.	2.2	2
4	<i>PNPT1</i> and <i>PCGF3</i> variants associated with angiotensin-converting enzyme inhibitor-induced cough: a nested case-control genome-wide study. <i>Pharmacogenomics</i> , 2020, 21, 601-614.	1.3	5
5	PHOSPHO1 Gene DNA Methylations are Associated with a Change in HDL-C Response to Simvastatin Treatment. <i>Current Pharmaceutical Design</i> , 2020, 26, 4944-4952.	1.9	1
6	TNFAIP3 genetic polymorphisms reduce ankylosing spondylitis risk in Eastern Chinese Han population. <i>Scientific Reports</i> , 2019, 9, 10209.	3.3	4
7	Serum Sclerostin and Bone Morphogenetic Protein-2 Levels in Patients with Ankylosing Spondylitis: A Meta-Analysis. <i>Calcified Tissue International</i> , 2019, 105, 37-50.	3.1	13
8	Genetic polymorphisms of G protein-coupled receptor 65 gene are associated with ankylosing spondylitis in a Chinese Han population: A case-control study. <i>Human Immunology</i> , 2019, 80, 146-150.	2.4	5
9	Associations of methylenetetrahydrofolate reductase C677T genotype with blood pressure levels in Chinese population with essential hypertension. <i>Clinical and Experimental Hypertension</i> , 2018, 40, 207-212.	1.3	7
10	<i>MTHFR</i> Gene and Serum Folate Interaction on Serum Homocysteine Lowering. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2018, 38, 679-685.	2.4	29
11	Associations of the <i>ABCA1</i> and <i>LPL</i> Gene Polymorphisms With Lipid Levels in a Hyperlipidemic Population. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2018, 24, 771-779.	1.7	5
12	Changes and clinical significance of CD8+CD122+ T cells in the peripheral blood of patients with ankylosing spondylitis. <i>Clinical Rheumatology</i> , 2018, 37, 639-646.	2.2	7
13	A gender-specific association of the polymorphism Ile197Met in the kininogen 1 gene with plasma irbesartan concentrations in Chinese patients with essential hypertension. <i>Journal of Human Hypertension</i> , 2018, 32, 781-788.	2.2	1
14	Associations of the <i>SLCO1B1</i> Polymorphisms With Hepatic Function, Baseline Lipid Levels, and Lipid-lowering Response to Simvastatin in Patients With Hyperlipidemia. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2018, 24, 240S-247S.	1.7	10
15	Multiple independent structural dynamic events in the evolution of snake mitochondrial genomes. <i>BMC Genomics</i> , 2018, 19, 354.	2.8	16
16	Associations of Estrogen Receptor Alpha Gene Polymorphisms with Type 2 Diabetes Mellitus and Metabolic Syndrome: A Systematic Review and Meta-Analysis. <i>Hormone and Metabolic Research</i> , 2018, 50, 469-477.	1.5	17
17	How to Control Asthma with Personalized Management: Where Do We Stand Now?. <i>Current Drug Metabolism</i> , 2018, 19, 1188-1198.	1.2	3
18	Individual and Joint Associations of Methylenetetrahydrofolate Reductase C677T Genotype and Plasma Homocysteine With Dyslipidemia in a Chinese Population With Hypertension. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2017, 23, 287-293.	1.7	10

#	ARTICLE	IF	CITATIONS
19	Effects of LEP G2548A and LEPR Q223R Polymorphisms on Serum Lipids and Response to Simvastatin Treatment in Chinese Patients With Primary Hyperlipidemia. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2017, 23, 336-344.	1.7	4
20	Methylenetetrahydrofolate reductase C677T polymorphism, hypertension and risk of stroke: a prospective, nested case-control study. <i>International Journal of Neuroscience</i> , 2017, 127, 253-260.	1.6	21
21	Association of peripheral differential leukocyte counts with dyslipidemia risk in Chinese patients with hypertension: insight from the China Stroke Primary Prevention Trial. <i>Journal of Lipid Research</i> , 2017, 58, 256-266.	4.2	21
22	Lipid levels and new-onset diabetes in a hypertensive population: the China Stroke Primary Prevention Trial. <i>Scientific Reports</i> , 2017, 7, 7014.	3.3	4
23	Serum levels of leptin, adiponectin and resistin in patients with ankylosing spondylitis: A systematic review and meta-analysis. <i>International Immunopharmacology</i> , 2017, 52, 310-317.	3.8	15
24	Effect modification by region in the associations of LEP G2548A and LEPR Q223R polymorphisms with statin-induced CK elevation. <i>Oncotarget</i> , 2017, 8, 107565-107576.	1.8	1
25	Elevation in Total Homocysteine Levels in Chinese Patients With Essential Hypertension Treated With Antihypertensive Benazepril. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2016, 22, 191-198.	1.7	5
26	Interactive Effect of the KCNJ11 Ile337Val Polymorphism and Cigarette Smoking on the Antihypertensive Response to Irbesartan in Chinese Hypertensive Patients. <i>American Journal of Hypertension</i> , 2016, 29, 553-559.	2.0	1
27	Predicting Hyperhomocysteinemia by Methylenetetrahydrofolate Reductase C677T Polymorphism in Chinese Patients With Hypertension. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2015, 21, 661-666.	1.7	11
28	H-type hypertension and risk of stroke in chinese adults: A prospective, nested case-control study. <i>Journal of Translational Internal Medicine</i> , 2015, 3, 171-178.	2.5	59
29	Associations of MTHFR and MTRR Polymorphisms With Serum Lipid Levels in Chinese Hypertensive Patients. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2014, 20, 400-410.	1.7	22
30	Effect of Simvastatin on Plasma Homocysteine Levels and Its Modification by MTHFR C677T Polymorphism in Chinese Patients with Primary Hyperlipidemia. <i>Cardiovascular Therapeutics</i> , 2013, 31, e27-33.	2.5	22
31	Effect of simvastatin on plasma homocysteine levels and its modification by MTHFR C677T polymorphism in Chinese patients with primary hyperlipidemia. <i>Cardiovascular Therapeutics</i> , 2012, 31, n/a-n/a.	2.5	1
32	Interactive effect of angiotensin II type 1 receptor (AGT1R) polymorphisms and plasma irbesartan concentration on antihypertensive therapeutic responses to irbesartan. <i>Journal of Hypertension</i> , 2011, 29, 890-895.	0.5	11
33	Individual and joint association of $\beta$ -adrenergic receptor Arg347Cys polymorphism and plasma irbesartan concentration with blood pressure therapeutic response in Chinese hypertensive subjects. <i>Clinical Pharmacology and Therapeutics</i> , 2005, 78, 239-248.	4.7	21
34	A Common Haplotype onMethylenetetrahydrofolate ReductaseGene Modifies the Effect of Angiotensin-Converting Enzyme Inhibitor on Blood Pressure in Essential Hypertension Patientsâ€”A Family-Based Association Study. <i>Clinical and Experimental Hypertension</i> , 2005, 27, 509-521.	1.3	13
35	Association between human atrial natriuretic peptide Val7Met polymorphism and baseline blood pressure, plasma trough irbesartan concentrations, and the antihypertensive efficacy of irbesartan in Rural Chinese patients with essential hypertension. <i>Clinical Therapeutics</i> , 2005, 27, 1774-1784.	2.5	22
36	Association of cigarette smoking with steady state plasma concentration of irbesartan in male Chinese with hypertension. <i>Methods and Findings in Experimental and Clinical Pharmacology</i> , 2005, 27, 173.	0.8	4

#	ARTICLE	IF	CITATIONS
37	The C677T polymorphism of the methylenetetrahydrofolate reductase gene is associated with the level of decrease on diastolic blood pressure in essential hypertension patients treated by angiotensin-converting enzyme inhibitor. <i>Thrombosis Research</i> , 2004, 113, 361-369.	1.7	25