

Hyoun-Ah Kim

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8430389/publications.pdf>

Version: 2024-02-01

114
papers

2,274
citations

270111

25
h-index

312153

41
g-index

116
all docs

116
docs citations

116
times ranked

3228
citing authors

#	ARTICLE	IF	CITATIONS
1	Flare of adult-onset Still's disease following mRNA COVID-19 vaccination: a case report and review of literature. <i>Clinical Rheumatology</i> , 2022, 41, 1583-1589.	1.0	12
2	Clinical Features and Drug Retention of TNF Inhibitors in Older Patients with Ankylosing Spondylitis: Results from the KOBIO Registry. <i>BioDrugs</i> , 2022, , 1.	2.2	1
3	Three Clinical Clusters Identified through Hierarchical Cluster Analysis Using Initial Laboratory Findings in Korean Patients with Systemic Lupus Erythematosus. <i>Journal of Clinical Medicine</i> , 2022, 11, 2406.	1.0	5
4	S100A8 in Serum, Urine, and Saliva as a Potential Biomarker for Systemic Lupus Erythematosus. <i>Frontiers in Immunology</i> , 2022, 13, 886209.	2.2	10
5	Prevalence and Factors of Osteoporosis and High Risk of Osteoporotic Fracture in Patients with Ankylosing Spondylitis: A Multicenter Comparative Study of Bone Mineral Density and the Fracture Risk Assessment Tool. <i>Journal of Clinical Medicine</i> , 2022, 11, 2830.	1.0	7
6	Comparison of remission criteria in patients with rheumatoid arthritis treated with biologic or targeted synthetic disease-modifying anti-rheumatic drugs: results from a nationwide registry. <i>Therapeutic Advances in Musculoskeletal Disease</i> , 2022, 14, 1759720X2210963.	1.2	1
7	Elevated expression of TLR2 and its correlation with disease activity and clinical manifestations in adult-onset Still's disease. <i>Scientific Reports</i> , 2022, 12, .	1.6	1
8	Correlations between salivary gland scintigraphy and histopathologic data of salivary glands in patients with primary Sjogren's syndrome. <i>Clinical Rheumatology</i> , 2022, 41, 3083-3093.	1.0	1
9	Implications of Persistent Pain in Patients With Rheumatoid Arthritis Despite Remission Status: Data From the KOBIO Registry. <i>Journal of Rheumatic Diseases</i> , 2022, , .	0.4	2
10	Systemic immune-inflammation index combined with ferritin can serve as a reliable assessment score for adult-onset Still's disease. <i>Clinical Rheumatology</i> , 2021, 40, 661-668.	1.0	37
11	Lupus Low Disease Activity State Achievement Is Important for Reducing Adverse Outcomes in Pregnant Patients With Systemic Lupus Erythematosus. <i>Journal of Rheumatology</i> , 2021, 48, 707-716.	1.0	23
12	Salivary Immunoglobulin Gamma-3 Chain C Is a Promising Noninvasive Biomarker for Systemic Lupus Erythematosus. <i>International Journal of Molecular Sciences</i> , 2021, 22, 1374.	1.8	3
13	Anti-Inflammatory Effects of Low-Dose Glucocorticoids Compensate for Their Detrimental Effects on Bone Mineral Density in Patients with Rheumatoid Arthritis. <i>Journal of Clinical Medicine</i> , 2021, 10, 2944.	1.0	5
14	Comparison of Retention Rates Between Tumor Necrosis Factor- α Inhibitors in Patients With Ankylosing Spondylitis: Data From the Korean College of Rheumatology Biologics Registry. <i>Frontiers in Medicine</i> , 2021, 8, 689609.	1.2	6
15	The Assessment of Muscle Mass and Function in Patients with Long-Standing Rheumatoid Arthritis. <i>Journal of Clinical Medicine</i> , 2021, 10, 3458.	1.0	9
16	KOBIO, the First Web-based Korean Biologics Registry Operated With a Unified Platform Among Distinct Disease Entities. <i>Journal of Rheumatic Diseases</i> , 2021, 28, 176-182.	0.4	19
17	Salivary Biomarkers in Patients with Sjogren's Syndrome: A Systematic Review. <i>International Journal of Molecular Sciences</i> , 2021, 22, 12903.	1.8	14
18	An Update on the Pathogenic Role of Neutrophils in Systemic Juvenile Idiopathic Arthritis and Adult-Onset Still's Disease. <i>International Journal of Molecular Sciences</i> , 2021, 22, 13038.	1.8	17

#	ARTICLE	IF	CITATIONS
19	Tocilizumab-induced anaphylaxis in patients with adult-onset Still's disease and systemic juvenile idiopathic arthritis: a case-based review. <i>Rheumatology International</i> , 2020, 40, 791-798.	1.5	19
20	Trabecular Bone Score Is a Useful Parameter for the Prediction of Vertebral Fractures in Patients With Polymyalgia Rheumatica. <i>Journal of Clinical Densitometry</i> , 2020, 23, 373-380.	0.5	13
21	Retention Rate and Safety of Biosimilar CT-P13 in Rheumatoid Arthritis: Data from the Korean College of Rheumatology Biologics Registry. <i>BioDrugs</i> , 2020, 34, 89-98.	2.2	13
22	Retention Rate and Efficacy of the Biosimilar CT-P13 Versus Reference Infliximab in Patients with Ankylosing Spondylitis: A Propensity Score-Matched Analysis from the Korean College of Rheumatology Biologics Registry. <i>BioDrugs</i> , 2020, 34, 529-539.	2.2	7
23	Prevalence of osteoporosis in patients with systemic lupus erythematosus: A multicenter comparative study of the World Health Organization and fracture risk assessment tool criteria. <i>Osteoporosis and Sarcopenia</i> , 2020, 6, 173-178.	0.7	5
24	Roles of Interactions Between Toll-Like Receptors and Their Endogenous Ligands in the Pathogenesis of Systemic Juvenile Idiopathic Arthritis and Adult-Onset Still's Disease. <i>Frontiers in Immunology</i> , 2020, 11, 583513.	2.2	20
25	Obesity increases the incidence of new-onset lupus nephritis and organ damage during follow-up in patients with systemic lupus erythematosus. <i>Lupus</i> , 2020, 29, 578-586.	0.8	16
26	Seasonal Variations and Associated Factors of Gout Attacks: a Prospective Multicenter Study in Korea. <i>Journal of Korean Medical Science</i> , 2020, 35, e133.	1.1	10
27	The Practice Guideline for Vaccinating Korean Patients with Autoimmune Inflammatory Rheumatic Disease. <i>Infection and Chemotherapy</i> , 2020, 52, 252.	1.0	14
28	The Practice Guideline for Vaccinating Korean Patients With Autoimmune Inflammatory Rheumatic Disease. <i>Journal of Rheumatic Diseases</i> , 2020, 27, 182-202.	0.4	2
29	Retention rate and long-term safety of biosimilar CT-P13 in patients with ankylosing spondylitis: data from the Korean College of Rheumatology Biologics registry. <i>Clinical and Experimental Rheumatology</i> , 2020, 38, 267-274.	0.4	6
30	Neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio are associated with disease activity in polymyalgia rheumatica. <i>Journal of Clinical Laboratory Analysis</i> , 2019, 33, e23000.	0.9	21
31	Biomarkers for adult-onset still's disease: up-to-date. <i>Expert Review of Molecular Diagnostics</i> , 2019, 19, 655-657.	1.5	3
32	Clinical Impact of the Fracture Risk Assessment Tool on the Treatment Decision for Osteoporosis in Patients with Knee Osteoarthritis: A Multicenter Comparative Study of the Fracture Risk Assessment Tool and World Health Organization Criteria. <i>Journal of Clinical Medicine</i> , 2019, 8, 918.	1.0	5
33	Associated clinical factors for serious infections in patients with systemic lupus erythematosus. <i>Scientific Reports</i> , 2019, 9, 9704.	1.6	30
34	Differences in Expression of Human Leukocyte Antigen Class II Subtypes and T Cell Subsets in Behçet's Disease with Arthritis. <i>International Journal of Molecular Sciences</i> , 2019, 20, 5044.	1.8	3
35	Rituximab biosimilar CT-P10 for the treatment of rheumatoid arthritis. <i>Expert Opinion on Biological Therapy</i> , 2019, 19, 979-986.	1.4	5
36	Clinical characteristics and role of whole-body bone scan in multifocal osteonecrosis. <i>BMC Musculoskeletal Disorders</i> , 2019, 20, 23.	0.8	8

#	ARTICLE	IF	CITATIONS
37	Mindfulness-based cognitive therapy in Korean patients with systemic lupus erythematosus: A pilot study. <i>Complementary Therapies in Clinical Practice</i> , 2019, 35, 18-21.	0.7	11
38	Histopathology and expression of the chemokines CXCL10, CXCL13, and CXCR3 and the endogenous TLR-4 ligand S100A8/A9 in lymph nodes of patients with adult-onset Still's disease. <i>Scientific Reports</i> , 2019, 9, 7517.	1.6	15
39	Neutrophil Extracellular Traps May Contribute to the Pathogenesis in Adult-onset Still Disease. <i>Journal of Rheumatology</i> , 2019, 46, 1560-1569.	1.0	31
40	The role of damage-associated molecular pattern for pathogenesis and biomarkers in adult-onset Still's disease. <i>Expert Review of Molecular Diagnostics</i> , 2019, 19, 459-468.	1.5	19
41	Body mass index and glucocorticoid dose contribute to subclinical atherosclerosis in Korean patients with systemic lupus erythematosus: A prospective 4-year follow-up study. <i>International Journal of Rheumatic Diseases</i> , 2019, 22, 1410-1418.	0.9	11
42	The Liver X Receptor Is Upregulated in Monocyte-Derived Macrophages and Modulates Inflammatory Cytokines Based on LXR Polymorphism. <i>Mediators of Inflammation</i> , 2019, 2019, 1-12.	1.4	4
43	The Role of Natural Killer Cells in Inflammation in Active Adult-onset Still Disease. <i>Journal of Rheumatology</i> , 2019, 46, 1253-1255.	1.0	3
44	Standardized, musculoskeletal ultrasonographic reference values for healthy Korean adults. <i>Korean Journal of Internal Medicine</i> , 2019, 34, 1372-1380.	0.7	5
45	The levels of CXCL12 and its receptor, CXCR4, as a biomarker of disease activity and cutaneous manifestation in adult-onset Still's disease. <i>Clinical and Experimental Rheumatology</i> , 2019, 37 Suppl 121, 67-73.	0.4	4
46	Outcome and predictors of renal survival in patients with lupus nephritis: Comparison between cyclophosphamide and mycophenolate mofetil. <i>International Journal of Rheumatic Diseases</i> , 2018, 21, 1031-1039.	0.9	14
47	Effects of risk factors for and components of metabolic syndrome on the quality of life of patients with systemic lupus erythematosus: a structural equation modeling approach. <i>Quality of Life Research</i> , 2018, 27, 105-113.	1.5	9
48	Patient with Systemic Lupus Erythematosus Combined with Erosive Arthritis was Treated Successfully with Tocilizumab: A Case Report. <i>Journal of Rheumatic Diseases</i> , 2018, 25, 144.	0.4	4
49	Effect of fenofibrate on uric acid level in patients with gout. <i>Scientific Reports</i> , 2018, 8, 16767.	1.6	17
50	Off-label use of tocilizumab to treat non-juvenile idiopathic arthritis in pediatric rheumatic patients: a literature review. <i>Pediatric Rheumatology</i> , 2018, 16, 79.	0.9	17
51	Prevalence and Fracture Risk of Osteoporosis in Patients with Rheumatoid Arthritis: A Multicenter Comparative Study of the FRAX and WHO Criteria. <i>Journal of Clinical Medicine</i> , 2018, 7, 507.	1.0	32
52	Association between brain-derived neurotrophic factor gene polymorphisms and fibromyalgia in a Korean population: a multicenter study. <i>Arthritis Research and Therapy</i> , 2018, 20, 220.	1.6	11
53	Characteristic patterns of HLA presentation and T cell differentiation in adult-onset Still's disease. <i>International Journal of Immunopathology and Pharmacology</i> , 2018, 32, 205873841879128.	1.0	13
54	Application of the international league against rheumatism classification criteria for systemic juvenile idiopathic arthritis as a prognostic factor in patients with adults-onset Still's disease. <i>Pediatric Rheumatology</i> , 2018, 16, 9.	0.9	6

#	ARTICLE	IF	CITATIONS
55	Association of depression with socioeconomic status, anticardiolipin antibodies, and organ damage in patients with systemic lupus erythematosus: results from the KORNET registry. <i>Clinical and Experimental Rheumatology</i> , 2018, 36, 627-635.	0.4	7
56	Comparative effectiveness of treatment options after conventional DMARDs failure in rheumatoid arthritis. <i>Rheumatology International</i> , 2017, 37, 975-982.	1.5	8
57	Characteristics and outcomes of rheumatoid arthritis patients who started biosimilar infliximab. <i>Rheumatology International</i> , 2017, 37, 1007-1014.	1.5	10
58	Switching profiles in a population-based cohort of rheumatoid arthritis receiving biologic therapy: results from the KOBIO registry. <i>Clinical Rheumatology</i> , 2017, 36, 1013-1022.	1.0	20
59	An update: current and prospective treatments for adult-onset Still's disease. <i>Expert Opinion on Orphan Drugs</i> , 2017, 5, 421-429.	0.5	1
60	Elevated circulating levels of the interferon- γ -induced chemokines are associated with disease activity and cutaneous manifestations in adult-onset Still's disease. <i>Scientific Reports</i> , 2017, 7, 46652.	1.6	49
61	Serum Levels of Interleukin 33 and Soluble ST2 Are Associated with the Extent of Disease Activity and Cutaneous Manifestations in Patients with Active Adult-onset Still's Disease. <i>Journal of Rheumatology</i> , 2017, 44, 740-747.	1.0	15
62	Usefulness of neutrophil-to-lymphocyte ratio as a biomarker for diagnosing infections in patients with systemic lupus erythematosus. <i>Clinical Rheumatology</i> , 2017, 36, 2479-2485.	1.0	36
63	The neutrophil-to-lymphocyte ratio could be a good diagnostic marker and predictor of relapse in patients with adult-onset Still's disease. <i>Medicine (United States)</i> , 2017, 96, e7546.	0.4	27
64	Highly Expression of CD11b and CD32 on Peripheral Blood Mononuclear Cells from Patients with Adult-Onset Still's Disease. <i>International Journal of Molecular Sciences</i> , 2017, 18, 202.	1.8	10
65	The Correlation of CD206, CD209, and Disease Severity in Behçet's Disease with Arthritis. <i>Mediators of Inflammation</i> , 2017, 2017, 1-13.	1.4	8
66	Determinants of quality of life in patients with fibromyalgia: A structural equation modeling approach. <i>PLoS ONE</i> , 2017, 12, e0171186.	1.1	51
67	TLR4 Endogenous Ligand S100A8/A9 Levels in Adult-Onset Still's Disease and Their Association with Disease Activity and Clinical Manifestations. <i>International Journal of Molecular Sciences</i> , 2016, 17, 1342.	1.8	37
68	Impact of Childbearing Decisions on Family Size of Korean Women with Systemic Lupus Erythematosus. <i>Journal of Korean Medical Science</i> , 2016, 31, 729.	1.1	5
69	Polymorphisms of the <i>TRPV2</i> and <i>TRPV3</i> genes associated with fibromyalgia in a Korean population. <i>Rheumatology</i> , 2016, 55, 1518-1527.	0.9	27
70	Elevated high-mobility group B1 levels in active adult-onset Still's disease associated with systemic score and skin rash. <i>Clinical Rheumatology</i> , 2016, 35, 1937-1942.	1.0	20
71	Severity of fibromyalgia symptoms is associated with socioeconomic status and not obesity in Korean patients. <i>Clinical and Experimental Rheumatology</i> , 2016, 34, S83-8.	0.4	4
72	<i>FCRIIB</i> Gene Polymorphisms Are Associated with Disease Risk and Clinical Manifestations of Systemic Lupus Erythematosus in Koreans. <i>Tohoku Journal of Experimental Medicine</i> , 2015, 236, 185-191.	0.5	6

#	ARTICLE	IF	CITATIONS
73	Association of CXCL10 and CXCL13 levels with disease activity and cutaneous manifestation in active adult-onset Still's disease. <i>Arthritis Research and Therapy</i> , 2015, 17, 260.	1.6	39
74	Elevated Salivary Alpha-Amylase Level, Association Between Depression and Disease Activity, and Stress as a Predictor of Disease Flare in Systemic Lupus Erythematosus. <i>Medicine (United States)</i> , 2015, 94, e1184.	0.4	26
75	Seasonality of Gout in Korea: A Multicenter Study. <i>Journal of Korean Medical Science</i> , 2015, 30, 240.	1.1	14
76	The Pathologic Findings of Skin, Lymph Node, Liver, and Bone Marrow in Patients With Adult-Onset Still Disease. <i>Medicine (United States)</i> , 2015, 94, e787.	0.4	36
77	Role of bone scan in the assessment of polymyositis/dermatomyositis. <i>Clinical Rheumatology</i> , 2015, 34, 699-706.	1.0	5
78	Biomarkers for systemic lupus erythematosus: an update. <i>International Journal of Clinical Rheumatology</i> , 2015, 10, 195-204.	0.3	2
79	Prediction for TNF Inhibitor Users in RA Patients According to Reimbursement Criteria Based on DAS28. <i>Journal of Rheumatic Diseases</i> , 2014, 21, 64.	0.4	6
80	Serum S100A12 May Be a Useful Biomarker of Disease Activity in Adult-onset Still's Disease. <i>Journal of Rheumatology</i> , 2014, 41, 2403-2408.	1.0	47
81	Autoantibodies to C-Reactive Protein in Incomplete Lupus and Systemic Lupus Erythematosus. <i>Journal of Investigative Medicine</i> , 2014, 62, 890-893.	0.7	10
82	Liver X receptors alpha gene (NR1H3) promoter polymorphisms are associated with systemic lupus erythematosus in Koreans. <i>Arthritis Research and Therapy</i> , 2014, 16, R112.	1.6	15
83	Serum growth arrest-specific protein 6 levels are elevated in adult-onset Still's disease. <i>Clinical Rheumatology</i> , 2014, 33, 865-868.	1.0	3
84	Serum Growth Arrest-Specific Protein 6 Levels are a Reliable Biomarker of Disease Activity in Systemic Lupus Erythematosus. <i>Journal of Clinical Immunology</i> , 2013, 33, 143-150.	2.0	22
85	Multicenter Retrospective Analysis of the Effectiveness and Safety of Rituximab in Korean Patients with Refractory Systemic Lupus Erythematosus. <i>Autoimmune Diseases</i> , 2012, 2012, 1-6.	2.7	18
86	Nonsteroidal Antiinflammatory Drugs (NSAID) Versus NSAID with Hydroxychloroquine in Treatment of Chemotherapy-related Arthropathy: Open-label Multicenter Pilot Study. <i>Journal of Rheumatology</i> , 2012, 39, 1902-1903.	1.0	4
87	Serum S100A8/A9, But Not Follistatin-like Protein 1 and Interleukin 18, May Be a Useful Biomarker of Disease Activity in Adult-onset Still's Disease. <i>Journal of Rheumatology</i> , 2012, 39, 1399-1406.	1.0	60
88	C-reactive Protein Is a More Sensitive and Specific Marker for Diagnosing Bacterial Infections in Systemic Lupus Erythematosus Compared to S100A8/A9 and Procalcitonin. <i>Journal of Rheumatology</i> , 2012, 39, 728-734.	1.0	55
89	Induction of Remission is Difficult due to Frequent Relapse during Tapering Steroids in Korean Patients with Polymyalgia Rheumatica. <i>Journal of Korean Medical Science</i> , 2012, 27, 22.	1.1	12
90	Therapeutic responses and prognosis in adult-onset Still's disease. <i>Rheumatology International</i> , 2012, 32, 1291-1298.	1.5	64

#	ARTICLE	IF	CITATIONS
91	Dickkopf-1 level is lower in patients with ankylosing spondylitis than in healthy people and is not influenced by anti-tumor necrosis factor therapy. <i>Rheumatology International</i> , 2012, 32, 2523-2527.	1.5	98
92	Korean Observational Study Network for Arthritis (KORONA): Establishment of a Prospective Multicenter Cohort for Rheumatoid Arthritis in South Korea. <i>Seminars in Arthritis and Rheumatism</i> , 2012, 41, 745-751.	1.6	54
93	Intestinal pseudo-obstruction caused by neuromyopathy in a patient with systemic sclerosis. <i>Modern Rheumatology</i> , 2012, 22, 912-918.	0.9	2
94	Smoking habits influence pain and functional and psychiatric features in fibromyalgia. <i>Joint Bone Spine</i> , 2011, 78, 259-265.	0.8	18
95	Vitamin D may not be a good marker of disease activity in Korean patients with systemic lupus erythematosus. <i>Rheumatology International</i> , 2011, 31, 1189-1194.	1.5	51
96	Les habitudes tabagiques influencent la douleur et les Ã©lÃ©ments fonctionnels et psychiatriques dans la fibromyalgie. <i>Revue Du Rhumatisme (Edition Francaise)</i> , 2011, 78, 173-179.	0.0	0
97	Women with Fibromyalgia Have Lower Levels of Calcium, Magnesium, Iron and Manganese in Hair Mineral Analysis. <i>Journal of Korean Medical Science</i> , 2011, 26, 1253.	1.1	32
98	Beta2-Microglobulin Can Be a Disease Activity Marker in Systemic Lupus Erythematosus. <i>American Journal of the Medical Sciences</i> , 2010, 339, 337-340.	0.4	32
99	Prevalence of Human Papilloma Virus Infections and Cervical Cytological Abnormalities among Korean Women with Systemic Lupus Erythematosus. <i>Journal of Korean Medical Science</i> , 2010, 25, 1431.	1.1	37
100	Interleukin 6 Gene Polymorphisms Are Associated with Systemic Lupus Erythematosus in Koreans. <i>Journal of Rheumatology</i> , 2010, 37, 2251-2258.	1.0	22
101	Chronic periaortitis with antiphospholipid syndrome. <i>International Journal of Rheumatic Diseases</i> , 2010, 13, 91-93.	0.9	3
102	C-Reactive Protein Gene Polymorphisms in Disease Susceptibility and Clinical Manifestations of Korean Systemic Lupus Erythematosus. <i>Journal of Rheumatology</i> , 2009, 36, 2238-2243.	1.0	25
103	An unusual clinical manifestation of Takayasu's arteritis: Spinal cord compression. <i>Joint Bone Spine</i> , 2009, 76, 209-212.	0.8	8
104	SLE developed in a preadolescent child with Klinefelter syndrome who had no detectable sex hormone. <i>Rheumatology International</i> , 2009, 29, 975-977.	1.5	1
105	Neurogenic Bladder in a Patient With Systemic Lupus Erythematosus and Cerebral Involvement. <i>Journal of Clinical Rheumatology</i> , 2009, 15, 40-41.	0.5	0
106	An Antinuclear Antibody-Negative Patient With Lupus Nephritis. <i>Korean Journal of Internal Medicine</i> , 2009, 24, 76.	0.7	23
107	Clinical characteristics of lupus myocarditis in Korea. <i>Rheumatology International</i> , 2008, 28, 275-280.	1.5	25
108	The antichromatin antibodies can be useful as a diagnostic tool and disease activity marker of systemic lupus erythematosus in Koreans. <i>Clinical Immunology</i> , 2008, 128, 277-283.	1.4	15

#	ARTICLE	IF	CITATIONS
109	Prevalence of work-related symptoms and serum-specific antibodies to wheat flour in exposed workers in the bakery industry. <i>Respiratory Medicine</i> , 2008, 102, 548-555.	1.3	41
110	Cytokines and their receptors as biomarkers of systemic lupus erythematosus. <i>Expert Review of Molecular Diagnostics</i> , 2008, 8, 189-198.	1.5	38
111	Treatment of Atopic Dermatitis with a Combination of Allergen-Specific Immunotherapy and a Histamine-Immunoglobulin Complex. <i>International Archives of Allergy and Immunology</i> , 2008, 146, 235-240.	0.9	26
112	Identification of <i>Dioscorea Batatas</i> (Sanyak) Allergen as an Inhalant and Oral Allergen. <i>Journal of Korean Medical Science</i> , 2008, 23, 72.	1.1	14
113	Association of β_2 -Adrenergic Receptor Polymorphism with the Phenotype of Aspirin-Intolerant Acute Urticaria. <i>Yonsei Medical Journal</i> , 2007, 48, 1079.	0.9	10
114	Cytokine IL-6 and IL-10 as Biomarkers in Systemic Lupus Erythematosus. <i>Journal of Clinical Immunology</i> , 2007, 27, 461-466.	2.0	321