

Ryohei Yamamoto

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8431187/publications.pdf>

Version: 2024-02-01

82
papers

1,927
citations

279487

23
h-index

288905

40
g-index

84
all docs

84
docs citations

84
times ranked

2722
citing authors

#	ARTICLE	IF	CITATIONS
1	Living alone and prediction of weight gain and overweight/obesity in university students: a retrospective cohort study. <i>Journal of American College Health</i> , 2023, 71, 1417-1426.	0.8	4
2	Care needs level in long-term care insurance system and family caregivers' self-perceived time-dependent burden in patients with home palliative care for cancer: a cross-sectional study. <i>Supportive Care in Cancer</i> , 2022, 30, 1587-1596.	1.0	5
3	Time to remission of proteinuria and incidence of relapse in patients with steroid-sensitive minimal change disease and focal segmental glomerulosclerosis: the Japan Nephrotic Syndrome Cohort Study. <i>Journal of Nephrology</i> , 2022, 35, 1135-1144.	0.9	3
4	Skipping Breakfast and Incidence of Frequent Alcohol Drinking in University Students in Japan: A Retrospective Cohort Study. <i>Nutrients</i> , 2022, 14, 2657.	1.7	4
5	Occupational sedentary behavior and prediction of proteinuria in young to middle-aged adults: a retrospective cohort study. <i>Journal of Nephrology</i> , 2021, 34, 719-728.	0.9	8
6	Associations of Skipping Breakfast, Lunch, and Dinner with Weight Gain and Overweight/Obesity in University Students: A Retrospective Cohort Study. <i>Nutrients</i> , 2021, 13, 271.	1.7	29
7	Efficacy of aerobic exercise on the cardiometabolic and renal outcomes in patients with chronic kidney disease: a systematic review of randomized controlled trials. <i>Journal of Nephrology</i> , 2021, 34, 155-164.	0.9	20
8	Serum Klotho Levels Contribute to the Prevention of Disease Progression. <i>International Journal of General Medicine</i> , 2021, Volume 14, 229-236.	0.8	7
9	Alpha-Klotho is a novel predictor of treatment responsiveness in patients with heart failure. <i>Scientific Reports</i> , 2021, 11, 2058.	1.6	11
10	Incidence and factors associated with prescribing renin-angiotensin system inhibitors in adult idiopathic nephrotic syndrome: A nationwide cohort study. <i>Journal of Clinical Hypertension</i> , 2021, 23, 999-1007.	1.0	4
11	Association of weekday-to-weekend sleep differences and stress response among a Japanese working population: a cross-sectional study. <i>Sleep Medicine</i> , 2021, 82, 159-164.	0.8	3
12	Demographic, clinical characteristics and treatment outcomes of immune-complex membranoproliferative glomerulonephritis and C3 glomerulonephritis in Japan: A retrospective analysis of data from the Japan Renal Biopsy Registry. <i>PLoS ONE</i> , 2021, 16, e0257397.	1.1	10
13	A digest of the Evidence-Based Clinical Practice Guideline for Nephrotic Syndrome 2020. <i>Clinical and Experimental Nephrology</i> , 2021, 25, 1277-1285.	0.7	21
14	Associations of kidney tests at medical facilities and health checkups with incidence of end-stage kidney disease: a retrospective cohort study. <i>Scientific Reports</i> , 2021, 11, 20717.	1.6	4
15	Vegetable Preference and Prediction of Proteinuria: A Retrospective Cohort Study. <i>Annals of Nutrition and Metabolism</i> , 2021, 77, 337-343.	1.0	1
16	Serum FGF21 levels are altered by various factors including lifestyle behaviors in male subjects. <i>Scientific Reports</i> , 2021, 11, 22632.	1.6	9
17	Sleep debt and prevalence of proteinuria in subjects with short sleep duration on weekdays: a cross-sectional study. <i>Clinical and Experimental Nephrology</i> , 2020, 24, 143-150.	0.7	4
18	Body mass index modifies the association between frequency of alcohol consumption and incidence of hypertension in men but not in women: a retrospective cohort study. <i>Hypertension Research</i> , 2020, 43, 322-330.	1.5	7

#	ARTICLE	IF	CITATIONS
19	Frequency of alcohol drinking modifies the association between salt intake and albuminuria: a 1-year observational study. <i>Hypertension Research</i> , 2020, 43, 1249-1256.	1.5	3
20	Efficacy of corticosteroid therapy for IgA nephropathy patients stratified by kidney function and proteinuria. <i>Clinical and Experimental Nephrology</i> , 2020, 24, 927-934.	0.7	4
21	Frequency of Breakfast, Lunch, and Dinner and Incidence of Proteinuria: A Retrospective Cohort Study. <i>Nutrients</i> , 2020, 12, 3549.	1.7	5
22	P0819THE EFFECT OF SMOKING UPON CKD IN A JAPANESE NATIONWIDE 6-YEARS FOLLOW UP SERVEY. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, .	0.4	0
23	Is alcohol drinking a key factor contributing to salt sensitivity?. <i>Journal of Clinical Hypertension</i> , 2020, 22, 1103-1105.	1.0	0
24	Drinking frequency modifies an association between salt intake and blood pressure: A cohort study. <i>Journal of Clinical Hypertension</i> , 2020, 22, 649-655.	1.0	5
25	Incidence of remission and relapse of proteinuria, end-stage kidney disease, mortality, and major outcomes in primary nephrotic syndrome: the Japan Nephrotic Syndrome Cohort Study (JNSCS). <i>Clinical and Experimental Nephrology</i> , 2020, 24, 526-540.	0.7	33
26	Better remission rates in elderly Japanese patients with primary membranous nephropathy in nationwide real-world practice: The Japan Nephrotic Syndrome Cohort Study (JNSCS). <i>Clinical and Experimental Nephrology</i> , 2020, 24, 893-909.	0.7	6
27	Causal relations of health indices inferred statistically using the DirectLiNGAM algorithm from big data of Osaka prefecture health checkups. <i>PLoS ONE</i> , 2020, 15, e0243229.	1.1	6
28	Proteinuria-associated renal magnesium wasting leads to hypomagnesemia: a common electrolyte abnormality in chronic kidney disease. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, 1154-1162.	0.4	36
29	A Randomized Trial of Magnesium Oxide and Oral Carbon Adsorbent for Coronary Artery Calcification in Predialysis CKD. <i>Journal of the American Society of Nephrology: JASN</i> , 2019, 30, 1073-1085.	3.0	98
30	Implication of alpha-Klotho as the predictive factor of stress. <i>Journal of Investigative Medicine</i> , 2019, 67, 1082-1086.	0.7	8
31	Regional variations in immunosuppressive therapy in patients with primary nephrotic syndrome: the Japan nephrotic syndrome cohort study. <i>Clinical and Experimental Nephrology</i> , 2018, 22, 1266-1280.	0.7	21
32	An implication of Klotho-related molecules in different smoking-related health outcomes between men and women. <i>Clinica Chimica Acta</i> , 2018, 476, 44-48.	0.5	13
33	Sleep Quality and Sleep Duration with CKD are Associated with Progression to ESKD. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2018, 13, 1825-1832.	2.2	59
34	Alcohol consumption and incidence of proteinuria: a retrospective cohort study. <i>Clinical and Experimental Nephrology</i> , 2018, 22, 1133-1142.	0.7	20
35	The association of alcohol and smoking with CKD in a Japanese nationwide cross-sectional survey. <i>Hypertension Research</i> , 2017, 40, 771-778.	1.5	22
36	Modified forelimb grip strength test detects aging-associated physiological decline in skeletal muscle function in male mice. <i>Scientific Reports</i> , 2017, 7, 42323.	1.6	144

#	ARTICLE	IF	CITATIONS
37	Body Mass Index Modifies an Association between Self-Reported Regular Exercise and Proteinuria. <i>Journal of Atherosclerosis and Thrombosis</i> , 2016, 23, 402-412.	0.9	10
38	IFI27 Is a Useful Genetic Marker for Diagnosis of Immunoglobulin A Nephropathy and Membranous Nephropathy Using Peripheral Blood. <i>PLoS ONE</i> , 2016, 11, e0153252.	1.1	18
39	Chiral amino acid metabolomics for novel biomarker screening in the prognosis of chronic kidney disease. <i>Scientific Reports</i> , 2016, 6, 26137.	1.6	162
40	Evidence-based clinical practice guidelines for IgA nephropathy 2014. <i>Clinical and Experimental Nephrology</i> , 2016, 20, 511-535.	0.7	56
41	Association between N-terminal pro-brain natriuretic peptide and adiponectin in healthy Japanese men. <i>Clinica Chimica Acta</i> , 2016, 460, 138-141.	0.5	8
42	Identification of biomarkers for development of end-stage kidney disease in chronic kidney disease by metabolomic profiling. <i>Scientific Reports</i> , 2016, 6, 26138.	1.6	50
43	Serum Uric Acid Level Predicts Progression of IgA Nephropathy in Females but Not in Males. <i>PLoS ONE</i> , 2016, 11, e0160828.	1.1	29
44	Association between Density of Coronary Artery Calcification and Serum Magnesium Levels among Patients with Chronic Kidney Disease. <i>PLoS ONE</i> , 2016, 11, e0163673.	1.1	42
45	Minimal Change Nephrotic Syndrome Which Was Most Likely Caused by Chronic Sinusitis. <i>Internal Medicine</i> , 2015, 54, 2373-2375.	0.3	2
46	<i>ACE</i> Insertion/Deletion Polymorphism and Cardiovascular Effect of Hypercholesterolemia. <i>Journal of Atherosclerosis and Thrombosis</i> , 2015, 22, 338-341.	0.9	3
47	<i>ACE</i> insertion/deletion polymorphism (rs1799752) modifies the renoprotective effect of renin-angiotensin system blockade in patients with IgA nephropathy. <i>JRAAS - Journal of the Renin-Angiotensin-Aldosterone System</i> , 2015, 16, 633-641.	1.0	12
48	Genome-wide association study of IgA nephropathy using 23â€‰‰465 microsatellite markers in a Japanese population. <i>Journal of Human Genetics</i> , 2015, 60, 573-580.	1.1	13
49	Urine Osmolarity Predicts the Body Weight-Reduction Response to Tolvaptan in Chronic Kidney Disease Patients: A Retrospective, Observational Study. <i>Nephron</i> , 2015, 130, 8-12.	0.9	14
50	CD16+CD56+ cells are a potential culprit for hematuria in IgA nephropathy. <i>Clinical and Experimental Nephrology</i> , 2015, 19, 216-224.	0.7	13
51	Periodontal Disease Bacteria Specific to Tonsil in IgA Nephropathy Patients Predicts the Remission by the Treatment. <i>PLoS ONE</i> , 2014, 9, e81636.	1.1	35
52	Comparison of Methylprednisolone Plus Prednisolone with Prednisolone Alone as Initial Treatment in Adult-Onset Minimal Change Disease. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2014, 9, 1040-1048.	2.2	29
53	An association between serum Î³-glutamyltransferase and proteinuria in drinkers and non-drinkers: a Japanese nationwide cross-sectional survey. <i>Clinical and Experimental Nephrology</i> , 2014, 18, 899-910.	0.7	5
54	Smoking Is a Risk Factor for the Progression of Idiopathic Membranous Nephropathy. <i>PLoS ONE</i> , 2014, 9, e100835.	1.1	13

#	ARTICLE	IF	CITATIONS
55	Patient Age and the Prognosis of Idiopathic Membranous Nephropathy. PLoS ONE, 2014, 9, e110376.	1.1	32
56	Is Low Testosterone Concentration a Risk Factor for Metabolic Syndrome in Healthy Middle-aged Men?. Urology, 2013, 82, 814-819.	0.5	26
57	Age and prediction of remission and relapse of proteinuria and corticosteroid-related adverse events in adult-onset minimal-change disease: a retrospective cohort study. Clinical and Experimental Nephrology, 2013, 17, 839-847.	0.7	21
58	A candidate gene approach to genetic contributors to the development of IgA nephropathy. Nephrology Dialysis Transplantation, 2012, 27, 1020-1030.	0.4	17
59	Cigarette smoking and chronic kidney diseases. Hypertension Research, 2012, 35, 261-265.	1.5	54
60	Low serum free testosterone level is associated with carotid intima-media thickness in middle-aged Japanese men. Endocrine Journal, 2012, 59, 809-815.	0.7	17
61	Use of xanthine oxidase inhibitor febuxostat inhibits renal interstitial inflammation and fibrosis in unilateral ureteral obstructive nephropathy. Clinical and Experimental Nephrology, 2012, 16, 549-556.	0.7	84
62	Complete remission within 2 years predicts a good prognosis after methylprednisolone pulse therapy in patients with IgA nephropathy. Clinical and Experimental Nephrology, 2012, 16, 883-891.	0.7	28
63	Serum hepcidin-25 levels and anemia in non-dialysis chronic kidney disease patients: a cross-sectional study. Nephrology Dialysis Transplantation, 2012, 27, 1076-1083.	0.4	62
64	Quantitative change of IgA hinge O-glycan composition is a novel marker of therapeutic responses of IgA nephropathy. Biochemical and Biophysical Research Communications, 2012, 428, 339-342.	1.0	19
65	Gene polymorphisms contributing to hypertension in immunoglobulin A nephropathy. Clinical and Experimental Nephrology, 2012, 16, 250-258.	0.7	13
66	Self-reported Sleep Duration and Prediction of Proteinuria: A Retrospective Cohort Study. American Journal of Kidney Diseases, 2012, 59, 343-355.	2.1	73
67	Towards developing new strategies to reduce the adverse side-effects of nonsteroidal anti-inflammatory drugs. Clinical and Experimental Nephrology, 2012, 16, 25-29.	0.7	19
68	Urinary Type IV Collagen in Nondiabetic Kidney Disease. Nephron Clinical Practice, 2011, 117, c160-c166.	2.3	18
69	Microarray Analysis of Tonsils of IgA Nephropathy Patients. Advances in Oto-Rhino-Laryngology, 2011, 72, 75-78.	1.6	1
70	Incidence of symptomatic stroke and cancer in chronic kidney disease patients treated with epoetins. Clinical and Experimental Nephrology, 2010, 14, 445-452.	0.7	19
71	Oxford IgA nephropathy classification: valid for children?. Nature Reviews Nephrology, 2010, 6, 638-639.	4.1	1
72	Impact of Age and Overt Proteinuria on Outcomes of Stage 3 to 5 Chronic Kidney Disease in a Referred Cohort. Clinical Journal of the American Society of Nephrology: CJASN, 2010, 5, 1558-1565.	2.2	54

#	ARTICLE	IF	CITATIONS
73	Specialist care and improved long-term survival of dialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2010, 25, 1930-1935.	0.4	9
74	Cigarette Smoking and Progression of IgA Nephropathy. <i>American Journal of Kidney Diseases</i> , 2010, 56, 313-324.	2.1	84
75	Microarray analysis of tonsils in immunoglobulin A nephropathy patients. <i>Biochemical and Biophysical Research Communications</i> , 2010, 393, 565-570.	1.0	16
76	Severe adverse effects of 5-fluorouracil in S-1 were lessened by haemodialysis due to elimination of the drug. <i>CKJ: Clinical Kidney Journal</i> , 2009, 2, 152-154.	1.4	6
77	Early Plasma Exchange for Progressive Liver Failure in Recipients of Adult-to-Adult Living-Related Liver Transplants. <i>Blood Purification</i> , 2009, 28, 40-46.	0.9	11
78	A novel classification for IgA nephropathy. <i>Kidney International</i> , 2009, 76, 477-480.	2.6	18
79	A candidate gene approach to genetic prognostic factors of IgA nephropathy—a result of Polymorphism REsearch to Distinguish genetic factors Contributing To progression of IgA Nephropathy (PREDICT-IgAN). <i>Nephrology Dialysis Transplantation</i> , 2009, 24, 3686-3694.	0.4	19
80	Even partial remission of proteinuria is associated with better renal outcome in patients with IgA nephropathy. <i>Nature Clinical Practice Nephrology</i> , 2008, 4, 246-247.	2.0	1
81	Thiazide diuretics enhance nocturnal blood pressure fall and reduce proteinuria in immunoglobulin A nephropathy treated with angiotensin II modulators. <i>Journal of Hypertension</i> , 2005, 23, 861-865.	0.3	41
82	ANCA-negative pauci-immune crescentic glomerulonephritis complicated with recurrent massive gastrointestinal hemorrhage. <i>Clinical and Experimental Nephrology</i> , 2005, 9, 174-178.	0.7	11