

# Wai Lydia Tai

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8435088/publications.pdf>

Version: 2024-02-01

13  
papers

275  
citations

949033

11  
h-index

1255698

13  
g-index

13  
all docs

13  
docs citations

13  
times ranked

585  
citing authors

#	ARTICLE	IF	CITATIONS
1	Central Endothelin-1 Confers Analgesia by Triggering Spinal Neuronal Histone Deacetylase 5 (HDAC5) Nuclear Exclusion in Peripheral Neuropathic Pain in Mice. <i>Journal of Pain</i> , 2021, 22, 454-471.	0.7	2
2	Histone deacetylase 5 (HDAC5) regulates neuropathic pain through SRY-related HMG-box 10 (SOX10)-dependent mechanism in mice. <i>Pain</i> , 2018, 159, 526-539.	2.0	19
3	Adiponectin regulates thermal nociception in a mouse model of neuropathic pain. <i>British Journal of Anaesthesia</i> , 2018, 120, 1356-1367.	1.5	34
4	Enriched Environment and Effects on Neuropathic Pain: Experimental Findings and Mechanisms. <i>Pain Practice</i> , 2018, 18, 1068-1082.	0.9	25
5	Suppression of Pax2 Attenuates Allodynia and Hyperalgesia through ET-1-ETAR-NFAT5 Signaling in a Rat Model of Neuropathic Pain. <i>Neuroscience</i> , 2018, 384, 139-151.	1.1	12
6	Pro-resolving mediator maresin 1 ameliorates pain hypersensitivity in a rat spinal nerve ligation model of neuropathic pain. <i>Journal of Pain Research</i> , 2018, Volume 11, 1511-1519.	0.8	32
7	Endocannabinoid activation of CB <sub>1</sub> receptors contributes to long-lasting reversal of neuropathic pain by repetitive spinal cord stimulation. <i>European Journal of Pain</i> , 2017, 21, 804-814.	1.4	32
8	MG53 anchored by dysferlin to cell membrane reduces hepatocyte apoptosis which induced by ischaemia/reperfusion injury <i>in vivo</i> and <i>in vitro</i> . <i>Journal of Cellular and Molecular Medicine</i> , 2017, 21, 2503-2513.	1.6	34
9	Sex differences in complex regional pain syndrome type I (CRPS-I) in mice. <i>Journal of Pain Research</i> , 2017, Volume 10, 1811-1819.	0.8	20
10	Targeted Overexpression of Astrocytic Endothelin-1 Attenuates Neuropathic Pain by Upregulating Spinal Excitatory Amino Acid Transporter-2. <i>Journal of Molecular Neuroscience</i> , 2015, 57, 90-96.	1.1	3
11	Central Administration of C-X-C Chemokine Receptor Type 4 Antagonist Alleviates the Development and Maintenance of Peripheral Neuropathic Pain in Mice. <i>PLoS ONE</i> , 2014, 9, e104860.	1.1	28
12	Over-expression of astrocytic ET-1 attenuates neuropathic pain by inhibition of ERK1/2 and Akt(s) via activation of ETA receptor. <i>Molecular and Cellular Neurosciences</i> , 2014, 60, 26-35.	1.0	16
13	Over-expression of endothelin-1 in astrocytes, but not endothelial cells, ameliorates inflammatory pain response after formalin injection. <i>Life Sciences</i> , 2012, 91, 618-622.	2.0	18