

# Takuya Awata

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8436839/publications.pdf>

Version: 2024-02-01

117  
papers

6,741  
citations

66343

42  
h-index

64796

79  
g-index

122  
all docs

122  
docs citations

122  
times ranked

7609  
citing authors

#	ARTICLE	IF	CITATIONS
1	Japanese Type 1 Diabetes Database Study (TIDE-J): rationale and study design. <i>Diabetology International</i> , 2022, 13, 288-294.	1.4	4
2	Pancreas transplantation for type 1 diabetes in Japan: past, present and future prospects. <i>Global Health &amp; Medicine</i> , 2020, 2, 360-366.	1.4	0
3	Characteristics and clinical course of type 1 diabetes mellitus related to anti-programmed cell death-1 therapy. <i>Diabetology International</i> , 2019, 10, 58-66.	1.4	65
4	Genome-Wide Association Study Confirming a Strong Effect of HLA and Identifying Variants in <i>CSAD/Inc-ITGB7-1</i> on Chromosome 12q13.13 Associated With Susceptibility to Fulminant Type 1 Diabetes. <i>Diabetes</i> , 2019, 68, 665-675.	0.6	31
5	Diffusion-weighted magnetic resonance imaging in the pancreas of fulminant type 1 diabetes. <i>Diabetology International</i> , 2018, 9, 257-265.	1.4	2
6	Clinical considerations for use of insulin degludec/insulin aspart in Japanese patients. <i>Expert Opinion on Biological Therapy</i> , 2018, 18, 77-85.	3.1	0
7	Effects of the Activation of Three Major Hepatic Akt Substrates on Glucose Metabolism in Male Mice. <i>Endocrinology</i> , 2017, 158, 2659-2671.	2.8	8
8	A prospective multicenter study on genome wide associations to ranibizumab treatment outcome for age-related macular degeneration. <i>Scientific Reports</i> , 2017, 7, 9196.	3.3	24
9	Possible Long-Term Efficacy of Sitagliptin, a Dipeptidyl Peptidase-4 Inhibitor, for Slowly Progressive Type 1 Diabetes (SPIDDM) in the Stage of Non-Insulin-Dependency: An Open-Label Randomized Controlled Pilot Trial (SPAN-S). <i>Diabetes Therapy</i> , 2017, 8, 1123-1134.	2.5	36
10	Clinical features of cases of seroconversion of anti-glutamic acid decarboxylase antibody during the clinical course of type 2 diabetes: a nationwide survey in Japan. <i>Diabetology International</i> , 2017, 8, 306-315.	1.4	1
11	Ezetimibe Promotes Brush Border Membrane-to-Lumen Cholesterol Efflux in the Small Intestine. <i>PLoS ONE</i> , 2016, 11, e0152207.	2.5	32
12	Feminizing Adrenocortical Carcinoma with Distinct Histopathological Findings. <i>Internal Medicine</i> , 2016, 55, 3301-3307.	0.7	4
13	Risk factors for sudden death and cardiac arrest at the onset of fulminant type 1 diabetes mellitus. <i>Diabetology International</i> , 2016, 7, 281-288.	1.4	10
14	Decreased glucagon levels and decreased insulin secretion after sitagliptin versus mitiglinide administration with similar glycemic levels following an oral glucose load: a randomized crossover pharmaceutical mechanistic study. <i>Diabetology International</i> , 2016, 7, 25-33.	1.4	1
15	Clinical and Genetic Characteristics of Non-Insulin-Requiring Glutamic Acid Decarboxylase (GAD) Autoantibody-Positive Diabetes: A Nationwide Survey in Japan. <i>PLoS ONE</i> , 2016, 11, e0155643.	2.5	18
16	Influence of Treatment with Extracts of <i>Hypsizygus marmoreus</i> Mushroom on Body Composition during Obesity Development in KK-A <sup>sup</sup> y <sup>sup</sup> Mice. <i>Journal of Nutritional Science and Vitaminology</i> , 2015, 61, 96-100.	0.6	5
17	Diagnostic criteria for slowly progressive insulin-dependent (type 1) diabetes mellitus (SPIDDM) (2012): report by the Committee on Slowly Progressive Insulin-Dependent (Type 1) Diabetes Mellitus of the Japan Diabetes Society. <i>Diabetology International</i> , 2015, 6, 1-7.	1.4	44
18	The Efficacy of Vildagliptin Concomitant With Insulin Therapy in Type 2 Diabetic Subjects. <i>Journal of Clinical Medicine Research</i> , 2015, 7, 303-307.	1.2	6

#	ARTICLE	IF	CITATIONS
19	Prognostic phenotypic and genotypic factors associated with photodynamic therapy response in patients with age-related macular degeneration. <i>Clinical Ophthalmology</i> , 2014, 8, 2471.	1.8	7
20	Diagnostic criteria for acute-onset type 1 diabetes mellitus (2012): Report of the Committee of Japan Diabetes Society on the Research of Fulminant and Acute-onset Type 1 Diabetes Mellitus. <i>Journal of Diabetes Investigation</i> , 2014, 5, 115-118.	2.4	82
21	A Genome-Wide Association Study for Diabetic Retinopathy in a Japanese Population: Potential Association with a Long Intergenic Non-Coding RNA. <i>PLoS ONE</i> , 2014, 9, e111715.	2.5	81
22	Identification of Two Nickel Ion-Induced Genes, <i>NCI16</i> and <i>Pc GST1</i> , in <i>Paramecium caudatum</i> . <i>Eukaryotic Cell</i> , 2014, 13, 1181-1190.	3.4	2
23	Mediobasal hypothalamic PTEN modulates hepatic insulin resistance independently of food intake in rats. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2014, 307, E47-E60.	3.5	11
24	Distinct Cell Clusters Touching Islet Cells Induce Islet Cell Replication in Association with Over-Expression of Regenerating Gene (REG) Protein in Fulminant Type 1 Diabetes. <i>PLoS ONE</i> , 2014, 9, e95110.	2.5	24
25	A low-frequency GLIS3 variant associated with resistance to Japanese type 1 diabetes. <i>Biochemical and Biophysical Research Communications</i> , 2013, 437, 521-525.	2.1	17
26	Diagnostic criteria for acute-onset type 1 diabetes mellitus (2012). <i>Diabetology International</i> , 2013, 4, 221-225.	1.4	13
27	<i>Clostridium Butyricum</i> MIYAIRI 588 Improves High-Fat Diet-Induced Non-Alcoholic Fatty Liver Disease in Rats. <i>Digestive Diseases and Sciences</i> , 2013, 58, 3534-3544.	2.3	63
28	Predictive power of home blood pressure and clinic blood pressure in hypertensive patients with impaired glucose metabolism and diabetes. <i>Journal of Hypertension</i> , 2013, 31, 1593-1602.	0.5	45
29	Tumor-induced osteomalacia: benign tumor recurrence after two surgical resections at two different medical institutions. <i>Endocrine Practice</i> , 2013, 19, 97-101.	2.1	3
30	A Novel Splicing Variant of Peroxisome Proliferator-Activated Receptor- $\gamma$ 3 (Ppar $\gamma$ 3sv) Cooperatively Regulates Adipocyte Differentiation with Ppar $\gamma$ 2. <i>PLoS ONE</i> , 2013, 8, e65583.	2.5	20
31	Hypoglycemic effects of colestimide on type 2 diabetic patients with obesity. <i>Endocrine Journal</i> , 2012, 59, 239-246.	1.6	8
32	Report of the Committee of the Japan Diabetes Society on the Research of Fulminant and Acute-onset Type 1 Diabetes Mellitus: New Diagnostic Criteria of Fulminant Type 1 Diabetes Mellitus (2012). <i>Diabetology International</i> , 2012, 3, 179-183.	1.4	20
33	Report of the Committee of the Japan Diabetes Society on the Research of Fulminant and Acute-onset Type 1 Diabetes Mellitus: New diagnostic criteria of fulminant type 1 diabetes mellitus (2012). <i>Journal of Diabetes Investigation</i> , 2012, 3, 536-539.	2.4	187
34	Free immunoglobulin light chain: Its biology and implications in diseases. <i>Clinica Chimica Acta</i> , 2011, 412, 843-849.	1.1	89
35	Complement Factor H and High-Temperature Requirement A-1 Genotypes and Treatment Response of Age-related Macular Degeneration. <i>Ophthalmology</i> , 2011, 118, 93-100.	5.2	53
36	Associations of Cigarette Smoking But Not Serum Fatty Acids with Age-related Macular Degeneration in a Japanese Population. <i>Ophthalmology</i> , 2011, 118, 1082-1088.	5.2	49

#	ARTICLE	IF	CITATIONS
37	Association of Elastin Gene Polymorphism to Age-Related Macular Degeneration and Polypoidal Choroidal Vasculopathy. , 2011, 52, 8780.		22
38	Analysis of the HLA and non-HLA susceptibility loci in Japanese type 1 diabetes. Diabetes/Metabolism Research and Reviews, 2011, 27, 844-848.	4.0	25
39	Type 1 Diabetes and Interferon Therapy. Diabetes Care, 2011, 34, 2084-2089.	8.6	59
40	RIG-I and MDA5-Initiated Innate Immunity Linked With Adaptive Immunity Accelerates $\beta$ -Cell Death in Fulminant Type 1 Diabetes. Diabetes, 2011, 60, 884-889.	0.6	71
41	Effects of Telmisartan on Insulin Resistance in Japanese Type 2 Diabetic Patients. Internal Medicine, 2010, 49, 1843-1847.	0.7	30
42	CFH, VEGF, and PEDF genotypes and the response to intravitreal injection of bevacizumab for the treatment of age-related macular degeneration. Journal of Ocular Biology, Diseases, and Informatics, 2010, 3, 53-59.	0.2	39
43	A Promoter in the Novel Exon of hPPAR $\beta$ Directs the Circadian Expression of PPAR $\beta$ . Journal of Atherosclerosis and Thrombosis, 2010, 17, 73-83.	2.0	34
44	Vascular Endothelial Growth Factor Gene Polymorphisms in Susceptibility to Coronary Artery Disease. American Journal of Hypertension, 2010, 23, 938-939.	2.0	3
45	The Fibronectin RGD Motif Is Required for Multiple Angiogenic Events During Early Embryonic Development. Arteriosclerosis, Thrombosis, and Vascular Biology, 2010, 30, e1.	2.4	4
46	Retrospective, Observation Study: Quantitative and Qualitative Effect of Ezetimibe and HMG-CoA Reductase Inhibitors on LDL-Cholesterol: Are There Disappearance Thresholds for Small, Dense LDL and IDL?. Recent Patents on Cardiovascular Drug Discovery, 2010, 5, 143-152.	1.5	2
47	Phenotype and Genotype Characteristics of Age-related Macular Degeneration in a Japanese Population. Ophthalmology, 2010, 117, 928-938.	5.2	107
48	Activating Effect of Momordin, Extract of Bitter Melon (Momordica Charantia L.), on the Promoter of Human PPAR $\beta$ . Journal of Atherosclerosis and Thrombosis, 2009, 16, 888-892.	2.0	27
49	Adiponectin Upregulates Ferritin Heavy Chain in Skeletal Muscle Cells. Diabetes, 2009, 58, 61-70.	0.6	24
50	Rapid and Simple Profiling of Lipoproteins by Polyacrylamide-Gel Disc Electrophoresis to Determine the Heterogeneity of Low-Density Lipoproteins (LDLs) Including Small, Dense LDL. Recent Patents on Cardiovascular Drug Discovery, 2009, 4, 31-36.	1.5	12
51	Genetic Association between the Interleukin-2 Receptor- $\beta$ Gene and Mode of Onset of Type 1 Diabetes in the Japanese Population. Journal of Clinical Endocrinology and Metabolism, 2009, 94, 947-952.	3.6	24
52	Differential association of HLA with three subtypes of type 1 diabetes: fulminant, slowly progressive and acute-onset. Diabetologia, 2009, 52, 2513-2521.	6.3	123
53	Dopamine D1-like receptor antagonist, SCH23390, exhibits a preventive effect on diabetes mellitus that occurs naturally in NOD mice. Biochemical and Biophysical Research Communications, 2009, 383, 460-463.	2.1	26
54	Association of Type 1 Diabetes with Two Loci on 12q13 and 16p13 and the Influence Coexisting Thyroid Autoimmunity in Japanese. Journal of Clinical Endocrinology and Metabolism, 2009, 94, 231-235.	3.6	47

#	ARTICLE	IF	CITATIONS
55	Involvement of low adiponectin levels in impaired glucose tolerance. <i>Metabolism: Clinical and Experimental</i> , 2008, 57, 1350-1354.	3.4	8
56	Evidence for association between vitamin D receptor Bsm1 polymorphism and type 1 diabetes in Japanese. <i>Journal of Autoimmunity</i> , 2008, 30, 207-211.	6.5	49
57	SH3 domain of the phosphatidylinositol 3-kinase regulatory subunit is responsible for the formation of a sequestration complex with insulin receptor substrate-1. <i>Biochemical and Biophysical Research Communications</i> , 2008, 365, 433-438.	2.1	3
58	Cytotoxic T-Lymphocyte Associated Antigen 4 Gene Polymorphisms and Autoimmune Thyroid Disease: A Meta-Analysis. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2007, 92, 3162-3170.	3.6	162
59	Insulin Gene/IDDM2Locus in Japanese Type 1 Diabetes: Contribution of Class I Alleles and Influence of Class I Subdivision in Susceptibility to Type 1 Diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2007, 92, 1791-1795.	3.6	31
60	Promoter polymorphisms of the pigment epithelium-derived factor gene are associated with diabetic retinopathy. <i>Biochemical and Biophysical Research Communications</i> , 2007, 361, 421-426.	2.1	20
61	Homozygosity Haplotype Allows a Genomewide Search for the Autosomal Segments Shared among Patients. <i>American Journal of Human Genetics</i> , 2007, 80, 1090-1102.	6.2	59
62	Coding and Noncoding Variants in the <i>CFH</i> Gene and Cigarette Smoking Influence the Risk of Age-Related Macular Degeneration in a Japanese Population. , 2007, 48, 5315.		67
63	Systematic search for single nucleotide polymorphisms in a lymphoid tyrosine phosphatase gene ( <i>PTPN22</i> ): Association between a promoter polymorphism and type 1 diabetes in Asian populations. <i>American Journal of Medical Genetics</i> 140A:586-593 (2006). <i>American Journal of Medical Genetics, Part A</i> , 2007, 143A, 1812-1813.	1.2	3
64	Association of the HTRA1 gene variant with age-related macular degeneration in the Japanese population. <i>Journal of Human Genetics</i> , 2007, 52, 636-641.	2.3	55
65	Acarbose improves fibrinolytic activity in patients with impaired glucose tolerance. <i>Metabolism: Clinical and Experimental</i> , 2006, 55, 935-939.	3.4	19
66	Acarbose ameliorates atherogenicity of low-density lipoprotein in patients with impaired glucose tolerance. <i>Metabolism: Clinical and Experimental</i> , 2006, 55, 946-952.	3.4	33
67	LKB1, an upstream AMPK kinase, regulates glucose and lipid metabolism in cultured liver and muscle cells. <i>Biochemical and Biophysical Research Communications</i> , 2006, 351, 595-601.	2.1	50
68	Association of SUMO4, as a Candidate Gene for IDDM5, with Susceptibility to Type 1 Diabetes in Asian Populations. <i>Annals of the New York Academy of Sciences</i> , 2006, 1079, 41-46.	3.8	15
69	Systematic search for single nucleotide polymorphisms in a lymphoid tyrosine phosphatase gene ( <i>PTPN22</i> ): Association between a promoter polymorphism and type 1 diabetes in Asian populations. <i>American Journal of Medical Genetics, Part A</i> , 2006, 140A, 586-593.	1.2	141
70	Evidence for the Role of Small Ubiquitin-Like Modifier 4 as a General Autoimmunity Locus in the Japanese Population. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2006, 91, 3138-3143.	3.6	35
71	The Association of CTLA4 Polymorphism with Type 1 Diabetes Is Concentrated in Patients Complicated with Autoimmune Thyroid Disease: A Multicenter Collaborative Study in Japan. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2006, 91, 1087-1092.	3.6	85
72	Genetic variants in the calpain-10 gene and the development of type 2 diabetes in the Japanese population. <i>Journal of Human Genetics</i> , 2005, 50, 92-98.	2.3	23

#	ARTICLE	IF	CITATIONS
73	CLOCK/BMAL1 is Involved in Lipid Metabolism via Transactivation of the Peroxisome Proliferator-activated Receptor (PPAR) Response Element. <i>Journal of Atherosclerosis and Thrombosis</i> , 2005, 12, 169-174.	2.0	128
74	Regulation of adiponectin receptor gene expression in diabetic mice. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2005, 288, E876-E882.	3.5	86
75	Genetic Heterogeneity in Association of the SUMO4 M55V Variant With Susceptibility to Type 1 Diabetes. <i>Diabetes</i> , 2005, 54, 3582-3586.	0.6	52
76	Glimepiride enhances intrinsic peroxisome proliferator-activated receptor- $\beta$ activity in 3T3-L1 adipocytes. <i>Biochemical and Biophysical Research Communications</i> , 2005, 328, 484-490.	2.1	42
77	Functional VEGF C-634G polymorphism is associated with development of diabetic macular edema and correlated with macular retinal thickness in type 2 diabetes. <i>Biochemical and Biophysical Research Communications</i> , 2005, 333, 679-685.	2.1	119
78	Efficacy of glimepiride in Japanese type 2 diabetic subjects. <i>Diabetes Research and Clinical Practice</i> , 2005, 68, 250-257.	2.8	24
79	Identification of a novel WFS1 mutation (AFF344-345ins) in Japanese patients with Wolfram syndrome. <i>Diabetes Research and Clinical Practice</i> , 2005, 69, 136-141.	2.8	9
80	Endothelial Nitric Oxide Synthase Gene Is Associated With Diabetic Macular Edema in Type 2 Diabetes. <i>Diabetes Care</i> , 2004, 27, 2184-2190.	8.6	86
81	ANGPTL3 is increased in both insulin-deficient and -resistant diabetic states. <i>Biochemical and Biophysical Research Communications</i> , 2004, 317, 1075-1079.	2.1	71
82	Pro12Ala substitution in peroxisome proliferator-activated receptor- $\beta$ 2 is associated with low adiponectin concentrations in young Japanese men. <i>Metabolism: Clinical and Experimental</i> , 2004, 53, 1548-1551.	3.4	24
83	VEGF is a modifier of amyotrophic lateral sclerosis in mice and humans and protects motoneurons against ischemic death. <i>Nature Genetics</i> , 2003, 34, 383-394.	21.4	794
84	Regulation of PPAR- $\beta$ transcriptional activity in 3T3-L1 adipocytes. <i>Biochemical and Biophysical Research Communications</i> , 2003, 300, 429-436.	2.1	48
85	Apoptosis of Endothelial Cells may be Mediated by Genes of Peroxisome Proliferator-activated Receptor .GAMMA.1 (PPAR .GAMMA.1) and PPAR.ALPHA. <i>Genes.. Journal of Atherosclerosis and Thrombosis</i> , 2003, 10, 99-108.	2.0	18
86	Fibrate and Statin Synergistically Increase the Transcriptional Activities of PPAR- $\alpha$ /RXR- $\alpha$ and Decrease the Transactivation of NF- $\kappa$ B. <i>Biochemical and Biophysical Research Communications</i> , 2002, 290, 131-139.	2.1	118
87	A Common Polymorphism in the 5'-Untranslated Region of the VEGF Gene Is Associated With Diabetic Retinopathy in Type 2 Diabetes. <i>Diabetes</i> , 2002, 51, 1635-1639.	0.6	501
88	The ligands/activators for peroxisome proliferator-activated receptor- $\alpha$ (PPAR- $\alpha$ ) and PPAR- $\beta$ increase Cu <sup>2+</sup> , Zn <sup>2+</sup> -superoxide dismutase and decrease p22phox message expressions in primary endothelial cells. <i>Metabolism: Clinical and Experimental</i> , 2001, 50, 3-11.	3.4	271
89	Diabetic ketoacidosis in a case of pheochromocytoma. <i>Diabetes Research and Clinical Practice</i> , 2001, 54, 137-142.	2.8	28
90	CTLA4 gene polymorphism contributes to the mode of onset of diabetes with antigitutamic acid decarboxylase antibody in Japanese patients: genetic analysis of diabetic patients with antigitutamic acid decarboxylase antibody. <i>Diabetic Medicine</i> , 2001, 18, 726-731.	2.3	19

#	ARTICLE	IF	CITATIONS
91	Missense Variations of the Gene Responsible for Wolfram Syndrome (WFS1/wolframin) in Japanese: Possible Contribution of the Arg456His Mutation to Type 1 Diabetes as a Nonautoimmune Genetic Basis. <i>Biochemical and Biophysical Research Communications</i> , 2000, 268, 612-616.	2.1	45
92	Lack of association of the Ala45Thr variant in the BETA2/NEUROD1 with type 1 diabetes in Japanese. <i>Diabetes Research and Clinical Practice</i> , 2000, 49, 61-63.	2.8	8
93	Tyrosine hydroxylase gene microsatellite polymorphism associated with insulin resistance in depressive disorder. <i>Metabolism: Clinical and Experimental</i> , 2000, 49, 1145-1149.	3.4	62
94	Lipophilic HMG-CoA reductase inhibitor has an anti-inflammatory effect. <i>Life Sciences</i> , 2000, 67, 863-876.	4.3	273
95	Blood Pressure Control in Japanese Hypertensives with or without Type 2 Diabetes Mellitus.. <i>Hypertension Research</i> , 2000, 23, 601-605.	2.7	32
96	Troglitazone and Metformin, But Not Glibenclamide, Decrease Blood Pressure in Otsuka Long Evans Tokushima Fatty Rats. <i>Clinical and Experimental Hypertension</i> , 1999, 21, 199-211.	1.3	17
97	CTLA4 gene polymorphism correlates with the mode of onset and presence of ICA512 Ab in Japanese Type 1 diabetes. <i>Diabetes Research and Clinical Practice</i> , 1999, 46, 169-175.	2.8	42
98	Bezafibrate has an antioxidant effect: Peroxisome proliferator-activated receptor $\hat{\pm}$ is associated with Cu <sup>2+</sup> , Zn <sup>2+</sup> -superoxide dismutase in the liver. <i>Life Sciences</i> , 1998, 63, 135-144.	4.3	96
99	Expression of Peroxisome Proliferator-Activated Receptor $\hat{\pm}$ (PPAR $\hat{\pm}$ ) in Primary Cultures of Human Vascular Endothelial Cells. <i>Biochemical and Biophysical Research Communications</i> , 1998, 246, 370-374.	2.1	206
100	Effect of the Vasodilatory $\hat{I}^2$ -Blocker, Nipradilol, and Ca-Antagonist, Barnidipine, on Insulin Sensitivity in Patients with Essential Hypertension. <i>Clinical and Experimental Hypertension</i> , 1998, 20, 751-761.	1.3	22
101	Acarbose controls postprandial hyperproinsulinemia in non-insulin dependent diabetes mellitus. <i>Diabetes Research and Clinical Practice</i> , 1997, 36, 143-151.	2.8	44
102	Glucose Intolerance in Spontaneously Hypertensive and Wistar-Kyoto Rats: Enhanced Gene Expression and Synthesis of Skeletal Muscle Glucose Transporter 4.. <i>Hypertension Research</i> , 1997, 20, 279-286.	2.7	26
103	Age-dependent association of HLA-A24 in Japanese IDDM patients. <i>Diabetologia</i> , 1996, 39, 371-373.	6.3	23
104	Absence of Shared HLA Class II (DR, DQ)-Linked Genetic Basis Between IDDM and Autoimmune Thyroid Disease in Japanese. <i>Diabetes Care</i> , 1995, 18, 582-583.	8.6	7
105	Age-dependent HLA genetic heterogeneity of IDDM in Japanese patients. <i>Diabetologia</i> , 1995, 38, 748-749.	6.3	17
106	Lack of association of the insulin gene region with Type 1 (insulin-dependent) diabetes mellitus in Japanese subjects. <i>Diabetologia</i> , 1994, 37, 210-213.	6.3	12
107	Association of polymorphism in the interferon $\gamma$ gene with IDDM. <i>Diabetologia</i> , 1994, 37, 1159-1162.	6.3	62
108	Genetic markers for insulin-dependent diabetes mellitus in Japanese. <i>Diabetes Research and Clinical Practice</i> , 1994, 24, S83-S87.	2.8	27



#	ARTICLE	IF	CITATIONS
109	A Subtype of Diabetes Mellitus Associated with a Mutation of Mitochondrial DNA. New England Journal of Medicine, 1994, 330, 962-968.	27.0	555
110	HLA Class II Alleles in Japanese Patients with Graves' Disease: Weak Associations of HLA-DR and -DQ.. Endocrine Journal, 1994, 41, 599-603.	1.6	29
111	Japanese case of diabetes mellitus and . deafness with mutation in mitochondrial tRNA <sup>Leu</sup> (UUR)gene. Lancet, The, 1993, 341, 1291-1292.	13.7	33
112	Effect of islet amyloid polypeptide (IAPP/amylin) on 2-deoxyglucose uptake in mouse pancreatic acini. Diabetes Research and Clinical Practice, 1992, 15, 71-75.	2.8	3
113	Additive effect of islet amyloid polypeptide (IAPP/amylin) and insulin on 2-deoxyglucose uptake in mouse pancreatic acini. Biochemical and Biophysical Research Communications, 1991, 180, 1513-1517.	2.1	3
114	Hypervariable region 5' flanking [Leu A3]insulin gene of insulin Tochigi is different from those of insulin Wakayama I,II. Diabetes Research and Clinical Practice, 1990, 8, 183-186.	2.8	2
115	HLA DR antigens in adult-onset and juvenile-onset Japanese insulin-dependent diabetic patients. Diabetes Research and Clinical Practice, 1988, 5, 107-112.	2.8	12
116	Restriction fragment length polymorphism of the insulin gene region in Japanese diabetic and non-diabetic subjects. Diabetologia, 1985, 28, 911-913.	6.3	31
117	Endocrinology and Diabetes, 1984, 84, 346-351.	1.2	1