

Kohji Mori

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8439784/publications.pdf>

Version: 2024-02-01

30
papers

3,665
citations

394421

19
h-index

477307

29
g-index

32
all docs

32
docs citations

32
times ranked

4570
citing authors

#	ARTICLE	IF	CITATIONS
1	The <i>C9orf72</i> GGGGCC Repeat Is Translated into Aggregating Dipeptide-Repeat Proteins in FTL/ALS. <i>Science</i> , 2013, 339, 1335-1338.	12.6	1,095
2	Bidirectional transcripts of the expanded <i>C9orf72</i> hexanucleotide repeat are translated into aggregating dipeptide repeat proteins. <i>Acta Neuropathologica</i> , 2013, 126, 881-893.	7.7	427
3	hnRNP A3 binds to GGGGCC repeats and is a constituent of p62-positive/TDP43-negative inclusions in the hippocampus of patients with <i>C9orf72</i> mutations. <i>Acta Neuropathologica</i> , 2013, 125, 413-423.	7.7	302
4	<i>C9orf72</i> FTL/ALS-associated Gly-Ala dipeptide repeat proteins cause neuronal toxicity and Unc119 sequestration. <i>Acta Neuropathologica</i> , 2014, 128, 485-503.	7.7	300
5	Dipeptide repeat protein pathology in <i>C9ORF72</i> mutation cases: clinico-pathological correlations. <i>Acta Neuropathologica</i> , 2013, 126, 859-879.	7.7	298
6	Effects of norepinephrine on rat cultured microglial cells that express β_1 , β_2 , β_1 and β_2 adrenergic receptors. <i>Neuropharmacology</i> , 2002, 43, 1026-1034.	4.1	184
7	Common pathobiochemical hallmarks of progranulin-associated frontotemporal lobar degeneration and neuronal ceroid lipofuscinosis. <i>Acta Neuropathologica</i> , 2014, 127, 845-60.	7.7	156
8	TREM2 deficiency reduces the efficacy of immunotherapeutic amyloid clearance. <i>EMBO Molecular Medicine</i> , 2016, 8, 992-1004.	6.9	144
9	β -Secretase Modulators and Presenilin 1 Mutants Act Differently on Presenilin/ β -Secretase Function to Cleave $A\beta_{42}$ and $A\beta_{43}$. <i>Cell Reports</i> , 2013, 3, 42-51.	6.4	110
10	Renal function is associated with blood neurofilament light chain level in older adults. <i>Scientific Reports</i> , 2020, 10, 20350.	3.3	96
11	Microglia, a potential source of neurons, astrocytes, and oligodendrocytes. <i>Glia</i> , 2004, 45, 96-104.	4.9	92
12	The 28-amino acid form of an APLP1-derived $A\beta$ -like peptide is a surrogate marker for $A\beta_{42}$ production in the central nervous system. <i>EMBO Molecular Medicine</i> , 2009, 1, 223-235.	6.9	72
13	Antibodies inhibit transmission and aggregation of <i>C9orf72</i> poly(GA) dipeptide repeat proteins. <i>EMBO Molecular Medicine</i> , 2017, 9, 687-702.	6.9	70
14	A protein quality control pathway regulated by linear ubiquitination. <i>EMBO Journal</i> , 2019, 38, .	7.8	63
15	Poly-glycine-alanine exacerbates <i>C9orf72</i> repeat expansion-mediated DNA damage via sequestration of phosphorylated ATM and loss of nuclear hnRNPA3. <i>Acta Neuropathologica</i> , 2020, 139, 99-118.	7.7	49
16	Reduced hnRNPA3 increases <i>C9orf72</i> repeat RNA levels and dipeptide-repeat protein deposition. <i>EMBO Reports</i> , 2016, 17, 1314-1325.	4.5	39
17	Treatment of delirium with ramelteon: initial experience in three patients. <i>General Hospital Psychiatry</i> , 2011, 33, 407-409.	2.4	30
18	Two populations of microglial cells isolated from rat primary mixed glial cultures. <i>Journal of Neuroscience Research</i> , 2003, 73, 22-30.	2.9	20

#	ARTICLE	IF	CITATIONS
19	Heterogeneous ribonuclear protein A3 (hnRNP A3) is present in dipeptide repeat protein containing inclusions in Frontotemporal Lobar Degeneration and Motor Neurone disease associated with expansions in C9orf72 gene. <i>Acta Neuropathologica Communications</i> , 2017, 5, 31.	5.2	20
20	l-Serine-mediated release of apolipoprotein E and lipids from microglial cells. <i>Experimental Neurology</i> , 2004, 185, 220-231.	4.1	18
21	The <sc>RNA</sc> exosome complex degrades expanded hexanucleotide repeat <sc>RNA</sc> in <i>C9orf72</i> <sc>FTLD</sc> / <sc>ALS</sc>. <i>EMBO Journal</i> , 2020, 39, e102700.	7.8	18
22	The porphyrin TMPyP4 inhibits elongation during the noncanonical translation of the FTLD/ALS-associated GGGGCC repeat in the C9orf72 gene. <i>Journal of Biological Chemistry</i> , 2021, 297, 101120.	3.4	17
23	The production ratios of AICD μ 51 and A β 242 by intramembrane proteolysis of β APP do not always change in parallel. <i>Psychogeriatrics</i> , 2010, 10, 117-123.	1.2	11
24	Destruxin E Decreases Beta-Amyloid Generation by Reducing Colocalization of Beta-Amyloid-Cleaving Enzyme 1 and Beta-Amyloid Protein Precursor. <i>Neurodegenerative Diseases</i> , 2009, 6, 230-239.	1.4	9
25	Repurposing bromocriptine for A β 2 metabolism in Alzheimer's disease (REBRAND) study: randomised placebo-controlled double-blind comparative trial and open-label extension trial to investigate the safety and efficacy of bromocriptine in Alzheimer's disease with presenilin 1 (PSEN1) mutations. <i>BMJ Open</i> , 2021, 11, e051343.	1.9	9
26	Two Neuropsychiatric Cases Seropositive for Bornavirus Improved by Ribavirin. <i>Japanese Journal of Infectious Diseases</i> , 2018, 71, 338-342.	1.2	5
27	Biological basis and psychiatric symptoms in frontotemporal dementia. <i>Psychiatry and Clinical Neurosciences</i> , 2022, 76, 351-360.	1.8	5
28	Macrophage colony stimulating factor is associated with excretion of amyloid β 2 peptides from cerebrospinal fluid to peripheral blood. <i>Psychogeriatrics</i> , 2008, 8, 188-195.	1.2	3
29	Plasma <sc>NfL</sc> is associated with mild cognitive decline in patients with diabetes. <i>Psychogeriatrics</i> , 2022, 22, 353-359.	1.2	3
30	Levels of the surrogate marker for A β 242 (i.e., APL1 β) in CSF of sporadic Alzheimer disease patients increase before the onset of its clinical symptoms. <i>Neuroscience Research</i> , 2010, 68, e67.	1.9	0