

Paul Saftig

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8443879/publications.pdf>

Version: 2024-02-01

205
papers

32,158
citations

4370

86
h-index

4203

174
g-index

212
all docs

212
docs citations

212
times ranked

33583
citing authors

#	ARTICLE	IF	CITATIONS
1	LAMP2 regulates autophagy in the thymic epithelium and thymic stroma-dependent CD4 T cell development. <i>Autophagy</i> , 2023, 19, 426-439.	4.3	12
2	S-palmitoylation determines TMEM55B-dependent positioning of lysosomes. <i>Journal of Cell Science</i> , 2022, 135, .	1.2	3
3	Ectodomain shedding by ADAM proteases as a central regulator in kidney physiology and disease. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2022, 1869, 119165.	1.9	6
4	Lung emphysema and impaired macrophage elastase clearance in mucolipin 3 deficient mice. <i>Nature Communications</i> , 2022, 13, 318.	5.8	25
5	Intravitreal gene therapy restores the autophagy-lysosomal pathway and attenuates retinal degeneration in cathepsin D-deficient mice. <i>Neurobiology of Disease</i> , 2022, 164, 105628.	2.1	8
6	Inhibition of ADAM17 impairs endothelial cell necroptosis and blocks metastasis. <i>Journal of Experimental Medicine</i> , 2022, 219, .	4.2	35
7	Recombinant pro-CTSD (cathepsin D) enhances SNCA/±-Synuclein degradation in ±-Synucleinopathy models. <i>Autophagy</i> , 2022, 18, 1127-1151.	4.3	20
8	Phagosomal signalling of the C-type lectin receptor Dectin-1 is terminated by intramembrane proteolysis. <i>Nature Communications</i> , 2022, 13, 1880.	5.8	17
9	How Lysosomes Sense, Integrate, and Cope with Stress. <i>Trends in Biochemical Sciences</i> , 2021, 46, 97-112.	3.7	84
10	ADAM10 hyperactivation acts on piccolo to deplete synaptic vesicle stores in Huntingtonâ€™s disease. <i>Human Molecular Genetics</i> , 2021, 30, 1175-1187.	1.4	11
11	ADAM10-Mediated Ectodomain Shedding Is an Essential Driver of Podocyte Damage. <i>Journal of the American Society of Nephrology: JASN</i> , 2021, 32, 1389-1408.	3.0	7
12	Rapid and Progressive Loss of Multiple Retinal Cell Types in Cathepsin D-Deficient Miceâ€”An Animal Model of CLN10 Disease. <i>Cells</i> , 2021, 10, 696.	1.8	10
13	GPR37 is processed in the Nâ€”terminal ectodomain by ADAM10 and furin. <i>FASEB Journal</i> , 2021, 35, e21654.	0.2	11
14	Posttranslational modifications by ADAM10 shape myeloid antigen-presenting cell homeostasis in the splenic marginal zone. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, .	3.3	7
15	A09â€”ADAM10 activity at the huntingtonâ€™s disease presynapse. , 2021, , .		1
16	Analysis of cathepsin B and cathepsin L treatment to clear toxic lysosomal protein aggregates in neuronal ceroid lipofuscinosis. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2021, 1867, 166205.	1.8	10
17	Ligands binding to the prion protein induce its proteolytic release with therapeutic potential in neurodegenerative proteinopathies. <i>Science Advances</i> , 2021, 7, eabj1826.	4.7	18
18	Enzyme replacement therapy with recombinant pro-CTSD (cathepsin D) corrects defective proteolysis and autophagy in neuronal ceroid lipofuscinosis. <i>Autophagy</i> , 2020, 16, 811-825.	4.3	70

#	ARTICLE	IF	CITATIONS
19	LAMP-2 Is Involved in Surface Expression of RANKL of Osteoblasts In Vitro. <i>International Journal of Molecular Sciences</i> , 2020, 21, 6110.	1.8	7
20	The FTLD Risk Factor TMEM106B Regulates the Transport of Lysosomes at the Axon Initial Segment of Motoneurons. <i>Cell Reports</i> , 2020, 30, 3506-3519.e6.	2.9	47
21	Cholesterol Handling in Lysosomes and Beyond. <i>Trends in Cell Biology</i> , 2020, 30, 452-466.	3.6	97
22	The tetraspanin Tspan15 is an essential subunit of an ADAM10 scissor complex. <i>Journal of Biological Chemistry</i> , 2020, 295, 12822-12839.	1.6	31
23	Mepripin \hat{I}^2 induces activities of A disintegrin and metalloproteinases 9, 10, and 17 by specific prodomain cleavage. <i>FASEB Journal</i> , 2019, 33, 11925-11940.	0.2	18
24	Lysosomal integral membrane protein-2 (LIMP-2/SCARB2) is involved in lysosomal cholesterol export. <i>Nature Communications</i> , 2019, 10, 3521.	5.8	99
25	Presynaptic Endosomal Cathepsin D Regulates the Biogenesis of GABAergic Synaptic Vesicles. <i>Cell Reports</i> , 2019, 28, 1015-1028.e5.	2.9	17
26	Ubiquitin C-terminal hydrolase L1 (UCH-L1) loss causes neurodegeneration by altering protein turnover in the first postnatal weeks. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 7963-7972.	3.3	36
27	The intramembrane protease $\langle scp \rangle$ SPPL $\langle /scp \rangle$ 2c promotes male germ cell development by cleaving \hat{A} phospholamban. <i>EMBO Reports</i> , 2019, 20, .	2.0	27
28	Genetic LAMP2 deficiency accelerates the age-associated formation of basal laminar deposits in the retina. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 23724-23734.	3.3	54
29	Lysosomal storage disorders \hat{A} challenges, concepts and avenues for therapy: beyond rare diseases. <i>Journal of Cell Science</i> , 2019, 132, jcs221739.	1.2	141
30	Inhibiting pathologically active ADAM10 rescues synaptic and cognitive decline in Huntington \hat{A} ™s disease. <i>Journal of Clinical Investigation</i> , 2019, 129, 2390-2403.	3.9	38
31	The lysosomal transporter MFSD1 is essential for liver homeostasis and critically depends on its accessory subunit GLMP. <i>ELife</i> , 2019, 8, .	2.8	23
32	Unconventional Trafficking of Mammalian Phospholipase D3 to Lysosomes. <i>Cell Reports</i> , 2018, 22, 1040-1053.	2.9	31
33	Structural and mechanistic aspects influencing the ADAM10-mediated shedding of the prion protein. <i>Molecular Neurodegeneration</i> , 2018, 13, 18.	4.4	45
34	In vivo regulation of the A disintegrin and metalloproteinase 10 (ADAM10) by the tetraspanin 15. <i>Cellular and Molecular Life Sciences</i> , 2018, 75, 3251-3267.	2.4	37
35	Intracellular compartments of pathogens: Highways to hell or stairways to heaven?. <i>International Journal of Medical Microbiology</i> , 2018, 308, 1-2.	1.5	0
36	Vacuolar ATPase in phago(lyso)some biology. <i>International Journal of Medical Microbiology</i> , 2018, 308, 58-67.	1.5	37

#	ARTICLE	IF	CITATIONS
37	Quantitative Proteome Analysis of Mouse Liver Lysosomes Provides Evidence for Mannose 6-phosphate-independent Targeting Mechanisms of Acid Hydrolases in Mucopolipidosis II. <i>Molecular and Cellular Proteomics</i> , 2017, 16, 438-450.	2.5	30
38	Disruption of the vacuolar-type H ⁺ -ATPase complex in liver causes MTORC1-independent accumulation of autophagic vacuoles and lysosomes. <i>Autophagy</i> , 2017, 13, 670-685.	4.3	19
39	Sequestration of cholesterol within the host late endocytic pathway restricts liver-stage <i>Plasmodium</i> development. <i>Molecular Biology of the Cell</i> , 2017, 28, 726-735.	0.9	37
40	Progranulin functions as a cathepsin D chaperone to stimulate axonal outgrowth in vivo. <i>Human Molecular Genetics</i> , 2017, 26, 2850-2863.	1.4	111
41	The metalloproteinase ADAM10: A useful therapeutic target?. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2017, 1864, 2071-2081.	1.9	111
42	The Influence of MHC Class II on B Cell Defects Induced by Invariant Chain/CD74 N-Terminal Fragments. <i>Journal of Immunology</i> , 2017, 199, 172-185.	0.4	11
43	Long-term enzyme replacement therapy improves neurocognitive functioning and hippocampal synaptic plasticity in immune-tolerant alpha-mannosidosis mice. <i>Neurobiology of Disease</i> , 2017, 106, 255-268.	2.1	8
44	Absence of RNase H2 triggers generation of immunogenic micronuclei removed by autophagy. <i>Human Molecular Genetics</i> , 2017, 26, 3960-3972.	1.4	160
45	Lysosomal integral membrane protein-2 as a phospholipid receptor revealed by biophysical and cellular studies. <i>Nature Communications</i> , 2017, 8, 1908.	5.8	43
46	Diverse functions of the prion protein " Does proteolytic processing hold the key?. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2017, 1864, 2128-2137.	1.9	60
47	Tetraspanin 3: A central endocytic membrane component regulating the expression of ADAM10, presenilin and the amyloid precursor protein. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2017, 1864, 217-230.	1.9	26
48	Functional characterization of the lysosomal membrane protein TMEM192 in mice. <i>Oncotarget</i> , 2017, 8, 43635-43652.	0.8	8
49	The Emerging Role of Tetraspanins in the Proteolytic Processing of the Amyloid Precursor Protein. <i>Frontiers in Molecular Neuroscience</i> , 2016, 9, 149.	1.4	40
50	Dissecting the role of ADAM10 as a mediator of <i>Staphylococcus aureus</i> Î±-toxin action. <i>Biochemical Journal</i> , 2016, 473, 1929-1940.	1.7	33
51	Substrate determinants of signal peptide peptidase-like 2a (SPPL2a)-mediated intramembrane proteolysis of the invariant chain CD74. <i>Biochemical Journal</i> , 2016, 473, 1405-1422.	1.7	24
52	Intramembrane proteolysis within lysosomes. <i>Ageing Research Reviews</i> , 2016, 32, 51-64.	5.0	14
53	Turn up the lysosome. <i>Nature Cell Biology</i> , 2016, 18, 1025-1027.	4.6	74
54	Parkinson's disease: acid-glucocerebrosidase activity and alpha-synuclein clearance. <i>Journal of Neurochemistry</i> , 2016, 139, 198-215.	2.1	59

#	ARTICLE	IF	CITATIONS
55	LAMP proteins account for the maturation delay during the establishment of the <i>Coxiella burnetii</i> -containing vacuole. <i>Cellular Microbiology</i> , 2016, 18, 181-194.	1.1	34
56	Characterization of the complex formed by β -glucocerebrosidase and the lysosomal integral membrane protein type-2. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, 3791-3796.	3.3	45
57	Impaired Lysosomal Integral Membrane Protein 2-dependent Peroxiredoxin 6 Delivery to Lamellar Bodies Accounts for Altered Alveolar Phospholipid Content in Adaptor Protein-3-deficient pearl Mice. <i>Journal of Biological Chemistry</i> , 2016, 291, 8414-8427.	1.6	24
58	ADAM17 controls IL-6 signaling by cleavage of the murine IL-6R α from the cell surface of leukocytes during inflammatory responses. <i>Journal of Leukocyte Biology</i> , 2016, 99, 749-760.	1.5	49
59	A disintegrin and metalloprotease 10 (ADAM10) is a central regulator of murine liver tissue homeostasis. <i>Oncotarget</i> , 2016, 7, 17431-17441.	0.8	17
60	Systematic substrate identification indicates a central role for the metalloprotease ADAM10 in axon targeting and synapse function. <i>ELife</i> , 2016, 5, .	2.8	124
61	Chronic enzyme replacement therapy ameliorates neuropathology in alpha β -mannosidosis mice. <i>Annals of Clinical and Translational Neurology</i> , 2015, 2, 987-1001.	1.7	8
62	Mannose 6-phosphate-independent Lysosomal Sorting of LIMP-2. <i>Traffic</i> , 2015, 16, 1127-1136.	1.3	23
63	TIMP-1 signaling via CD63 triggers granulopoiesis and neutrophilia in mice. <i>Haematologica</i> , 2015, 100, 1005-13.	1.7	37
64	Proteases at work: cues for understanding neural development and degeneration. <i>Frontiers in Molecular Neuroscience</i> , 2015, 8, 13.	1.4	16
65	β -Secretase BACE1 Regulates Hippocampal and Reconstituted M-Currents in a β -Subunit-Like Fashion. <i>Journal of Neuroscience</i> , 2015, 35, 3298-3311.	1.7	34
66	Vacuolar ATPase in Phagosome-Lysosome Fusion. <i>Journal of Biological Chemistry</i> , 2015, 290, 14166-14180.	1.6	75
67	Lysosomal integral membrane protein type-2 (LIMP-2/SCARB2) is a substrate of cathepsin-F, a cysteine protease mutated in type-B-Kufs-disease. <i>Biochemical and Biophysical Research Communications</i> , 2015, 457, 334-340.	1.0	13
68	Myeloid A Disintegrin and Metalloproteinase Domain 10 Deficiency Modulates Atherosclerotic Plaque Composition by Shifting the Balance from Inflammation toward Fibrosis. <i>American Journal of Pathology</i> , 2015, 185, 1145-1155.	1.9	46
69	LAMP-2 deficiency leads to hippocampal dysfunction but normal clearance of neuronal substrates of chaperone-mediated autophagy in a mouse model for Danon disease. <i>Acta Neuropathologica Communications</i> , 2015, 3, 6.	2.4	63
70	The alpha secretase ADAM10: A metalloprotease with multiple functions in the brain. <i>Progress in Neurobiology</i> , 2015, 135, 1-20.	2.8	190
71	Processing of CD74 by the Intramembrane Protease SPPL2a Is Critical for B Cell Receptor Signaling in Transitional B Cells. <i>Journal of Immunology</i> , 2015, 195, 1548-1563.	0.4	25
72	Cathepsin D deficiency induces oxidative damage in brain pericytes and impairs the blood-brain barrier. <i>Molecular and Cellular Neurosciences</i> , 2015, 64, 51-60.	1.0	21

#	ARTICLE	IF	CITATIONS
73	The sheddase ADAM10 is a potent modulator of prion disease. <i>ELife</i> , 2015, 4, .	2.8	66
74	ADAM10: Alzheimer a-Sekretase und neurobiologischer Regulator. <i>E-Neuroforum</i> , 2014, 20, 212-221.	0.2	0
75	The Intramembrane Proteases Signal Peptide Peptidase-Like 2a and 2b Have Distinct Functions <i><i>In Vivo</i></i> . <i>Molecular and Cellular Biology</i> , 2014, 34, 1398-1411.	1.1	30
76	The endolysosomal cysteine cathepsins L and K are involved in macrophage-mediated clearance of <i><i>Staphylococcus aureus</i></i> and the concomitant cytokine induction. <i>FASEB Journal</i> , 2014, 28, 162-175.	0.2	44
77	ADAM metalloproteases promote a developmental switch in responsiveness to the axonal repellent Sema3A. <i>Nature Communications</i> , 2014, 5, 4058.	5.8	39
78	LIMP-2 expression is critical for β -glucocerebrosidase activity and β -synuclein clearance. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 15573-15578.	3.3	109
79	Signal-peptide-peptidase-like 2a is required for CD74 intramembrane proteolysis in human B cells. <i>Biochemical and Biophysical Research Communications</i> , 2014, 451, 48-53.	1.0	19
80	Lassa virus entry requires a trigger-induced receptor switch. <i>Science</i> , 2014, 344, 1506-1510.	6.0	251
81	High susceptibility to fatty liver disease in two-pore channel 2-deficient mice. <i>Nature Communications</i> , 2014, 5, 4699.	5.8	164
82	Regulated Proteolysis of NOTCH2 and NOTCH3 Receptors by ADAM10 and Presenilins. <i>Molecular and Cellular Biology</i> , 2014, 34, 2822-2832.	1.1	72
83	Natural history of alpha mannosidosis a longitudinal study. <i>Orphanet Journal of Rare Diseases</i> , 2013, 8, 88.	1.2	50
84	Postnatal Disruption of the Disintegrin/Metalloproteinase ADAM10 in Brain Causes Epileptic Seizures, Learning Deficits, Altered Spine Morphology, and Defective Synaptic Functions. <i>Journal of Neuroscience</i> , 2013, 33, 12915-12928.	1.7	107
85	Parallel regulation of renin and lysosomal integral membrane protein 2 in renin-producing cells: further evidence for a lysosomal nature of renin secretory vesicles. <i>Pflugers Archiv European Journal of Physiology</i> , 2013, 465, 895-905.	1.3	9
86	Structure of LIMP-2 provides functional insights with implications for SR-BI and CD36. <i>Nature</i> , 2013, 504, 172-176.	13.7	226
87	Lysosomal Membrane Proteins and Their Central Role in Physiology. <i>Traffic</i> , 2013, 14, 739-748.	1.3	175
88	Killing from the inside. <i>Nature</i> , 2013, 502, 312-313.	13.7	79
89	Regulation of adult hematopoiesis by the a disintegrin and metalloproteinase 10 (ADAM10). <i>Biochemical and Biophysical Research Communications</i> , 2013, 442, 234-241.	1.0	13
90	Extracellular cathepsin K exerts antimicrobial activity and is protective against chronic intestinal inflammation in mice. <i>Gut</i> , 2013, 62, 520-530.	6.1	31

#	ARTICLE	IF	CITATIONS
91	Cathepsin F mutations cause Type B Kufs disease, an adult-onset neuronal ceroid lipofuscinosis. <i>Human Molecular Genetics</i> , 2013, 22, 1417-1423.	1.4	105
92	The intramembrane protease SPPL2a promotes B cell development and controls endosomal traffic by cleavage of the invariant chain. <i>Journal of Experimental Medicine</i> , 2013, 210, 41-58.	4.2	100
93	Mannose 6 Dephosphorylation of Lysosomal Proteins Mediated by Acid Phosphatases Acp2 and Acp5. <i>Molecular and Cellular Biology</i> , 2012, 32, 774-782.	1.1	43
94	The lysosomal polypeptide transporter TAPL is stabilized by the interaction with LAMP-1 and LAMP-2. <i>Journal of Cell Science</i> , 2012, 125, 4230-40.	1.2	39
95	Ectodomain shedding and ADAMs in development. <i>Development (Cambridge)</i> , 2012, 139, 3693-3709.	1.2	211
96	Activity-Dependent Proteolytic Cleavage of Neuroligin-1. <i>Neuron</i> , 2012, 76, 410-422.	3.8	179
97	Tetraspanin15 regulates cellular trafficking and activity of the ectodomain sheddase ADAM10. <i>Cellular and Molecular Life Sciences</i> , 2012, 69, 2919-2932.	2.4	99
98	Physiological functions of the amyloid precursor protein secretases ADAM10, BACE1, and Presenilin. <i>Experimental Brain Research</i> , 2012, 217, 331-341.	0.7	52
99	A Critical Histidine Residue Within <i>LIMP2</i> Mediates pH-Sensitive Binding to Its Ligand β -Glucocerebrosidase. <i>Traffic</i> , 2012, 13, 1113-1123.	1.3	41
100	Sensitivity to Lysosome-Dependent Cell Death Is Directly Regulated by Lysosomal Cholesterol Content. <i>PLoS ONE</i> , 2012, 7, e50262.	1.1	66
101	The disintegrin/metalloproteinase Adam10 is essential for epidermal integrity and Notch-mediated signaling. <i>Development (Cambridge)</i> , 2011, 138, 495-505.	1.2	130
102	The Tetraspanin CD63 Regulates ESCRT-Independent and -Dependent Endosomal Sorting during Melanogenesis. <i>Developmental Cell</i> , 2011, 21, 708-721.	3.1	687
103	Disrupted in renal carcinoma 2 (DIRC2), a novel transporter of the lysosomal membrane, is proteolytically processed by cathepsin L. <i>Biochemical Journal</i> , 2011, 439, 113-128.	1.7	29
104	Cerebellar Alterations and Gait Defects as Therapeutic Outcome Measures for Enzyme Replacement Therapy in α -Mannosidosis. <i>Journal of Neuropathology and Experimental Neurology</i> , 2011, 70, 83-94.	0.9	22
105	CD63 is an essential cofactor to leukocyte recruitment by endothelial P-selectin. <i>Blood</i> , 2011, 118, 4265-4273.	0.6	79
106	Deletion of Adam10 in endothelial cells leads to defects in organ-specific vascular structures. <i>Blood</i> , 2011, 118, 1163-1174.	0.6	69
107	Two dileucine motifs mediate late endosomal/lysosomal targeting of transmembrane protein 192 (TMEM192) and a C-terminal cysteine residue is responsible for disulfide bond formation in TMEM192 homodimers. <i>Biochemical Journal</i> , 2011, 434, 219-231.	1.7	25
108	Role for LAMP-2 in endosomal cholesterol transport. <i>Journal of Cellular and Molecular Medicine</i> , 2011, 15, 280-295.	1.6	70

#	ARTICLE	IF	CITATIONS
109	The α 5 β 1 Disintegrin And Metalloproteases ADAM10 and ADAM17: Novel drug targets with therapeutic potential?. <i>European Journal of Cell Biology</i> , 2011, 90, 527-535.	1.6	256
110	Signal-peptide-peptidase-like 2a (SPPL2a) is targeted to lysosomes/late endosomes by a tyrosine motif in its C-terminal tail. <i>FEBS Letters</i> , 2011, 585, 2951-2957.	1.3	39
111	β -Site APP-cleaving enzyme 1 (BACE1) cleaves cerebellar Na ⁺ channel β 4-subunit and promotes Purkinje cell firing by slowing the decay of resurgent Na ⁺ current. <i>Pflugers Archiv European Journal of Physiology</i> , 2011, 461, 355-371.	1.3	27
112	Lack of α 5-disintegrin-and-metalloproteinase ADAM10 leads to intracellular accumulation and loss of shedding of the cellular prion protein in vivo. <i>Molecular Neurodegeneration</i> , 2011, 6, 36.	4.4	93
113	Neuronal Brain-derived Neurotrophic Factor Is Synthesized in Excess, with Levels Regulated by Sortilin-mediated Trafficking and Lysosomal Degradation. <i>Journal of Biological Chemistry</i> , 2011, 286, 29556-29567.	1.6	91
114	Sphingolipid Storage Affects Autophagic Metabolism of the Amyloid Precursor Protein and Promotes A β Generation. <i>Journal of Neuroscience</i> , 2011, 31, 1837-1849.	1.7	82
115	The proteome of lysosomes. <i>Proteomics</i> , 2010, 10, 4053-4076.	1.3	188
116	Disease-causing mutations within the lysosomal integral membrane protein type 2 (LIMP-2) reveal the nature of binding to its ligand β -glucocerebrosidase. <i>Human Molecular Genetics</i> , 2010, 19, 563-572.	1.4	86
117	Molecular characterisation of β -transmembrane protein 192 TM (TMEM192), a novel protein of the lysosomal membrane. <i>Biological Chemistry</i> , 2010, 391, 695-704.	1.2	43
118	Activity-dependent β -Cleavage of Nectin-1 Is Mediated by A Disintegrin and Metalloprotease 10 (ADAM10). <i>Journal of Biological Chemistry</i> , 2010, 285, 22919-22926.	1.6	46
119	Critical role of the disintegrin metalloprotease ADAM17 for intestinal inflammation and regeneration in mice. <i>Journal of Experimental Medicine</i> , 2010, 207, 1617-1624.	4.2	286
120	The Disintegrin/Metalloproteinase ADAM10 Is Essential for the Establishment of the Brain Cortex. <i>Journal of Neuroscience</i> , 2010, 30, 4833-4844.	1.7	327
121	ADAM17 is regulated by a rapid and reversible mechanism that controls access to its catalytic site. <i>Journal of Cell Science</i> , 2010, 123, 3913-3922.	1.2	165
122	Lysosomal membrane proteins: life between acid and neutral conditions. <i>Biochemical Society Transactions</i> , 2010, 38, 1420-1423.	1.6	73
123	CNS-Expressed Cathepsin D Prevents Lymphopenia in a Murine Model of Congenital Neuronal Ceroid Lipofuscinosis. <i>American Journal of Pathology</i> , 2010, 177, 271-279.	1.9	42
124	Alcadein Cleavages by Amyloid β -Precursor Protein (APP) β - and γ -Secretases Generate Small Peptides, p3-Alcs, Indicating Alzheimer Disease-related γ -Secretase Dysfunction. <i>Journal of Biological Chemistry</i> , 2009, 284, 36024-36033.	1.6	46
125	Cytoplasmic Relaxation of Active Eph Controls Ephrin Shedding by ADAM10. <i>PLoS Biology</i> , 2009, 7, e1000215.	2.6	72
126	ADAMs 10 and 17 Represent Differentially Regulated Components of a General Shedding Machinery for Membrane Proteins Such as Transforming Growth Factor β , L-Selectin, and Tumor Necrosis Factor β . <i>Molecular Biology of the Cell</i> , 2009, 20, 1785-1794.	0.9	230

#	ARTICLE	IF	CITATIONS
127	Deficiency of the Tetraspanin CD63 Associated with Kidney Pathology but Normal Lysosomal Function. <i>Molecular and Cellular Biology</i> , 2009, 29, 1083-1094.	1.1	99
128	Non-proteolytic effect of β -site APP-cleaving enzyme 1 (BACE1) on sodium channel function. <i>Neurobiology of Disease</i> , 2009, 33, 282-289.	2.1	39
129	Autophagy: A lysosomal degradation pathway with a central role in health and disease. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2009, 1793, 664-673.	1.9	581
130	Klotho is a substrate for β -casein and β -secretase. <i>FEBS Letters</i> , 2009, 583, 3221-3224.	1.3	215
131	Lysosome biogenesis and lysosomal membrane proteins: trafficking meets function. <i>Nature Reviews Molecular Cell Biology</i> , 2009, 10, 623-635.	16.1	1,320
132	The α -Disintegrin And Metalloprotease (ADAM) family of sheddases: Physiological and cellular functions. <i>Seminars in Cell and Developmental Biology</i> , 2009, 20, 126-137.	2.3	356
133	Regulated intramembrane proteolysis: A story about sheddases and I-CliPs. <i>Seminars in Cell and Developmental Biology</i> , 2009, 20, 125.	2.3	5
134	Cathepsin D expression level affects alpha-synuclein processing, aggregation, and toxicity in vivo. <i>Molecular Brain</i> , 2009, 2, 5.	1.3	232
135	ADAM10, the Rate-limiting Protease of Regulated Intramembrane Proteolysis of Notch and Other Proteins, Is Processed by ADAMS-9, ADAMS-15, and the β -Secretase. <i>Journal of Biological Chemistry</i> , 2009, 284, 11738-11747.	1.6	161
136	Array-Based Gene Discovery with Three Unrelated Subjects Shows SCARB2/LIMP-2 Deficiency Causes Myoclonus Epilepsy and Glomerulosclerosis. <i>American Journal of Human Genetics</i> , 2008, 82, 673-684.	2.6	230
137	ADAM10-Mediated E-Cadherin Release Is Regulated by Proinflammatory Cytokines and Modulates Keratinocyte Cohesion in Eczematous Dermatitis. <i>Journal of Investigative Dermatology</i> , 2008, 128, 1737-1746.	0.3	79
138	ADAM10 Regulates Endothelial Permeability and T-Cell Transmigration by Proteolysis of Vascular Endothelial Cadherin. <i>Circulation Research</i> , 2008, 102, 1192-1201.	2.0	264
139	LAMP-2: A control step for phagosome and autophagosome maturation. <i>Autophagy</i> , 2008, 4, 510-512.	4.3	190
140	A soluble form of the receptor for advanced glycation endproducts (RAGE) is produced by proteolytic cleavage of the membrane-bound form by the sheddase a disintegrin and metalloprotease 10 (ADAM10). <i>FASEB Journal</i> , 2008, 22, 3716-3727.	0.2	483
141	Regulated Intramembrane Proteolysis of Bri2 (Itm2b) by ADAM10 and SPPL2a/SPPL2b. <i>Journal of Biological Chemistry</i> , 2008, 283, 1644-1652.	1.6	132
142	Reversal of peripheral and central neural storage and ataxia after recombinant enzyme replacement therapy in α -mannosidosis mice. <i>Human Molecular Genetics</i> , 2008, 17, 3437-3445.	1.4	60
143	Impaired Phagosomal Maturation in Neutrophils Leads to Periodontitis in Lysosomal-Associated Membrane Protein-2 Knockout Mice. <i>Journal of Immunology</i> , 2008, 180, 475-482.	0.4	67
144	Synaptic Changes in the Thalamocortical System of Cathepsin D-Deficient Mice. <i>Journal of Neuropathology and Experimental Neurology</i> , 2008, 67, 16-29.	0.9	79

#	ARTICLE	IF	CITATIONS
145	Lysosomal integral membrane protein 2 is a novel component of the cardiac intercalated disc and vital for load-induced cardiac myocyte hypertrophy. <i>Journal of Experimental Medicine</i> , 2007, 204, 1227-1235.	4.2	37
146	Substrate Selectivity of Epidermal Growth Factor-Receptor Ligand Sheddases and their Regulation by Phorbol Esters and Calcium Influx. <i>Molecular Biology of the Cell</i> , 2007, 18, 176-188.	0.9	276
147	Regulated Shedding of Transmembrane Chemokines by the Disintegrin and Metalloproteinase 10 Facilitates Detachment of Adherent Leukocytes. <i>Journal of Immunology</i> , 2007, 178, 8064-8072.	0.4	151
148	LIMP-2 Is a Receptor for Lysosomal Mannose-6-Phosphate-Independent Targeting of β -2-Glucocerebrosidase. <i>Cell</i> , 2007, 131, 770-783.	13.5	428
149	ADAM10 Inhibition of Human CD30 Shedding Increases Specificity of Targeted Immunotherapy In vitro. <i>Cancer Research</i> , 2007, 67, 332-338.	0.4	62
150	LAMP proteins are required for fusion of lysosomes with phagosomes. <i>EMBO Journal</i> , 2007, 26, 313-324.	3.5	542
151	Metalloproteases regulate T-cell proliferation and effector function via LAG-3. <i>EMBO Journal</i> , 2007, 26, 494-504.	3.5	203
152	Arrested maturation of Neisseria-containing phagosomes in the absence of the lysosome-associated membrane proteins, LAMP-1 and LAMP-2. <i>Cellular Microbiology</i> , 2007, 9, 2153-2166.	1.1	70
153	Cathepsin D Deficiency Is Associated with a Human Neurodegenerative Disorder. <i>American Journal of Human Genetics</i> , 2006, 78, 988-998.	2.6	255
154	Control of Peripheral Nerve Myelination by the β -Secretase BACE1. <i>Science</i> , 2006, 314, 664-666.	6.0	652
155	ADAM10 is a principal 'shedase' of the low-affinity immunoglobulin E receptor CD23. <i>Nature Immunology</i> , 2006, 7, 1293-1298.	7.0	189
156	Deafness in LIMP2-deficient mice due to early loss of the potassium channel KCNQ1/KCNE1 in marginal cells of the stria vascularis. <i>Journal of Physiology</i> , 2006, 576, 73-86.	1.3	54
157	LAMP-2 deficient mice show depressed cardiac contractile function without significant changes in calcium handling. <i>Basic Research in Cardiology</i> , 2006, 101, 281-291.	2.5	49
158	Multivariate neurocognitive and emotional profile of a mannosidosis murine model for therapy assessment. <i>Neurobiology of Disease</i> , 2006, 23, 422-432.	2.1	11
159	Breaking up the tie: Disintegrin-like metalloproteinases as regulators of cell migration in inflammation and invasion. , 2006, 111, 985-1006.		115
160	Furin-, ADAM 10-, and β -Secretase-Mediated Cleavage of a Receptor Tyrosine Phosphatase and Regulation of β -Catenin's Transcriptional Activity. <i>Molecular and Cellular Biology</i> , 2006, 26, 3917-3934.	1.1	102
161	Regulated ADAM10-dependent Ectodomain Shedding of β -Protocadherin C3 Modulates Cell-Cell Adhesion. <i>Journal of Biological Chemistry</i> , 2006, 281, 21735-21744.	1.6	94
162	Proteomic Identification of Desmoglein 2 and Activated Leukocyte Cell Adhesion Molecule as Substrates of ADAM17 and ADAM10 by Difference Gel Electrophoresis. <i>Molecular and Cellular Biology</i> , 2006, 26, 5086-5095.	1.1	106

#	ARTICLE	IF	CITATIONS
163	ADAM10 cleavage of N-cadherin and regulation of cell-cell adhesion and β -catenin nuclear signalling. <i>EMBO Journal</i> , 2005, 24, 742-752.	3.5	438
164	The NH ₂ -Terminal Transmembrane and Luminal Domains of LGP85 are Needed for the Formation of Enlarged Endosomes/Lysosomes. <i>Traffic</i> , 2005, 6, 895-906.	1.3	21
165	Mannose 6-phosphate receptors, Niemann-Pick C2 protein, and lysosomal cholesterol accumulation. <i>Journal of Lipid Research</i> , 2005, 46, 2559-2569.	2.0	52
166	β Subunits of Voltage-gated Sodium Channels Are Novel Substrates of β -Site Amyloid Precursor Protein-cleaving Enzyme (BACE1) and γ -Secretase. <i>Journal of Biological Chemistry</i> , 2005, 280, 23009-23017.	1.6	260
167	The apoptosis/autophagy paradox: autophagic vacuolization before apoptotic death. <i>Journal of Cell Science</i> , 2005, 118, 3091-3102.	1.2	487
168	Neurocognitive and Psychotiform Behavioral Alterations and Enhanced Hippocampal Long-Term Potentiation in Transgenic Mice Displaying Neuropathological Features of Human α -Mannosidosis. <i>Journal of Neuroscience</i> , 2005, 25, 6539-6549.	1.7	62
169	ADAM10 mediates E-cadherin shedding and regulates epithelial cell-cell adhesion, migration, and β -catenin translocation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 9182-9187.	3.3	604
170	Soluble Axl Is Generated by ADAM10-Dependent Cleavage and Associates with Gas6 in Mouse Serum. <i>Molecular and Cellular Biology</i> , 2005, 25, 9324-9339.	1.1	70
171	L1 Is Sequentially Processed by Two Differently Activated Metalloproteases and Presenilin/ γ -Secretase and Regulates Neural Cell Adhesion, Cell Migration, and Neurite Outgrowth. <i>Molecular and Cellular Biology</i> , 2005, 25, 9040-9053.	1.1	212
172	Phenotypic and Biochemical Analyses of BACE1- and BACE2-deficient Mice. <i>Journal of Biological Chemistry</i> , 2005, 280, 30797-30806.	1.6	309
173	Participation of Autophagy in Storage of Lysosomes in Neurons from Mouse Models of Neuronal Ceroid-Lipofuscinoses (Batten Disease). <i>American Journal of Pathology</i> , 2005, 167, 1713-1728.	1.9	305
174	Disturbed Cholesterol Traffic but Normal Proteolytic Function in LAMP-1/LAMP-2 Double-deficient Fibroblasts. <i>Molecular Biology of the Cell</i> , 2004, 15, 3132-3145.	0.9	241
175	Natural Soluble Interleukin-15 Is Generated by Cleavage That Involves the Tumor Necrosis Factor- α -converting Enzyme (TACE/ADAM17). <i>Journal of Biological Chemistry</i> , 2004, 279, 40368-40375.	1.6	65
176	Efficacy of enzyme replacement therapy in α -mannosidosis mice: a preclinical animal study. <i>Human Molecular Genetics</i> , 2004, 13, 1979-1988.	1.4	87
177	Partial loss of presenilins causes seborrheic keratosis and autoimmune disease in mice. <i>Human Molecular Genetics</i> , 2004, 13, 1321-1331.	1.4	76
178	The Transmembrane CXC-Chemokine Ligand 16 Is Induced by IFN- γ and TNF- α and Shed by the Activity of the Disintegrin-Like Metalloproteinase ADAM10. <i>Journal of Immunology</i> , 2004, 172, 6362-6372.	0.4	369
179	Distinct roles for ADAM10 and ADAM17 in ectodomain shedding of six EGFR ligands. <i>Journal of Cell Biology</i> , 2004, 164, 769-779.	2.3	895
180	Role for Rab7 in maturation of late autophagic vacuoles. <i>Journal of Cell Science</i> , 2004, 117, 4837-4848.	1.2	781

#	ARTICLE	IF	CITATIONS
181	Evidence for a Critical Role of the Tumor Necrosis Factor α Convertase (TACE) in Ectodomain Shedding of the p75 Neurotrophin Receptor (p75NTR). <i>Journal of Biological Chemistry</i> , 2004, 279, 4241-4249.	1.6	134
182	Cell-matrix interaction via CD44 is independently regulated by different metalloproteinases activated in response to extracellular Ca ²⁺ influx and PKC activation. <i>Journal of Cell Biology</i> , 2004, 165, 893-902.	2.3	260
183	At the acidic edge: emerging functions for lysosomal membrane proteins. <i>Trends in Cell Biology</i> , 2003, 13, 137-145.	3.6	564
184	Involvement of two different cell death pathways in retinal atrophy of cathepsin D-deficient mice. <i>Molecular and Cellular Neurosciences</i> , 2003, 22, 146-161.	1.0	142
185	The Cell Adhesion Protein P-selectin Glycoprotein Ligand-1 Is a Substrate for the Aspartyl Protease BACE1. <i>Journal of Biological Chemistry</i> , 2003, 278, 48713-48719.	1.6	230
186	Cellular Cholesterol Depletion Triggers Shedding of the Human Interleukin-6 Receptor by ADAM10 and ADAM17 (TACE). <i>Journal of Biological Chemistry</i> , 2003, 278, 38829-38839.	1.6	332
187	LIMP-2/LGP85 deficiency causes ureteric pelvic junction obstruction, deafness and peripheral neuropathy in mice. <i>Human Molecular Genetics</i> , 2003, 12, 631-646.	1.4	110
188	Thyroid functions of mouse cathepsins B, K, and L. <i>Journal of Clinical Investigation</i> , 2003, 111, 1733-1745.	3.9	188
189	The disintegrin-like metalloproteinase ADAM10 is involved in constitutive cleavage of CX3CL1 (fractalkine) and regulates CX3CL1-mediated cell-cell adhesion. <i>Blood</i> , 2003, 102, 1186-1195.	0.6	624
190	LIMP-2/LGP85 deficiency causes ureteric pelvic junction obstruction, deafness and peripheral neuropathy in mice. <i>Human Molecular Genetics</i> , 2003, 12, 631-46.	1.4	49
191	The disintegrin/metalloprotease ADAM 10 is essential for Notch signalling but not for alpha-secretase activity in fibroblasts. <i>Human Molecular Genetics</i> , 2002, 11, 2615-2624.	1.4	580
192	A role for the lysosomal membrane protein LGP85 in the biogenesis and maintenance of endosomal and lysosomal morphology. <i>Journal of Cell Science</i> , 2002, 115, 4117-4131.	1.2	132
193	Role of LAMP-2 in Lysosome Biogenesis and Autophagy. <i>Molecular Biology of the Cell</i> , 2002, 13, 3355-3368.	0.9	309
194	Disease model: LAMP-2 enlightens Danon disease. <i>Trends in Molecular Medicine</i> , 2001, 7, 37-39.	3.5	98
195	Involvement of Nitric Oxide Released from Microglia-Macrophages in Pathological Changes of Cathepsin D-Deficient Mice. <i>Journal of Neuroscience</i> , 2001, 21, 7526-7533.	1.7	113
196	The Disintegrins ADAM10 and TACE Contribute to the Constitutive and Phorbol Ester-regulated Normal Cleavage of the Cellular Prion Protein. <i>Journal of Biological Chemistry</i> , 2001, 276, 37743-37746.	1.6	222
197	The discrepancy between presenilin subcellular localization and β -secretase processing of amyloid precursor protein. <i>Journal of Cell Biology</i> , 2001, 154, 731-740.	2.3	155
198	Accumulation of autophagic vacuoles and cardiomyopathy in LAMP-2-deficient mice. <i>Nature</i> , 2000, 406, 902-906.	13.7	836

#	ARTICLE	IF	CITATIONS
199	Cathepsin D Deficiency Induces Lysosomal Storage with Ceroid Lipofuscin in Mouse CNS Neurons. <i>Journal of Neuroscience</i> , 2000, 20, 6898-6906.	1.7	353
200	Cathepsin L deficiency as molecular defect offurless:hyperproliferation of keratinocytes and pertubation of hair follicle cycling. <i>FASEB Journal</i> , 2000, 14, 2075-2086.	0.2	290
201	1/4 1A-adaptin-deficient mice: lethality, loss of AP-1 binding and rerouting of mannose 6-phosphate receptors. <i>EMBO Journal</i> , 2000, 19, 2193-2203.	3.5	388
202	Cell Surface Presenilin-1 Participates in the β -Secretase-like Proteolysis of Notch. <i>Journal of Biological Chemistry</i> , 1999, 274, 36801-36807.	1.6	246
203	A presenilin-1-dependent β -secretase-like protease mediates release of Notch intracellular domain. <i>Nature</i> , 1999, 398, 518-522.	13.7	2,002
204	Presenilin-1 deficiency leads to loss of Cajalâ€Retzius neurons and cortical dysplasia similar to human type 2 lissencephaly. <i>Current Biology</i> , 1999, 9, 719-727.	1.8	132
205	Deficiency of presenilin-1 inhibits the normal cleavage of amyloid precursor protein. <i>Nature</i> , 1998, 391, 387-390.	13.7	1,765