

Shuheï Nakanishi

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8445431/shuheï-nakanishi-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

54
papers

246
citations

9
h-index

14
g-index

59
ext. papers

365
ext. citations

3.9
avg, IF

2.96
L-index

#	Paper	IF	Citations
54	Vitamin D deficiency osteomalacia triggered by long-term social withdrawal and unbalanced diet in a Japanese middle-aged subject: A case report.. <i>Medicine (United States)</i> , 2022 , 101, e28589	1.8	0
53	Glucagon Test Is a Useful Predictor of Withdrawal From Insulin Therapy in Subjects With Type 2 Diabetes Mellitus.. <i>Frontiers in Endocrinology</i> , 2022 , 13, 871660	5.7	0
52	Case Report: A Variety of Immune-Related Adverse Events Triggered by Immune Checkpoint Inhibitors in a Subject With Malignant Melanoma: Destructive Thyroiditis, Aseptic Meningitis and Isolated ACTH Deficiency. <i>Frontiers in Endocrinology</i> , 2021 , 12, 722586	5.7	1
51	Case Report: A Case of Pituitary Adenoma Producing Growth Hormone and Thyroid-Stimulating Hormone Simultaneously. <i>Frontiers in Endocrinology</i> , 2021 , 12, 659076	5.7	0
50	Effects of sedentary behavior and daily walking steps on body mass index and body composition: Prospective observational study using outpatient clinical data of Japanese patients with type 2 diabetes. <i>Journal of Diabetes Investigation</i> , 2021 , 12, 1732-1738	3.9	0
49	Acute and extremely severe necrotic esophagitis accompanied by hyperglycemic hyperosmolar syndrome in a subject with type 2 diabetes mellitus. <i>Journal of Diabetes Investigation</i> , 2021 , 12, 1925-1926	3.9	1
48	Clinical Characteristics of Withdrawal of Basal Insulin Therapy Among Japanese Patients with Type 2 Diabetes: A Multicenter Retrospective Observational Study. <i>Diabetes Therapy</i> , 2021 , 12, 1849-1860	3.6	0
47	Case Report: Various Abnormalities in Lipid and Glucose Metabolism Induced by Capecitabine. <i>Frontiers in Oncology</i> , 2021 , 11, 664475	5.3	2
46	Case Report: Malignant Pheochromocytoma Without Hypertension Accompanied by Increment of Serum VEGF Level and Catecholamine Cardiomyopathy. <i>Frontiers in Endocrinology</i> , 2021 , 12, 688536	5.7	2
45	A case of tamoxifen-induced hypertriglyceridemia monitoring the changes in lipoprotein fractions over time. <i>BMC Endocrine Disorders</i> , 2021 , 21, 115	3.3	1
44	Significance of body mass index for diagnosing sarcopenia is equivalent to slow gait speed in Japanese individuals with type 2 diabetes: Cross-sectional study using outpatient clinical data. <i>Journal of Diabetes Investigation</i> , 2021 , 12, 417-424	3.9	4
43	Dulaglutide exerts beneficial anti atherosclerotic effects in ApoE knockout mice with diabetes: the earlier, the better. <i>Scientific Reports</i> , 2021 , 11, 1425	4.9	6
42	Persistent Hypoglycemia Induced by Long-acting Insulin Degludec. <i>Internal Medicine</i> , 2021 ,	1.1	1
41	Comprehensive efficacy of ipragliflozin on various conditioned type 2 diabetes compared with dipeptidyl peptidase-4 inhibitors and with both agents, based on a real-world multicenter trial. <i>Diabetology International</i> , 2021 , 12, 364-378	2.3	0
40	Favorable Effects of GLP-1 Receptor Agonist against Pancreatic β Cell Glucose Toxicity and the Development of Arteriosclerosis: "The Earlier, the Better" in Therapy with Incretin-Based Medicine. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	1
39	Concurrence of overt Cushing's syndrome and primary aldosteronism accompanied by aldosterone-producing cell cluster in adjacent adrenal cortex: case report. <i>BMC Endocrine Disorders</i> , 2021 , 21, 163	3.3	0
38	Switching From Daily DPP-4 Inhibitor to Once-Weekly GLP-1 Receptor Activator Dulaglutide Significantly Ameliorates Glycemic Control in Subjects With Poorly Controlled Type 2 Diabetes Mellitus: A Retrospective Observational Study. <i>Frontiers in Endocrinology</i> , 2021 , 12, 714447	5.7	0

37	Early combination therapy of empagliflozin and linagliptin exerts beneficial effects on pancreatic β cells in diabetic db/db mice. <i>Scientific Reports</i> , 2021 , 11, 16120	4.9	1
36	Multiple endocrine neoplasia type 1 with a frameshift mutation in its gene accompanied by a giant cervical lipoma and multiple fatty deposits in the pancreas: Case report. <i>BMC Endocrine Disorders</i> , 2021 , 21, 164	3.3	0
35	Idiopathic Bilateral Extraocular Myositis in a Subject With Poorly Controlled Type 2 Diabetes Mellitus: Case Report. <i>Frontiers in Medicine</i> , 2021 , 8, 700307	4.9	1
34	Syndrome of inappropriate secretion of thyroid-stimulating hormone in a subject with galactorrhea and menstrual disorder and undergoing infertility treatment: Case report.. <i>Medicine (United States)</i> , 2021 , 100, e28414	1.8	
33	Association Between Severity of Diabetic Neuropathy and Success in Weight Loss During Hospitalization Among Japanese Patients with Type 2 Diabetes: A Retrospective Observational Study. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2020 , 13, 1669-1676	3.4	
32	Insulin allergy brought out 8 years after starting insulin therapy in a subject with type 1 diabetes mellitus. <i>Acta Diabetologica</i> , 2020 , 57, 1025-1026	3.9	3
31	Comparison of the effects of three kinds of glucose-lowering drugs on non-alcoholic fatty liver disease in patients with type 2 diabetes: A randomized, open-label, three-arm, active control study. <i>Journal of Diabetes Investigation</i> , 2020 , 11, 1612-1622	3.9	21
30	Effect of Combination Therapy of Canagliflozin Added to Tenzeligliptin Monotherapy in Japanese Subjects with Type 2 Diabetes Mellitus: A Retrospective Study. <i>Journal of Diabetes Research</i> , 2020 , 2020, 4861681	3.9	2
29	Impact of physical activity and sedentary time on glycated hemoglobin levels and body composition: Cross-sectional study using outpatient clinical data of Japanese patients with type 2 diabetes. <i>Journal of Diabetes Investigation</i> , 2020 , 11, 633-639	3.9	2
28	Impact of sarcopenia on glycemic control and atherosclerosis in Japanese patients with type 2 diabetes: Cross-sectional study using outpatient clinical data. <i>Geriatrics and Gerontology International</i> , 2020 , 20, 1196-1201	2.9	2
27	Basedow's disease with associated features of Hashimoto's thyroiditis based on histopathological findings. <i>BMC Endocrine Disorders</i> , 2020 , 20, 120	3.3	2
26	Severe hypertriglyceridemia in a subject with disturbed life style and poor glycemic control without recurrence of acute pancreatitis: a case report. <i>BMC Endocrine Disorders</i> , 2019 , 19, 92	3.3	2
25	Vascular endothelial PDK1 plays a pivotal role in the maintenance of pancreatic beta cell mass and function in adult male mice. <i>Diabetologia</i> , 2019 , 62, 1225-1236	10.3	6
24	Idiopathic and isolated adrenocorticotrophic hormone deficiency presenting as continuous epigastric discomfort without symptoms of hypoglycemia: a case report. <i>Journal of Medical Case Reports</i> , 2019 , 13, 113	1.2	
23	Eicosapentaenoic acid/arachidonic acid ratio and weight loss during hospitalization for glycemic control among overweight Japanese patients with type 2 diabetes: a retrospective observational study. <i>Lipids in Health and Disease</i> , 2019 , 18, 36	4.4	0
22	Verification of Kumamoto Declaration 2013 and Glycemic Targets for Elderly Patients with Diabetes in Japan for prevention of diabetic complications: A retrospective longitudinal study using outpatient clinical data. <i>Journal of Diabetes Investigation</i> , 2019 , 10, 290-301	3.9	3
21	Comparison of HbA1c levels and body mass index for prevention of diabetic kidney disease: A retrospective longitudinal study using outpatient clinical data in Japanese patients with type 2 diabetes mellitus. <i>Diabetes Research and Clinical Practice</i> , 2019 , 155, 107807	7.4	6
20	Effect of mild exercise on glycemic and bodyweight control in Japanese type 2 diabetes patients: A retrospective analysis. <i>Journal of Diabetes Investigation</i> , 2019 , 10, 104-107	3.9	1

19	Body composition and development of diabetes: a 15-year follow-up study in a Japanese population. <i>European Journal of Clinical Nutrition</i> , 2018 , 72, 374-380	5.2	17
18	There is a Close Association Between the Recovery of Liver Injury and Glycemic Control after SGLT2 Inhibitor Treatment in Japanese Subjects with Type 2 Diabetes: A Retrospective Clinical Study. <i>Diabetes Therapy</i> , 2018 , 9, 1569-1580	3.6	6
17	Protective effects of the SGLT2 inhibitor luseogliflozin on pancreatic β cells in db/db mice: The earlier and longer, the better. <i>Diabetes, Obesity and Metabolism</i> , 2018 , 20, 2442-2457	6.7	20
16	Onset of type 1 diabetes mellitus and heparin-induced thrombocytopenia in a patient with Basedow's disease and idiopathic thrombocytopenic purpura: Novel combination as autoimmune polyglandular syndrome. <i>Journal of Diabetes Investigation</i> , 2018 , 9, 1381-1382	3.9	0
15	Switching from low-dose thiazide diuretics to sodium-glucose cotransporter 2 inhibitor improves various metabolic parameters without affecting blood pressure in patients with type 2 diabetes and hypertension. <i>Journal of Diabetes Investigation</i> , 2018 , 9, 875-881	3.9	9
14	Administration of RAS Inhibitor before the Onset of Diabetic Nephropathy Counteracts the Adverse Effect of Chronic Hyperglycemia and Reduces the Augmentation of Urinary Albumin Excretion: A Retrospective Clinical Study. <i>Journal of Diabetes Research</i> , 2018 , 2018, 9435401	3.9	1
13	Efficacy and Safety of Switching from Insulin Glargine 100 U/mL to the Same Dose of Glargine 300 U/mL in Japanese Type 1 and 2 Diabetes Patients: A Retrospective Analysis. <i>Internal Medicine</i> , 2018 , 57, 1381-1389	1.1	2
12	Effect of Tofogliflozin on Body Composition and Glycemic Control in Japanese Subjects with Type 2 Diabetes Mellitus. <i>Journal of Diabetes Research</i> , 2018 , 2018, 6470137	3.9	18
11	Advanced breast cancer in a relatively young man with severe obesity and type 2 diabetes mellitus. <i>Journal of Diabetes Investigation</i> , 2017 , 8, 395-396	3.9	7
10	Werner Syndrome and Diabetes Mellitus Accompanied by Adrenal Cortex Cancer. <i>Internal Medicine</i> , 2017 , 56, 1987-1992	1.1	3
9	Decreased glucagon-like peptide 1 receptor expression in endothelial and smooth muscle cells in diabetic db/db mice: TCF7L2 is a possible regulator of the vascular glucagon-like peptide 1 receptor. <i>Diabetes and Vascular Disease Research</i> , 2017 , 14, 540-548	3.3	11
8	Temporal lobe epilepsy associated with GAD autoimmunity. <i>Acta Diabetologica</i> , 2017 , 54, 321-323	3.9	
7	Efficacy and safety of insulin degludec in Japanese patients with type 1 and type 2 diabetes: 24-week results from the observational study in routine clinical practice. <i>Journal of Diabetes Investigation</i> , 2016 , 7, 94-9	3.9	27
6	Sex Differences in the Association between the Eicosapentaenoic Acid/Arachidonic Acid Ratio and the Visceral Fat Area among Patients with Type 2 Diabetes. <i>Internal Medicine</i> , 2016 , 55, 1269-74	1.1	2
5	Association of GA/HbA1c ratio and cognitive impairment in subjects with type 2 diabetes mellitus. <i>Journal of Diabetes and Its Complications</i> , 2016 , 30, 1452-1455	3.2	16
4	Effects of microvascular complications on structural changes in visceral and subcutaneous adiposity among type 2 diabetes patients. <i>Obesity Medicine</i> , 2016 , 1, 1-5	2.6	1
3	Strawberry milk-like blood in a subject with diabetic lipemia: dramatic change to transparent color after insulin therapy. <i>SpringerPlus</i> , 2016 , 5, 1499		2
2	Regardless of central obesity, metabolic syndrome is a significant predictor of type 2 diabetes in Japanese Americans. <i>Journal of Diabetes Investigation</i> , 2015 , 6, 527-32	3.9	4

- 1 Prevalence of metabolic syndrome compared between native Japanese and Japanese-Americans. *Diabetes Research and Clinical Practice*, **2008**, 79, 518-22 7.4 29