Naho Ichikawa

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8454232/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Harmonization of resting-state functional MRI data across multiple imaging sites via the separation of site differences into sampling bias and measurement bias. PLoS Biology, 2019, 17, e3000042.	5.6	127
2	A prediction model of working memory across health and psychiatric disease using whole-brain functional connectivity. ELife, 2018, 7, .	6.0	73
3	Global connectivity and local excitability changes underlie antidepressant effects of repetitive transcranial magnetic stimulation. Neuropsychopharmacology, 2020, 45, 1018-1025.	5.4	71
4	Patients with major depressive disorder exhibit reduced reward size coding in the striatum. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2017, 79, 317-323.	4.8	62
5	Generalizable brain network markers of major depressive disorder across multiple imaging sites. PLoS Biology, 2020, 18, e3000966.	5.6	54
6	A multi-site, multi-disorder resting-state magnetic resonance image database. Scientific Data, 2021, 8, 227.	5.3	48
7	Primary functional brain connections associated with melancholic major depressive disorder and modulation by antidepressants. Scientific Reports, 2020, 10, 3542.	3.3	39
8	Overlapping but Asymmetrical Relationships Between Schizophrenia and Autism Revealed by Brain Connectivity. Schizophrenia Bulletin, 2020, 46, 1210-1218.	4.3	28
9	Effects of behavioral activation on default mode network connectivity in subthreshold depression: A preliminary resting-state fMRI study. Journal of Affective Disorders, 2018, 227, 156-163.	4.1	20
10	Enhancing Multi-Center Generalization of Machine Learning-Based Depression Diagnosis From Resting-State fMRI. Frontiers in Psychiatry, 2020, 11, 400.	2.6	20
11	Functional Alterations of Postcentral Gyrus Modulated by Angry Facial Expressions during Intraoral Tactile Stimuli in Patients with Burning Mouth Syndrome: A Functional Magnetic Resonance Imaging Study. Frontiers in Psychiatry, 2017, 8, 224.	2.6	16
12	Regional brain functions in the resting state indicative of potential differences between depression and chronic pain. Scientific Reports, 2017, 7, 3003.	3.3	13
13	Importance of the Habenula for Avoidance Learning Including Contextual Cues in the Human Brain: A Preliminary fMRI Study. Frontiers in Human Neuroscience, 2020, 14, 165.	2.0	9
14	Effects of behavioural activation on the neural circuit related to intrinsic motivation. BJPsych Open, 2018, 4, 317-323.	0.7	8
15	Neural and sympathetic activity associated with exploration in decision-making: further evidence for involvement of insula. Frontiers in Behavioral Neuroscience, 2014, 8, 381.	2.0	7
16	Predicting Ventral Striatal Activation During Reward Anticipation From Functional Connectivity at Rest. Frontiers in Human Neuroscience, 2019, 13, 289.	2.0	5
17	Common Brain Networks Between Major Depressive-Disorder Diagnosis and Symptoms of Depression That Are Validated for Independent Cohorts. Frontiers in Psychiatry, 2021, 12, 667881.	2.6	3
18	Generalizable brain network markers of major depressive disorder across multiple imaging sites. , 2020, 18, e3000966.		0

#	Article	IF	CITATIONS
19	Generalizable brain network markers of major depressive disorder across multiple imaging sites. , 2020, 18, e3000966.		0
20	Generalizable brain network markers of major depressive disorder across multiple imaging sites. , 2020, 18, e3000966.		0
21	Generalizable brain network markers of major depressive disorder across multiple imaging sites. , 2020, 18, e3000966.		0
22	Generalizable brain network markers of major depressive disorder across multiple imaging sites. , 2020, 18, e3000966.		0
23	Generalizable brain network markers of major depressive disorder across multiple imaging sites. , 2020, 18, e3000966.		0