Naho Ichikawa

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8454232/publications.pdf

Version: 2024-02-01

759233 839539 23 612 12 18 h-index citations g-index papers 25 25 25 943 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Common Brain Networks Between Major Depressive-Disorder Diagnosis and Symptoms of Depression That Are Validated for Independent Cohorts. Frontiers in Psychiatry, 2021, 12, 667881.	2.6	3
2	A multi-site, multi-disorder resting-state magnetic resonance image database. Scientific Data, 2021, 8, 227.	5. 3	48
3	Importance of the Habenula for Avoidance Learning Including Contextual Cues in the Human Brain: A Preliminary fMRI Study. Frontiers in Human Neuroscience, 2020, 14, 165.	2.0	9
4	Enhancing Multi-Center Generalization of Machine Learning-Based Depression Diagnosis From Resting-State fMRI. Frontiers in Psychiatry, 2020, 11, 400.	2.6	20
5	Primary functional brain connections associated with melancholic major depressive disorder and modulation by antidepressants. Scientific Reports, 2020, 10, 3542.	3.3	39
6	Global connectivity and local excitability changes underlie antidepressant effects of repetitive transcranial magnetic stimulation. Neuropsychopharmacology, 2020, 45, 1018-1025.	5 . 4	71
7	Overlapping but Asymmetrical Relationships Between Schizophrenia and Autism Revealed by Brain Connectivity. Schizophrenia Bulletin, 2020, 46, 1210-1218.	4.3	28
8	Generalizable brain network markers of major depressive disorder across multiple imaging sites. PLoS Biology, 2020, 18, e3000966.	5 . 6	54
9	Generalizable brain network markers of major depressive disorder across multiple imaging sites. , 2020, 18, e3000966.		O
10	Generalizable brain network markers of major depressive disorder across multiple imaging sites. , 2020, 18, e3000966.		0
11	Generalizable brain network markers of major depressive disorder across multiple imaging sites. , 2020, 18, e3000966.		O
12	Generalizable brain network markers of major depressive disorder across multiple imaging sites. , 2020, 18, e3000966.		0
13	Generalizable brain network markers of major depressive disorder across multiple imaging sites. , 2020, 18, e3000966.		O
14	Generalizable brain network markers of major depressive disorder across multiple imaging sites. , 2020, 18, e3000966.		0
15	Predicting Ventral Striatal Activation During Reward Anticipation From Functional Connectivity at Rest. Frontiers in Human Neuroscience, 2019, 13, 289.	2.0	5
16	Harmonization of resting-state functional MRI data across multiple imaging sites via the separation of site differences into sampling bias and measurement bias. PLoS Biology, 2019, 17, e3000042.	5.6	127
17	Effects of behavioral activation on default mode network connectivity in subthreshold depression: A preliminary resting-state fMRI study. Journal of Affective Disorders, 2018, 227, 156-163.	4.1	20
18	Effects of behavioural activation on the neural circuit related to intrinsic motivation. BJPsych Open, 2018, 4, 317-323.	0.7	8

#	Article	IF	CITATION
19	A prediction model of working memory across health and psychiatric disease using whole-brain functional connectivity. ELife, $2018, 7, \ldots$	6.0	73
20	Regional brain functions in the resting state indicative of potential differences between depression and chronic pain. Scientific Reports, 2017, 7, 3003.	3.3	13
21	Patients with major depressive disorder exhibit reduced reward size coding in the striatum. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2017, 79, 317-323.	4.8	62
22	Functional Alterations of Postcentral Gyrus Modulated by Angry Facial Expressions during Intraoral Tactile Stimuli in Patients with Burning Mouth Syndrome: A Functional Magnetic Resonance Imaging Study. Frontiers in Psychiatry, 2017, 8, 224.	2.6	16
23	Neural and sympathetic activity associated with exploration in decision-making: further evidence for involvement of insula. Frontiers in Behavioral Neuroscience, 2014, 8, 381.	2.0	7